

Fall 2019

Topology

WVU Mathematics Department

Follow this and additional works at: <https://researchrepository.wvu.edu/math-grad-exams>

Recommended Citation

WVU Mathematics Department, "Topology" (2019). *M.S. Advanced and Ph.D. Entrance Exams*. 9.
<https://researchrepository.wvu.edu/math-grad-exams/9>

This Other is brought to you for free and open access by the Mathematics at The Research Repository @ WVU. It has been accepted for inclusion in M.S. Advanced and Ph.D. Entrance Exams by an authorized administrator of The Research Repository @ WVU. For more information, please contact researchrepository@mail.wvu.edu.

#	1	2	3	4	5	total
score						

Directions: Solve all of the following problems. Start each solution on a **new sheet of paper**, write the problem number and page number (of the particular problem). The pages should be numbered **separately for each problem** with the first page of each problem having number 1. Write the solution on **one side** of the paper and stay **within the borders**. Anything written outside the borders will not be taken into account. Return your solutions with pages in correct order **arranged according to problem numbers** and together with this cover page.

- (1) Prove, or give a counterexample, to each of the following statements.
 - (a) Continuous image of a separable space is separable. (Include definition of a separable space.)
 - (b) Continuous image of a Lindelöf space is Lindelöf. (Include definition of a Lindelöf space.)
 - (c) Continuous image of a completely regular space is completely regular. (Include definition of a completely regular space.)
- (2) Let $\langle X, d \rangle$ and $\langle Y, \rho \rangle$ be compact metric spaces and let $f : X \rightarrow Y$ and $g : X \rightarrow Y$ be continuous. Show that if $f(x) \neq g(x)$ for all $x \in X$, then there exists a $\varepsilon > 0$ such that $\rho(f(x), g(x)) > \varepsilon$.
- (3) Let $\{T_a\}_{a \in A}$ be a family of connected subsets of a topological space X . Let $Y := \bigcup_{a \in A} T_a$ and assume that for every $x, y \in Y$ there is a finite sequence $a_0, \dots, a_n \in A$ such that $x \in T_{a_0}$, $y \in T_{a_n}$, and $T_{a_i} \cap T_{a_{i+1}} \neq \emptyset$ that for every $i < n$. Prove that Y is connected.
- (4) Let X be a regular Hausdorff space and A be an infinite subset of X . Prove that there is a sequence U_1, U_2, \dots of open subsets of X such that $\overline{U_n} \cap \overline{U_m} = \emptyset$ when $m \neq n$ and $U_n \cap A \neq \emptyset$ for each $n \in \mathbb{N}$.
Hint: Construct the sequence $\langle U_n : n \in \mathbb{N} \rangle$ by induction so that $A \setminus \overline{U_n}$ is infinite for every n .
- (5) Let $X = \mathbb{N}^{\mathbb{N}}$ have the product topology assuming that \mathbb{N} has discrete topology. Let $Y \subseteq X$ be a compact subspace of X .
 - (a) Prove that for each $n \in \mathbb{N}$, there is finite $F_n \subseteq \mathbb{N}$ such that $Y \subseteq \prod_{n \in \mathbb{N}} F_n$.
 - (b) Show that X is not a countable union of compact subsets of X .
Hint: In the proof of part (b) use part (a) and a diagonal argument to show that for every sequence $\langle Y_n : n \in \mathbb{N} \rangle$ of compact subsets of X there is an $x \in X$ that belongs to no Y_n .

page #

of problem #

name:
