

Fall 2013

Differential Equations

WVU Mathematics Department

Follow this and additional works at: <https://researchrepository.wvu.edu/math-grad-exams>

Recommended Citation

WVU Mathematics Department, "Differential Equations" (2013). *M.S. Advanced and Ph.D. Entrance Exams*. 19.

<https://researchrepository.wvu.edu/math-grad-exams/19>

This Other is brought to you for free and open access by the Mathematics at The Research Repository @ WVU. It has been accepted for inclusion in M.S. Advanced and Ph.D. Entrance Exams by an authorized administrator of The Research Repository @ WVU. For more information, please contact researchrepository@mail.wvu.edu.

ENTRANCE EXAM: Differential Equations, Fall 2013.

Solve any 6 (six) problems. All problems carry the same weight.

1. Given the initial value problem

$$\begin{aligned} y' &= f(t, y, p), & y(t_0) &= \eta, \\ t, t_0, p &\in \mathbb{R}^1, & y, \eta, f &\in \mathbb{R}^n. \end{aligned}$$

Assume that $f(t, y, p)$ is a continuous vector function and that the entries of the Jacobian matrix

$$Jf := \left(\frac{\partial f_i}{\partial y_j} \right)_{ij} = \begin{bmatrix} \nabla f_1 \\ \nabla f_2 \\ \cdot \\ \cdot \\ \nabla f_n \end{bmatrix}$$

are continuous in the set of points (t, y, p) , where

$$t, t_0, p \in \mathbb{R}^1, \quad y, \eta, f \in \mathbb{R}^n.$$

Consider the solution $y = \phi(t, t_0, \eta, p)$ as a vector function of the variables (t, t_0, η, p) . Determine an equation that is satisfied by a) $\frac{\partial \phi}{\partial t_0}$, b) $\frac{\partial \phi}{\partial p}$. What information could be derived from these equations? Why is that information important?

2. Convert the scalar ode $x'' + x^7 = 0$ into a vector system of differential equations. Show, that a solution to any initial value problem of the ode exists on any interval $[a, b]$.

3. a) Explain the essence of the method of successive approximations and under what assumptions it is valid. b) Calculate the first 3 successive approximations for the initial value problem

$$y' = \begin{bmatrix} -y_1 + y_2^2 \\ y_1 + y_2 \end{bmatrix}, \quad y(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

4. Provide the best estimates for the maximal intervals of existence of solutions of

$$y' = \begin{bmatrix} (t+5)^{-2} & t \\ t^2 & (t-1)^{-3} \end{bmatrix} y, \quad y(t_0) = \eta,$$

with a) $t_0 = -10$, b) $t_0 = 0$, c) $t_0 = 2$.

5. (a) Find and classify all the equilibria of the system

$$\dot{x} = -3y + xy - 4, \quad \dot{y} = x^2 - xy.$$

(b) Sketch the phase diagram.

6. Consider the system

$$\dot{x} = x - xy, \quad \dot{y} = y + xy.$$

(a) Find the equilibria and study their stability.

(b) Find all the invariant lines.

(c) Find the null and vertical (possibly curvilinear) clines.

(d) Draw the phase diagram and use arrows to indicate the direction of increasing time.

7. Consider the system

$$\dot{x} = 2x + y - 2x^3 - 3xy^2, \quad \dot{y} = -2x + 4y - 2x^2y - 4y^3.$$

(a) (6 points) Find all critical points and discuss their stability. *Hint: First use the Lyapunov function $V(x, y) = 2x^2 + y^2$ to prove that if there are other critical points beside the origin, then they must lie on the unit circle.*

(b) (4 points) Are there any periodic solutions? If so, approximately where are their orbits located?

8. Consider the linear system $\dot{\mathbf{x}} = A\mathbf{x}$, where A is an $n \times n$ nonsingular, antisymmetric (i.e. $A^T = -A$), real matrix.

(a) Show that every solution satisfies $|\mathbf{x}(t)| = \text{const}$ for all $t \in \mathbb{R}$.

(b) Find all critical points and study their stability.

(c) If $n = 2$, are there any non-periodic solutions?