1-1-1936

Mass feeding of sheep with copper sulphate and salt to control gastro-intestinal parasites

J. H. Rietz

Follow this and additional works at: https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins

Digital Commons Citation

Rietz, J. H., "Mass feeding of sheep with copper sulphate and salt to control gastro-intestinal parasites" (1936). *West Virginia Agricultural and Forestry Experiment Station Bulletins*. 271.
https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins/272

This Bulletin is brought to you for free and open access by the Davis College of Agriculture, Natural Resources And Design at The Research Repository @ WVU. It has been accepted for inclusion in West Virginia Agricultural and Forestry Experiment Station Bulletins by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

West Virginia University Libraries

3 0802 100521224 7

MASS FEEDING OF SHEEP WITH COPPER SULPHATE AND SALT TO CONTROL GASTRO-INTESTINAL PARASITES

By J. H. RIETZ

AGRICULTURAL EXPERIMENT STATION
COLLEGE OF AGRICULTURE, WEST VIRGINIA UNIVERSITY
F. D. FROMME, Director
MORGANTOWN

AGRICULTURAL EXPERIMENT STATION STAFF

C. S. BOUCHER, Ph. D., President of the University

F. D. FROMME, Ph. D. Dean and Director

GERALD JENNY, M. S., Agricultural Editor.....JOHN C. JOHNSTON, Chief Clerk

AGRONOMY AND GENETICS

R. J. Garber, Ph. D.
Agronomist and Geneticist

W. H. Pierre, Ph. D.
Associate Agronomist

T. C. McIlvalne, Ph. D.*†
Associate Agronomist

G. G. Pohlman, Ph. D.
Assistant Agronomist

C. R. Burnham, Ph. D.
Assistant Geneticist

J. A. Rigney, M. S.
Assistant in Agronomy

ANIMAL HUSBANDRY

E. A. Livesay, D. Sc.
Animal Husbandman

C. V. Wilson, M. S.
Assistant Animal Husbandman

J. H. Longwell, M. A.
Assistant Animal Husbandman

R. H. Tuckwiller, B. S. Agr.*
Assistant Animal Husbandman

J. H. Rietz, D. V. M.
Associate Veterinarian

E. T. Wightman, M. S.
Assistant Poultry Husbandman

T. B. Clark, M. S.
Assistant Poultry Husbandman

CHEMISTRY

R. B. Dustman, Ph. D.
Agricultural Chemist

C. E. Weakley, Jr., M. A.
Assistant Chemist

A. H. VanLandingham, Ph. D.
Assistant Chemist

I. J. Duncan, M. S.
Assistant in Agricultural Chemistry

DAIRY HUSBANDRY

H. O. Henderson, Ph. D.
Dairy Husbandman

L. M. Thurston, Ph. D.
Associate Dairy Husbandman

G. A. Bowling, M. S.
Assistant Dairy Husbandman

R. A. Ackerman, M. S.‡
Assistant Dairy Husbandman

ENTOMOLOGY

L. M. Peairs, Ph. D.
Entomologist

W. E. Rumsey, B. S. Agr.
Assistant Entomologist

FARM ECONOMICS

W. W. Armentrout, Ph. D.
Farm Economist

F. D. Cornell, Jr., M. S.
Assistant Farm Economist

C. H. Bruce, B. S.
Assistant in Farm Economics

W. P. Gainer, B. S. Agr.
Assistant in Farm Economics

L. F. Herrmann, M. S.
Assistant in Farm Economics

HOME ECONOMICS

Rachel H. Colwell, M. A.
Home Economist

Hazel C. Cameron, M. A.
Research Specialist in Nutrition

HORTICULTURE

H. E. Knowlton, Ph. D.
Horticulturist

K. C. Westover, M. S.
Assistant Horticulturist

L. P. Batjer, Ph. D.
Assistant Horticulturist

E. N. McCubbin, M. A.
Assistant in Horticulture

A. P. Dye, M. S. Agr.
Assistant in Horticulture

W. H. Childs, M. S.
Assistant in Horticulture

PLANT PATHOLOGY

C. R. Orton, Ph. D.
Plant Pathologist

L. H. Leonian, Ph. D.
Mycologist

Anthony Berg, M. S.
Associate Plant Pathologist

F. J. Schneiderhan, Ph. D.††
Associate Plant Pathologist

*In cooperation with the U. S. Department of Agriculture, Washington, D. C.

†In charge of the Lakin Experiment Farm, Lakin, W. Va.

‡In charge of the Reymann Memorial Farms, Wardensville, W. Va.

††In charge of the University Experiment Farm, Kearneysville, W. Va.

Mass Feeding of Sheep With Copper Sulphate and Salt to Control Gastro-Intestinal Parasites

by J. H. RIETZ

THE QUESTION OF MASS FEEDING of sheep with a mixture of copper sulphate with salt or with grain to control gastro-intestinal parasites is raised frequently. Mass feeding of any anthelmintic that will control internal parasites would be the ideal method of control.

The object of this experiment was to find a method of administration of the anthelmintic easier than the regular system of drenching, which has been proved effective (Hutcheson (6), Bell (1), Nighbert (7), Hardy and Schmidt (5), Rietz (8)), and thereby to bring about a more general use of control measures.

PLAN OF THE EXPERIMENT

In this experiment 20 ewes were allowed access to a mixture of 1 pound of copper sulphate to 30 pounds of salt; 7 ewes were used as treated controls and drenched at intervals of 21 days with 3 ounces of a 1½% solution of copper sulphate; and 4 ewes were left as untreated controls.

The 20 sheep receiving the copper-sulphate—salt mixture were pastured in a field of approximately 12 acres. When the grass became inadequate to provide proper feed it was supplemented by grain or grain and hay. The grain used was a mixture of equal parts of shelled corn and oats. The hay was a fair-to-good grade of alfalfa.

The copper-sulphate—salt mixture was placed in a box in the barn for protection from moisture and was otherwise protected from contamination and waste by the sheep. The 20 sheep consumed approximately 10 pounds of the copper-sulphate—salt mixture per month or one-half pound each per month throughout the experiment. This is approximately the normal salt consumption for these sheep when allowed access to salt *ad libitum* (Evvard, J. M., *et al.* (4), Bell (2)).

The pasture over which these sheep grazed had been used continuously during the three previous years as quarters for sheep used in investigational work on parasites. Fifteen species of parasites were identified by autopsy and laboratory findings during these investigations (8). Consequently the range was considered heavily infested.

The 7 treated and 4 untreated controls were pastured together in an adjoining pasture of approximately 6 acres, with all the conditions approximating as nearly as possible those in the pasture used for the 20 sheep fed the copper-sulphate—salt mixture. Salt was fed to this group *ad libitum*, the consumption being approximately one-half pound for each sheep per month.

TABLE 1—Condition and count of nematode ova for sheep fed 1:30 and 1:50 mixtures of copper sulphate and salt

No.	Nematode ova per gram of feces June, 1933	Condition of sheep June, 1933	Nematode ova per gram of feces June, 1934	Condition of sheep June, 1934	Nematode ova per gram of feces Sept. 1934, or at time of autopsy	Condition of sheep Sept. 1934, or at time of autopsy	Remarks
1	0	Good	4000	Good	14000	Good	Died August, 1934
3	0	Good			0	Good	Died May, 1934 Cu So ₄ poisoning
4	0	Good	6000	Good	26000	Good	Died May, 1934 Cu So ₄ poisoning
5	0	Good	800	Good	21000	Fair	Died April, 1934
6	0	Good	0	Good	0	Good	Cu So ₄ poisoning
85	0	Good	0	Good	0	Good	
30	0	Good	0	Good	400	Fair	
45	0	Good	100	Good	100	Good	
55	0	Good	700	Good	2500	Fair	
70	500	Fair	1200	Poor	2000	Emaciated	Stolen from pasture
52	200	Good	2000	Good	13000	Poor	September, 1934
86	100	Good	200	Good	600	Poor	Died September, 1934
31	200	Good	1400	Poor	1400	Poor	
71	700	Good	1700	Fair	1400	Fair	
10	300	Good	400	Fair	900	Poor	Killed accidentally
11	0	Good	300	Good			September, 1934
20	800	Good	1700	Poor	800	Fair	Killed by dogs, June, 1934
48	200	Good	400	Good	700	Fair	
51	0	Good	400	Good	1500	Fair	
50	0	Good	0	Good	700	Fair	

A count of the nematode ova in the feces of each sheep in the three groups was made (Stoll (9)) at the beginning, at intervals during, and at the termination of the experiment.

RESULTS OF THE EXPERIMENT

The trials in which the mixture of 1 pound of copper sulphate to 30 pounds of salt was used began in June 1933 and ended in June 1934. Table 1 shows the condition of each sheep and the count of nematode ova at the beginning and end of this period. The average count of nematode ova per gram of feces was 150 in June 1933, the beginning of the experiment, and 1100 in June 1934, the end of the experiment.

On the morning of April 6, 1934, sheep no. 6 of the lot allowed free access to the copper-sulphate—salt mixture was found dead in the pasture. This animal was apparently normal when fed grain on the evening of April 5. The autopsy showed the carcass to be in a good state of nourishment. There was an ample layer of fat over the body and the omentum contained much fat. All the body tissues showed a marked icterus. The liver was enlarged, soft, and yellow; kidneys swollen, dark, and soft; bladder hemorrhagic, contents bloody. No other marked changes were noted except the general icterus (Boughton and Hardy (3)). The tentative diagnosis of poisoning from feeding an excessive amount of copper sulphate was made.

On the morning of May 10 sheep no. 3 was found dead in the pasture. No previous illness had been noted by the attendant. The autopsy showed the carcass to be in good condition; tissues markedly icteric; liver enlarged, soft, and yellow; kidneys swollen, dark, and soft. The pelvis of the kidneys contained a light blue precipitate. The bladder was congested and empty. The yellow color was the only change noted in the other organs.

Analysis* of the liver and kidney of sheep no. 3 showed a deposit of 140 milligrams of copper per kilogram of liver. Qualitative analysis of the blue precipitate from the kidney showed this to be a heavy deposit of copper. The diagnosis was poisoning from feeding an excessive amount of copper sulphate.

Table 2 shows the condition of each sheep and the count of nematode ova per gram of feces for the 7 treated control sheep at the beginning and end of the experiment. The average count of nematode ova per gram of feces was 350 in June 1933, the beginning of the experiment, and 85 in June 1934, the end of the experiment.

Table 3 shows the condition of each sheep and the count of nematode ova per gram of feces for the 4 untreated sheep at the beginning and end of the experiment. The average count of nematode ova per gram of feces was 75 for June 1933, the beginning of the experiment, and 3400 in June 1934, the end of the experiment.

*The analysis was made by Dr. R. B. Dustman, head of the department of agricultural chemistry, West Virginia Agricultural Experiment Station.

TABLE 2—Condition of and count of nematode ova for sheep treated with 1½% copper-sulphate solution (control)

No.	Nematode ova per gram of feces June, 1933	Condition of sheep June, 1933	Nematode ova per gram of feces June, 1934	Condition of sheep June, 1934	Nematode ova per gram of feces Sept. 1934 or at time of autopsy	Condition of sheep Sept. 1934 or at time of autopsy	Remarks
14	0	Good	0	Good	0	Good	
16	200	Good	100	Good	0	Good	Killed by dogs, June 1934
18	0	Good	0	Good	...		Killed by dogs, June 1934
33	0	Good	0	Good	...		
34	0	Good	0	Good	0	Good	
44	2100	Poor	300	Fair	...		Killed by dogs, June 1934
96	200	Good	200	Good	200	Good	

TABLE 3—Condition of and count of nematode ova for the untreated control sheep

No.	Nematode ova per gram of feces June, 1933	Condition of sheep June, 1933	Nematode ova per gram of feces June, 1934	Condition of sheep June, 1934	Nematode ova per gram of feces Sept. 1934 or at time of autopsy	Condition of sheep Sept. 1934 or at time of autopsy	Remarks
35	100	Good	10000	Poor	...		Killed by dogs, June 1934
41	0	Good	300	Good	500	Good	Killed for post-mortem, Sept. 1934
23	0	Good	3000	Poor	...		Killed by dogs, June 1934
100	100	Good	500	Fair	600	Fair	Killed for post-mortem, Sept. 1934

Because of the loss from poisoning of two sheep, nos. 6 and 3, from the lot receiving the 1:30 mixture of copper sulphate and salt, the proportion of copper sulphate to salt was reduced from 1:30 to 1:50 in June 1934. The condition of the sheep and the count of nematode ova per gram of feces for the three lots of sheep in this experiment will be found in Tables 1, 2, and 3 for June 1934, bearing data on the beginning of the feeding of the 1:50 copper-sulphate—salt mixture as well as the end of the feeding of the 1:30 mixture.

Table 1 shows the condition of the sheep receiving the copper-sulphate—salt mixture and the count of nematode ova per gram of feces for June 1934 and September 1934, the beginning and end of the feeding of the 1:50 mixture.

The average count of nematode ova per gram of feces for this lot was 1100 in June 1934 and 4500 in September 1934, the beginning and termination of the feeding experiment. During this three-month period the average count of nematode ova per gram of feces had increased from 1100 to 4500, and the general condition of the flock had decreased materially.

Three weeks after changing the copper-sulphate—salt mixture from 1:30 to 1:50, sheep no. 4 was found dead in the pasture. This sheep did not show symptoms of sickness on the previous day, when seen by the attendant. The autopsy of this sheep showed a general icterus; liver enlarged, soft, and yellow; kidneys swollen, dark, and soft; bladder hemorrhagic, and contents a dark-brown liquid. The diagnosis was poisoning from feeding an excessive amount of copper sulphate.

SUPPLEMENTARY EXPERIMENT

In September 1934 the mass feeding of the copper-sulphate—salt mixture was discontinued. The 12 ewes surviving in this experiment with their 11 spring lambs were divided into two lots. Lot 1, consisting of 5 ewes and 5 lambs, was treated with a 1½% solution of copper sulphate at intervals of 21 days, beginning September 26, 1934. The adults received 3 ounces each as a drench. The lambs received 2 ounces on September 26; 2½ ounces on October 17 and on November 7; and 3 ounces each from November 28 to the end of the experiment on February 20, 1935. Lot 2, consisting of 7 ewes and 6 lambs, was treated with a mixture of equal parts of a 1½% solution of copper sulphate and a 1½% solution of nicotine sulphate (Black-leaf 40) at intervals of 21 days beginning September 26, 1934. The adults received 3 ounces each as a drench. The lambs received 2 ounces each on September 26; 2½ ounces each on October 17 and on November 7; and 3 ounces each from November 28 to the end of the experiment on February 20, 1935.

Table 4 shows the condition of the sheep and the count of nematode ova per gram of feces for each sheep in Lot 1. These sheep received the 1½% solution of copper sulphate from September 26,

TABLE 4—Condition of sheep and count of nematode ova at the beginning and end of the treatment with the copper-sulphate solution

No.	Nematode ova per gram of feces September, 1934	Condition of sheep September, 1934	Nematode ova per gram of feces February, 1935	Condition of sheep February, 1935	Remarks
5	21000	Fair	0	Good	
10	900	Poor	0	Good	
20	800	Fair	0	Good	
31	1400	Poor	100	Good	
	700	Fair	100	Good	
217	1800	Poor	100	Good	
122	17000	Poor	0	Good	
123	17500	Poor	200	Good	
124	400	Good	200	Good	
130	1600	Poor	100	Good	

TABLE 5—Condition of sheep and count of nematode ova at the beginning and end of the treatment with the copper-sulphate—
8 nicotine-sulphate mixture

No.	Nematode ova per gram of feces September, 1934	Condition of sheep September, 1934	Nematode ova per gram of feces February, 1935	Condition of sheep February, 1935	Remarks
30	400	Fair	0	Good	
50	700	Fair	100	Good	
45	100	Good	0	Good	
51	1500	Fair	0	Good	
52	13500	Poor	100	Good	
85	0	Good	0	Good	
86	600	Poor	...		
125	11000	Poor	0	Good	
126	4200	Poor	0	Good	
127	13200	Poor	400	Good	
219	13400	Poor	200	Good	
220	1500	Fair	0	Good	
221	17100	Fair	400	Good	

1934, the beginning, to February 20, 1935, the end of the period of treatment. The average count of nematode ova per gram of feces was 6300 on September 26, 1934, and 80 on February 20, 1935, the termination of the experiment.

Table 5 shows the condition of the sheep and the count of nematode ova per gram of feces for each sheep in Lot 2. These sheep received a mixture of equal parts of a 1½% solution of copper sulphate and a 1½% solution of nicotine sulphate (Blackleaf 40) for September 26, 1934, the beginning, and February 20, 1935, the end of the period of treatment. The average count of nematode ova per gram of feces was 6000 on September 26, 1934, and 100 on February 20, 1935, the termination of the treatments.

The general condition of both lots improved materially during the period of treatment from September 1934 to February 1935, as shown in Tables 4 and 5.

At the time of the September 1934 examination of the feces in Lots 1 and 2, tapeworm segments were recorded for sheep nos. 31, 122, 124, and 130 in Lot 1, which subsequently were treated with a 1½% solution of copper sulphate, and sheep nos. 86, 125, 127, 220, and 221 in Lot 2, which subsequently were treated with a mixture of equal parts of a 1½% solution of copper sulphate and a 1½% solution of nicotine sulphate (Blackleaf 40). The feces in Lots 1 and 2 were examined on November 7, 1934, and again on February 20, 1935, the time of the termination of the experiment. No tapeworm segments were found at either of these examinations.

SUMMARY

Twenty sheep were mass-fed or allowed access to a mixture of copper sulphate and salt, 1:30. They showed an average of 150 nematode ova per gram of feces at the time the feeding was begun. One year later the average count of nematode ova in the feces was 1100 per gram, or an increase of over 600% in nematode ova. Even with this increase in nematode infestation the condition of the flock remained good. During this period two sheep died of poisoning from excessive amounts of copper sulphate.

A change in the proportion of copper sulphate to salt from 1:30 to 1:50 resulted in an increase of nematode ova in the feces from 1100 to 4500 ova per gram in three months. This was an increase of slightly more than 300% in nematode ova. One sheep died from poisoning three weeks following the reduction in the proportion of the copper sulphate in the mixture.

The 7 sheep in the treated control lot, or those receiving a 1½% solution of copper sulphate every 21 days, showed a decrease in the average count of nematode ova in the feces from 380 to 50 per gram, and the condition of the sheep had improved over the period of one year.

The 4 sheep in the untreated control lot showed an average increase in the nematode ova in the feces from 50 to 3400 per gram;

these sheep also showed a marked decline in their condition during this one-year period.

As a further check on the effectiveness of regular drenching versus mass feeding, the 12 ewes and their 11 lambs were divided into two lots as nearly uniform as possible. One lot was treated with a 1½% solution of copper sulphate and the other with a mixture of equal parts of a 1½% solution of copper sulphate and a 1½% solution of nicotine sulphate (Blackleaf 40).

In the lot receiving the copper-sulphate solution the nematode ova count was reduced from 6300 to 80. In the lot receiving the mixture of copper sulphate and nicotine sulphate the nematode ova were reduced from 6000 to 100 per gram. These treatments covered a period of eight months.

Tapeworm segments disappeared from the stools equally well in the lots treated with copper sulphate and the copper-sulphate—nicotine-sulphate mixture.

CONCLUSIONS

While the number of sheep in this experiment was small, the results would indicate that the mixture of copper sulphate and salt, 1:30, when fed to sheep *ad libitum* and consumed at the rate of one-half pound per sheep per month, did not adequately control the development of nematode parasites in the gastro-intestinal tract, yet caused the death of sheep from poisoning. A mixture of copper sulphate and salt 1:50 was wholly inadequate in controlling nematode parasites in the gastro-intestinal tracts of these sheep.

The 1½% solution of copper sulphate and the mixture of equal parts of a 1½% solution of copper sulphate and a 1½% solution of nicotine sulphate were equally efficient in the control of nematode infestation. Cestode infestation also disappeared equally under the two treatments.

LITERATURE CITED

- (1) BELL, D. S.
1926. STOMACH WORMS AND NODULAR WORMS IN LAMBS. Ohio Agr. Exp. Sta. Bul. 401.
- (2) BELL, D. S.
1935. PERSONAL COMMUNICATION. Ohio Agr. Exp. Sta.
- (3) BOUGHTON, I. B., and HARDY, W. T.
1934. CHRONIC COPPER-SULPHATE POISONING IN SHEEP. Texas Agr. Exp. Sta. Bul. 499.
- (4) EVVARD, J. M., BROWN, L. C., CULBERTSON, C. C., and HAMMOND, W. E.
1926. THE SALT CONSUMPTION OF SHEEP AND FATTENING LAMBS. Iowa Agr. Exp. Sta. Res. Bul. 94.
- (5) HARDY, W. T., and SCHMIDT, H.
1932. STOMACH WORMS IN SHEEP AND GOATS. Texas Agr. Exp. Sta. Bul. 448.
- (6) HUTCHESON, D.
1891. THE TREATMENT OF WIRE WORMS. Agr. Jour. Good Hope 4: 8-9. South Africa.
- (7) NIGHBERT, E. M.
1932. CONTROLLING STOMACH WORMS IN SHEEP AND LAMBS. U. S. Dept. of Agr., B. A. I. Leaflet No. 89.
- (8) RIETZ, J. H.
1935. COPPER SULPHATE AS AN ANTHELMINTIC FOR GASTRO-INTESTINAL PARASITES OF SHEEP. W. Va. Agr. Exp. Sta. Bul. 264 (Out of print).
- (9) STOLL, N. R.
1923. INVESTIGATIONS ON THE CONTROL OF HOOKWORM DISEASE: XV. AN EFFECTIVE METHOD OF COUNTING HOOKWORM EGGS IN FECES. Am. Jour. Hyg. 3:59-70.