1-1-1989

Yield and quality of winter annual forages

Julio Cesar Medel

John Allan Balasko

Follow this and additional works at: https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins

Digital Commons Citation

Medel, Julio Cesar and Balasko, John Allan, "Yield and quality of winter annual forages" (1989). *West Virginia Agricultural and Forestry Experiment Station Bulletins*. 701.
https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins/595

This Bulletin is brought to you for free and open access by the Davis College of Agriculture, Natural Resources And Design at The Research Repository @ WVU. It has been accepted for inclusion in West Virginia Agricultural and Forestry Experiment Station Bulletins by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

YIELD AND QUALITY OF WINTER ANNUAL FORAGES

S127
.E1
no. 701
Nov. 1989
C. 2

AGRICULTURAL AND FORESTRY EXPERIMENT STATION
WEST VIRGINIA UNIVERSITY

Bulletin 701
November 1989

West Virginia University Libraries

3 0802 100365076 8

Authors

Julio C. Medal is a former graduate student and John A. Balasko is professor of agronomy, Division of Plant and Soil Sciences, College of Agriculture and Forestry, West Virginia University.

West Virginia University
Agricultural and Forestry Experiment Station
College of Agriculture and Forestry
Robert H. Maxwell, Director
Morgantown

DEC 21 '94 MAY 18 '95

AUG 13 1997

Contents

Abstract	iii
List of Tables	v
Introduction	1
Literature Review	3
<i>Rye</i>	3
<i>Austrian winter pea</i>	5
<i>Hairy vetch</i>	7
Materials and Methods	9
Results and Discussion	13
<i>Weather conditions</i>	13
<i>Dry matter yields of winter forages</i>	
<i>harvested in early spring</i>	13
<i>Dry matter yields of winter rye harvested</i>	
<i>in early winter and spring.</i>	17
<i>Dry matter yields of winter forages</i>	
<i>harvested in late spring.</i>	17
<i>Forage digestibility</i>	18
<i>Crude protein of winter forages</i>	
<i>harvested in early spring.</i>	21
<i>Crude protein of winter rye harvested</i>	
<i>in early winter and spring.</i>	22
<i>Crude protein of winter forages</i>	
<i>harvested in late spring</i>	25
<i>Ground cover</i>	25
Summary and Conclusions	29
References	31

WEST VIRGINIA UNIVERSITY
EMMONS LIBRARY

DEC 18 1991

Abstract

Winter annual cover crops have been used successfully to control soil erosion and to provide forage during winter and spring on otherwise idle corn land. The effects of late summer or early fall seeding and late winter or early spring harvest on dry matter yield, crude protein, *in vitro* dry matter digestibility, and ground cover of Abruzzi and Wheeler rye, hairy vetch, and Austrian winter field pea were measured in field experiments during a two-year period (1983 to 1985) at three locations (Reedsville, Morgantown, and Point Pleasant) in West Virginia.

Dry matter yields of Abruzzi rye harvested in late April were 28% higher than those of Wheeler rye for all seeding dates and locations during the two-year study. Mean yields for the mid-September, early October, and mid-October planting dates were 2,808, 2,658, and 1,782 lb. per acre, respectively. Abruzzi rye matured earlier than Wheeler rye and was more susceptible to snow mold fungus at Reedsville during the second year of the experiment. At Morgantown and Reedsville, mean yields for field pea and hairy vetch planted in mid-September and harvested in late April were 670 and 934 lb. per acre, respectively. Delaying harvest of these crops until 22 May at Morgantown resulted in respective yields of 2,442 and 3,481 lb. per acre. Field pea appeared to be somewhat better adapted to later planting than hairy vetch in that its yields from early October and mid-October plantings were higher. Few legume plants survived the winter on the sandy and somewhat droughty soil at Point Pleasant.

In vitro dry matter digestibility (IVDMD) of rye forage harvested in late April usually increased as seeding date was delayed from mid-September (72.5%) to mid-October (76.5%). Digestibility and crude protein values were always higher for Wheeler rye than for Abruzzi rye. Mean digestibility values were 79.0% for Wheeler and 69.8% for Abruzzi. Crude protein values ranged from 6.9 to 15.6% for Abruzzi and from 9.8 to 18.6% for Wheeler. Winter rye varieties harvested in an advanced stage of maturity (in late May) produced poor quality forage. Crude protein concentrations ranged from 3.9 to 6.9%, and IVDMD ranged from 44.8 to 55.2%. These values were significantly lower than those for rye harvested in late April. Digestibility and crude protein values for the winter annual legumes were usually higher than those for the rye varieties. Digestibility values for forage harvested in late April ranged from 76.6 to 86.0% for field pea and from 72.9 to 83.4% for hairy vetch. Crude protein values ranged from 15.7 to 27.0% for field pea, and from 17.2 to 26.0% for hairy vetch.

Rye varieties provided more ground cover during winter than field pea and hairy vetch. Ground cover of winter rye varieties two months after seeding was greater than 78% at all locations for the mid-September and early October plantings.

Differences in forage yield and quality of the rye varieties appeared to be related to their difference in maturity. Based on the data collected, Abruzzi rye must be harvested earlier than Wheeler rye to optimize yield of quality forage. Rye varieties should be chosen based upon expected planting date of corn. Abruzzi rye should be the choice for early-planted corn and Wheeler rye the choice for late-planted corn. Rye should be planted on or before October 1. Hairy vetch and field pea provided a significant amount of forage when harvest was delayed until late May. Winter annual legumes should be planted in mid-September or earlier because they establish slowly.

Tables

Table 1. Descriptions, climates, and soils of the locations where experimental plots were established.	10
Table 2. Soil analyses of samples collected prior to planting in 1983 and 1984.	11
Table 3. Schedule of seeding and harvest dates.	11
Table 4. Climatological data for the experimental areas at Morgantown and Reedsville, West Virginia, 1983-1985.	14
Table 5. Climatological data for the experimental area at Point Pleasant, West Virginia, 1984-1985.	15
Table 6. Dry matter yields of field pea, hairy vetch, and winter rye forages harvested in April 1984 and April 1985.	16
Table 7. Dry matter yields of winter rye forages harvested in December 1983 and April 1984.	18
Table 8. Dry matter yields of field pea, hairy vetch, and winter rye at Morgantown, during the 1983-1984 period.	19
Table 9. In vitro dry matter digestibility of field pea, hairy vetch, and winter rye forages harvested in April 1984 and April 1985 .	20
Table 10. Crude protein and in vitro dry matter digestibility values of winter rye harvested in December 1983 and April 1984.	22
Table 11. In vitro dry matter digestibility of field pea, hairy vetch, and winter rye forages at Morgantown during the 1983-1984 period.	23
Table 12. Crude protein concentrations of field pea, hairy vetch, and winter rye forages harvested in April 1984 and April 1985.	24

Table 13. Crude protein of field pea, hairy vetch, and winter rye forages at Morgantown during the 1983-1984 period.	26
Table 14. Ground cover provided by field pea and hairy vetch in April 1984.	27
Table 15. Ground cover provided by field pea, hairy vetch, and winter rye at Morgantown during the 1984-1985 period.	27
Table 16. Ground cover of field pea, hairy vetch, and winter rye at Reedsville during the 1984-1985 period.	28
Table 17. Ground cover of field pea, hairy vetch, and winter rye at Point Pleasant during the 1984-1985 period.	28

YIELD AND QUALITY OF WINTER ANNUAL FORAGES

Julio C. Medal and John A. Balasko

Introduction

Ruminant livestock production in West Virginia generates about 50% of the cash receipts from farm sales (West Virginia Department of Agriculture, 1987). Producers choose and manage forages that are used for livestock production based on economic considerations, availability of resources, and other factors.

Traditionally, corn grown for silage in West Virginia is harvested in late summer or fall. Much of the land devoted to this crop lies idle and is subject to erosion during the winter season. Furthermore, farmers often face shortages of forage during late winter and early spring and begin grazing when temperatures are not warm enough to allow for rapid plant growth. Additional erosion occurs as a result of overgrazing.

Planting winter annual crops in late summer or fall is an alternative management practice that can be used to provide forage during late winter and/or early spring. Both grass and legume species are available. Winter rye (*Secale cereale* L.) matures earlier and is better able to survive winter conditions in West Virginia than other small grains. Forage legumes that have been used in different regions of the United States are field pea (*Pisum sativum* subsp. *arvense* (L.) Poir.) and hairy vetch (*Vicia villosa* Roth). In addition to providing ground protection during the winter, winter annual legumes may also provide a significant amount of high quality forage and fixed nitrogen that will be available to the subsequent crop.

One of the most important factors affecting winter survival and plant development is time of planting; therefore, experiments were conducted in West Virginia to evaluate the effects of late summer and fall seeding dates on forage yield and quality of winter rye, field pea, and hairy vetch harvested during late winter or early spring. Ground cover provided by the winter forages was also estimated.

Literature Review

Rye

Winter cereals are used for winter and early spring pasture in many regions of the U.S. (Briggle, 1959). They provide opportunities for grazing when other pastures are semidormant and unproductive. Studies in the northern U.S. and Canada (Klebesadel, 1969; Gusta and Fowler, 1976; Agriculture Institute of Canada, 1982) have shown that winter rye is the most valuable of the fall-sown winter cereals for extending the grazing season because it is more winter-hardy than wheat (*Triticum aestivum* L.), barley (*Hordeum vulgare* L.), and oat (*Avena sativa* L.). Rye grows rapidly in early spring, and it yields well.

Smith (1974) compared the yields of five triticale varieties, Arthur wheat, and Balbo rye planted in the fall at Knoxville, Tennessee for a two-year period. Balbo rye yielded an average of 892 lb. per acre more forage than the next highest variety. Sprague (1954) found that in New Jersey, rye, wheat, and oat produced forage in approximately a 3:2:1 relationship. He concluded that the forage production of rye (3,043 lb. per acre) and wheat (1,911 lb. per acre) made a significant contribution to the forage needs of a farm for the period considered. This study also indicated that when rye and wheat were grazed in both fall and spring, approximately one-third of the forage was produced in fall and two-thirds in spring.

Bishnoi (1980) studied the effect of seeding rates and row spacing on forage production of Wrens Abruzzi rye, 6TA 131 triticale, and Arthur 71 wheat in Alabama. Yields tended to increase with increases in seeding rate (45 to 90 lb. per acre) and with a decrease in row spacing (10 to 5 inches). The rye produced the highest forage yield (4,335 lb. per acre) across all seeding rates and row spacings. Rye yield was 8 and 9% higher than wheat and triticale yields, respectively. A similar trend, but with larger differences, was found by Brown and Almodares (1976) in a two-year experiment conducted in Georgia. In both years, Athens Abruzzi rye yielded significantly more than varieties of triticale, wheat, and oat. In contrast, Bishnoi and Hughes (1979), in Alabama, found that average forage yields of winter rye (3,273 lb. per acre) and winter triticale (3,181 lb. per acre) did not differ significantly and that both exceeded wheat forage yield by more than 20%. Fribourg *et al.* (1982) found that the average dry matter yields of rye varieties (3,849 lb. per acre) in Tennessee were 6% lower than those obtained with barley varieties and about the same as those obtained with wheat and oat varieties.

In a review of rye in the southeastern states, Morey (1972) concluded that for early grazing, rye should be planted as early as possible — late August in north Georgia and September in central Georgia. Briggles (1959) found that when rye is planted as a cover crop, seeding early in the fall allows for growth which will provide protection of the soil during the winter months.

Rye is more productive on infertile, sandy, and acid soils than are wheat, oat, or barley (Briggles, 1959). Rye will grow on a wide range of soil types with adequate drainage. Morey (1972) found that a pH of 5.6 to 6.5 was optimum for growing rye.

Fall-sown rye is susceptible to a cool-weather disease known as pink snow mold or *Fusarium* patch, caused by a fungus formerly assigned as *Fusarium* spp. (Westcott, 1971). Recently the causal agent was reassigned from the genus *Fusarium* to *Gerlachia* (Smiley, 1984). Plants are infected in the winter by wind blown ascospores or by mycelia growing on previously infected residue. Cool, wet weather conditions promote this disease. It has also been associated with a snow cover persisting for long periods (Nyvall, 1979; Smiley, 1984).

Smith (1974) in Tennessee compared crude protein concentration of fall-planted cereals harvested at different stages of growth. Rye forage contained 21 to 23% crude protein when harvested in early spring, and its crude protein concentration was reduced to 15% when the harvest was delayed until the middle of May. The same investigator evaluated the effect of seeding dates on crude protein concentration of five triticale varieties, a wheat variety, and Balbo rye. The average crude protein percentage of the varieties tested was significantly higher when planted 10 October than when planted 6 November 1972, and harvested in the vegetative growth stage on 2 April 1973. The author stated that the average percentage of protein of forage from either planting date was acceptable with regard to quality standards. Contrary to these results, Klebesadel (1969) found that the crude protein concentration (20-26%) of winter rye forage growing in south-central Alaska was not influenced by planting date.

In Georgia, Brown and Almodares (1976) found that the crude protein concentrations of rye forages were similar to those of wheat, oat, and triticale at comparable growth stages, with a few exceptions. Average crude protein percentages of rye forage harvested 7 December and 22 January were 29.6 and 25.3, respectively. Values decreased to 11.9% when forage was harvested 10 April. This value is similar to the crude protein value of 10.5% assigned by the National Academy of Science (1971) for rye hay in a dough stage. It is lower than the crude protein concentration reported by Kimber (1965) for April rye forage (14.9%) or April rye regrowth (16.6%) in England. In conclusion, the

crude protein concentration of rye forage appears to be of the same magnitude as that of other cereals, and it is influenced by seeding and harvest dates.

A number of workers have pointed out that digestibility of forage dry matter declines with plant maturation. For example, Kimber (1965) reported that digestibility of rye dry matter decreased from 80.4% when harvested in early March to 73.2% when harvested a month later in England.

Austrian winter pea

Legumes seeded alone or in mixtures with perennial grasses or small grains have received renewed interest as green manure crops in different regions of the U.S. because of the increasing cost of nitrogen fertilizers (Auld et al., 1982). In addition, they often provide forage in late winter or early spring (Markarian and Andersen, 1966). Field pea and vetch are among the most winter-hardy annual forage-legume crops. A pea that is capable of surviving severe winters, mainly with adequate snow cover present, is Austrian winter field pea. It was introduced into Idaho from Austria in 1932 (Slinkard and Murray, 1979). It is used as livestock feed mainly in the northern states from New York west to Washington and Oregon. In the southern states (cotton belt), it has been used as a green manure crop. Researchers at the University of Idaho have promoted new varieties of this crop and improved management practices for its use. Austrian winter field pea grows well on clay loam soils with adequate drainage and on fertilized sandy loam soils of neutral or slightly acidic reaction (Martin et al., 1976).

Auld et al. (1982) evaluated four Austrian winter pea lines planted in the fall and spring in Idaho. They found that the lines produced an average of 11,596 lb. of organic matter per acre when planted in fall, and only 6,155 lb. per acre when planted in spring. Similar results were obtained by Heneise and Murray (1980). They reported that in Idaho, spring-planted Austrian winter pea produced about one-third to one-half as much as fall-planted pea. Auld et al. (1982) found that late summer planting (2 September) of pea produced an organic matter yield 71% higher than that obtained from early fall planting (4 October).

In Alabama, Funchess (1928) found that Austrian pea yielded more than twice as much as hairy vetch and that dry matter yields decreased as date of seeding was delayed. In contrast, Albrecht (1942) reported an average dry matter yield of 7,553 lb. per acre for hairy vetch which was 13% higher than the average yield obtained for Austrian

winter pea over a 15-year period in three regions of Alabama. The study of Funchess (1928) also showed that dry matter yields decreased as the date of seeding was delayed.

The protective influence of snow cover was evident from the study conducted by Markarian and Andersen (1966) at East Lansing, Michigan. They found that when air temperature was below freezing, even a light snow cover had a stabilizing effect on soil temperature, preventing the soil from freezing. Data showed that variable depths of snow resulted in differential plant survival in segregating pea populations. Similar results were obtained by Auld et al. (1983b). They found an average survival of 96% in winter pea lines that were protected during most of the winter by snow cover. The degree of freezing injury depends on the degree of coldness and/or the length of exposure (Hagerdon, 1984). If pea plants are subjected to gradual cooling conditions, young plants may become acclimatized or cold-hardened. They then are better able to withstand exposure to colder temperatures without apparent injury, than plants growing in warmer temperatures and exposed suddenly to cool weather. Swensen and Murray (1983) found that all spring pea seedlings were killed at temperatures of 16 and 10 °F and that pea survival averaged less than 5% at 21 °F. Survival of winter pea seedlings after 4 weeks of being subjected to freezing temperatures of 21, 16, 10 °F were 91, 62, and 7%, respectively.

Klages (1928) compared winter-hardiness of 20 species and varieties of pea and vetches in Oklahoma, and showed that only the Austrian field pea and two varieties of vetches (hairy vetch and woolly-podded vetch) were capable of surviving winter conditions without showing a significant percentage of winter kill. The failure of the non-hardy varieties was attributed to the combined effect of low temperatures and dry winds which produced a rapid desiccation of the plants.

Murray et al. (1978), found that early seeding (September) of winter pea in Idaho allowed greater vegetative growth than late seeding, and also increased the chance of winter survival. A study conducted in England by Dowker (1969) to evaluate the effect of fall seeding dates on winter survival of field pea indicated that there was a significant interaction between time of sowing and variety.

Carter and Larson (1964) pointed out that the protein concentration of field pea hay is similar to that of alfalfa hay, when both crops are cut at comparable growth stages. Henderson and Davies (1955) found crude protein values of 25.1, 22.3, and 19.7% for vegetative, full flower, and fully developed seed growth stages, respectively.

Thorlacius and Beacom (1981), conducted digestibility trials with lambs to evaluate the feeding value of field pea, corn, and oat silages

in Saskatchewan, Canada. They found that field pea silage had an organic matter digestibility of 72.6%. Average values of 62.4 and 68.4% were obtained for oat and corn silages, respectively.

Hairy vetch

Vetches are native to southern Europe and adjacent parts of Asia and Africa, and are currently widely distributed in temperate regions of the world (Heath et al., 1985). They belong to the genus *Vicia* which comprises about 150 species. Some 36 species of native or naturalized vetches are grown in North America. Most of the commonly grown vetches are annuals. In the U.S. vetches have been grown mainly as winter cover crops, and to some extent for hay, silage, and pasture. Vetches produce well on light sandy soils and heavier soils, and they are more tolerant of acid soil conditions than most legume crops (McKee and Schoth, 1961). They respond to P and K fertilizer applications on soils showing low levels of these elements (USDA, 1948). Many of the diseases and insects affecting other forage legumes, also attack vetches (Heath et al., 1985).

Hairy vetch is the most commonly grown vetch in the U.S., and it is the most winter-hardy of the commercial vetches. It is the only one that is recommended for fall planting in the northern U.S. (McKee and Schoth, 1961). North of the 40th parallel, from the Rocky Mountains to the Atlantic coast, hairy vetch should be sown in August or early September.

Hairy vetch, alone or in mixture, extends the grazing season by supplying late-fall and early-spring forage (McKee and Schoth, 1961). Hermann (1960), in his review of the cultivated species of vetches in the U.S., indicated that they are particularly palatable to livestock, very useful as cover crops for providing erosion control, and reduce the leaching of nutrients during the winter months.

Ebelhar et al. (1984) conducted a study in Kentucky from 1977 to 1981 to determine the amount of biologically fixed N provided to no-tillage corn by the winter cover crops, hairy vetch, big flower vetch (*Vicia grandiflora* W. Koch), and crimson clover (*Trifolium incarnatum* L.). They found that hairy vetch produced more dry matter, with a higher N percentage, than the other legumes. These researchers estimated that hairy vetch supplied biologically fixed N equivalent to approximately 80 to 89 lb. per acre of N fertilizer annually to corn. Hargrove (1986) found that hairy vetch top growth contained 136 lb. N per acre and that, after fallow, this N replaced the equivalent of 86 lb. of N fertilizer for a subsequent no-till grain sorghum (*Sorghum bicolor* (L.) Moench) crop. Furthermore, when the legume was chemi-

cally killed, it provided a mulch that continued to control erosion, increase water infiltration, and decrease evaporation during the growing season.

Dry matter yields for species of vetches have been estimated to be from 2,203 to 6,369 lb. per acre by the USDA (1948). When dry matter yields of hairy vetch or other vetches are compared to those of other legumes, there are some contradictory results. For example, field experiments conducted by Ebelhar et al. (1984) in Kentucky indicated that hairy vetch produced significantly more dry matter than crimson clover and big flower vetch. On the other hand, a study conducted in Florida (Dunavin, 1982) to compare cool-season forages, indicated that crimson clover and arrowleaf clover (*Trifolium vesiculosum* Savi) yielded significantly more than the four vetch varieties tested during a 2-year period.

Planting hairy vetch early in the fall allows for more vegetative growth to accumulate and increases the chance of plant survival. McKee and Schoth (1961) in their review of climatic requirements of vetches, concluded that winter killing of any variety of vetch cannot be associated only with an absolute temperature value. There are several factors, such as soil moisture, winter protection, vigor, and age of the plant, that are directly associated with winter killing. For example, young plants from late seedings are often winter-killed while older plants of earlier seedings survive without injury. They stated that plants which are growing fast, because of high temperature and ample moisture, may receive more injury from freezing than plants that are growing slowly and have had a gradual approach or acclimation to a period of freezing. Hairy vetch survives temperatures lower than 0 °F in regions where temperature fluctuations are slight or where a snow cover exists.

Materials and Methods

Experiments were designed to measure the effects of late summer and fall seeding dates on yield, crude protein, *in vitro* dry matter digestibility and ground cover of field pea, hairy vetch and winter rye. They were conducted on West Virginia University experiment farms at Morgantown and Reedsville from September 1983 to April 1985 and at Point Pleasant from September 1984 to April 1985.

Climate and soils of the locations where the experimental plots were established are presented in Table 1. Weather data for the 1983-1985 period were taken from Reedsville experiment farm records and from National Oceanographic and Atmospheric Administration bulletins. Soil samples were collected at Morgantown and Reedsville before planting. Results of the analyses of these samples are presented in Table 2. Prior to each year's seeding, 446 lb. per acre of 10-20-20 fertilizer was applied to the soil.

The experimental design was a split plot with four replications. The treatments were three seeding dates (mid-September, early October, and mid-October) and four varieties (Austrian winter field pea, common hairy vetch, Wrens Abruzzi rye, and Wheeler rye), arranged in a 3 x 4 factorial. The seeding dates made up the main plots and the varieties the sub-plots. The plot size for the first period (from September 1983 to April 1984) was 108 square ft. (16.4 x 6.6 ft.), and plot size for the second period (from September 1984 to April 1985) was 72 square ft. (16.4 x 4.4 ft.). Each plot had eight rows with 10 inches between rows for the first period and nine rows with 6 inches between rows for the second period.

The fields were plowed and harrowed about 2 weeks before the first seeding. Seedings were made with a Tye no-till seeder the first year and a small gasoline-engine forage seeder the second year. Seeding rates for hairy vetch, field pea, and rye were 40, 70 and 90 lb. per acre, respectively.

Legume seeds were inoculated with the appropriate Rhizobia species prior to seeding in the first year. Seeding and harvest schedules are shown in Table 3. Plots (three and five center rows for the first and second periods, respectively) were harvested with a sickle bar mower to a two-inch stubble. Rye cultivars were harvested between the boot and early flowering stages, and legumes were harvested between bud and early flowering in late April. Rye cultivars planted in mid-September 1983 were also harvested in December 1983. An additional harvest for all of the crops was made in late May 1984 at Morgantown. This harvest included additional treatments (plots harvested for the

Table 1. Descriptions, climates, and soils of the locations where the experimental plots were established in West Virginia*.

Variable	Location		
	Point Pleasant	Morgantown	Reedsville
Elevation (ft.)	569	1,240	1,761
Longitude	82°08'W	79°50'W	79°47'W
Latitude	38°51'N	39°40'N	39°30'N
Climate			
Temperature (°F)			
Winter	34	32	29
Spring	55	51	49
Summer	74	72	68
Fall	56	55	51
Year	55	52	49
Precipitation (in)			
Annual average rainfall	41.5	39.2	48.4
Annual average snowfall	17.8	31.8	57.8
Soil Series	Lakin	Lily	Gilpin (1983-84) Rayne (1984-85)
Drainage	excessively drained	moderately to well drained	well drained
Texture	loamy sand	loam	silt loam

* Soil survey of Jackson and Mason counties, West Virginia 1961.
 Soil survey of Marion and Monongalia counties, West Virginia 1982.
 Climatological summary of the Reedsville Experimental Farm,
 1966-1978, unpublished data of W.B. Bryan.

Table 2. Soil analyses of samples collected prior to planting in 1983 and 1984.

Location	Year	Available macro nutrients				pH
		P	K	Ca	Mg	
----- lb. per acre -----						
Reedsville	1983	84	173	3,858	246	6.1
	1984	55	76	2,700	14	5.8
Morgantown	1983	77	134	3,618	415	6.6
	1984	48	60	4,069	136	6.2

Table 3. Schedule of seeding and harvest dates.

Seeding dates	Harvest dates	
	Fall	Spring
Morgantown		
19 September 1983	2 December 1983	20 April, 22 May 1984
4 October 1983	-----	20 April, 22 May 1984
19 October 1983	-----	20 April, 22 May 1984
18 September 1984	-----	24 April 1985
5 October 1984	-----	24 April 1985
15 October 1984	-----	24 April 1985
Reedsville		
15 September 1983	1 December 1983	27 April 1984
30 September 1983	-----	27 April 1984
14 October 1983	-----	27 April 1984
14 September 1984	-----	26 April 1985
5 October 1984	-----	26 April 1985
15 October 1984	-----	26 April 1985
Point Pleasant		
13 September 1984	-----	19 April 1985
27 September 1984	-----	19 April 1985
11 October 1984	-----	19 April 1985

first time in May, regrowth from April, regrowth from December, and second regrowth from April of plots also harvested in December).

A sample of forage from each plot was dried in a forced-draft oven at 131-151 °F for percent dry matter determination. Dry matter yield was expressed in lb. per acre. Samples for quality determinations were then ground in a Wiley mill using a 1 mm screen. Duplicate samples of 0.25 g were used to measure *in vitro* dry matter digestibility (IVDMD) according to the procedure outlined by Barnes (1966). Nitrogen concentration of the forage cultivars was measured using the Kjeldahl procedure (AOAC, 1970). A visual estimation of the percentage of ground cover provided by the legume forages was made in April 1984. During the second year of the study, ground cover for all crops was estimated visually at monthly intervals during the growing period.

Data for each year and location were analyzed separately according to an analysis of variance appropriate for the split-plot design (Dowdy and Wearden, 1983). Duncan's New Multiple Range Test was used to detect significant differences among cultivars and seeding dates.

Results and Discussion

Weather conditions

Temperature and precipitation data, including snowfall, for September through April 1983-1984 and 1984-1985 are shown in Tables 4 and 5. During the 1983-1984 period, temperatures at Morgantown were normal to below normal with the exception of February 1984 which had an average temperature 7 °F above normal. The 1984-1985 period had temperatures below normal during September, November, January, and February. This period was characterized by temperatures warmer than normal during March and April, when plants resumed growth. These warmer temperatures during the 1984-1985 period may account in part for differences in stage of growth at harvest time and higher yields of the winter ryes in relation to the 1983-1984 period.

The total precipitation at Morgantown (Table 4) was similar during both periods, but the monthly distribution was different. Snowfall at Morgantown during the 1984-1985 period (39 inches) was almost double that of the previous winter (20 inches).

Average temperatures at Reedsville during October, December, March, and April of the 1984-1985 period were higher than those recorded the previous period. The total precipitation from September 1984 to April 1985 (23 inches) was 6 inches lower than for the first period. Snowfall (90 inches) was 56% above normal. Heavy snow and low temperatures promoted snow mold infestation of winter rye in some low-lying plots seeded in mid-September.

During the 1984-1985 winter season, Point Pleasant (Table 5) had temperatures below normal for the months of September, November, January, and February and above normal for the months of October, December, March, and April. Precipitation was below normal (1 inch) in September 1984, and during the first and second weeks of October there were two dry periods of 5 and 6 days respectively. Of the total October precipitation (5.8 inches) only 1 inch occurred during the first 18 days of the month, which corresponded with the time of the last two planting dates. A relatively dry period, sandy soil, and a high percentage of winter-killed plants were the factors responsible for the failure of legume seedings at this location.

Dry matter yields of winter forages harvested in early spring

Yields generally decreased as seeding date was delayed (Table 6). With one exception, the highest dry matter yields at all seeding dates

Table 4. Climatological data for the experimental areas at Morgantown and Reedsville, West Virginia 1983-1985.

Month	Year	Temperature (°F)*				Rainfall (in)		Snow (in)
		Max	Min	Mean	Dep.	Total	Dep.	
Morgantown								
Sept	1983	75.5	52.4	64.0	-2.0	2.1	-1.3	---
Oct	1983	63.7	44.0	53.9	-0.5	2.2	-0.4	---
Nov	1983	52.2	34.8	43.5	-0.2	3.2	0.5	0.4
Dec	1983	34.6	20.5	27.6	-6.6	3.3	0.2	2.8
Jan	1984	34.1	21.5	27.8	-1.9	1.4	-1.6	8.5
Feb	1984	48.1	29.9	39.0	7.1	1.8	-0.7	2.7
Mar	1984	45.8	26.7	36.3	-5.1	3.4	-0.2	5.6
Apr	1984	60.3	40.4	50.4	-2.0	5.9	2.4	---
Period	1983-1984					23.3		20.0
Sept	1984	72.4	50.4	61.4	-4.6	2.4	-1.0	---
Oct	1984	68.5	51.4	60.0	5.6	3.8	1.2	---
Nov	1984	50.4	30.7	40.6	-3.1	4.4	1.7	6.0
Dec	1984	52.3	33.5	42.9	8.7	3.3	0.2	2.2
Jan	1985	29.8	16.8	23.3	-6.4	1.5	-1.4	16.0
Feb	1985	38.8	21.0	29.9	-2.0	1.3	-1.2	11.8
Mar	1985	54.5	33.4	44.0	2.6	5.8	2.1	---
Apr	1985	67.7	43.1	55.4	3.0	1.7	-1.8	3.0
Period	1984-1985					24.1		39.0
Reedsville								
Sept	1983	73.8	47.7	60.8	-1.6	2.1	-1.4	---
Oct	1983	63.9	42.1	53.0	1.7	4.0	0.8	---
Nov	1983	53.6	31.6	42.6	1.9	5.1	1.5	7.2
Dec	1983	35.1	18.9	27.0	-5.7	4.7	0.2	9.0
Jan	1984	32.9	15.6	24.2	-2.5	0.8	-2.8	20.5
Feb	1984	47.1	25.3	36.2	7.7	2.4	-0.3	14.0
Mar	1984	43.9	24.3	34.1	-5.1	4.2	0.4	23.0
Apr	1984	58.9	37.3	48.1	-0.8	5.6	1.7	---
Period	1983-1984					28.9		73.7
Sept	1984	72.1	46.3	59.2	-3.2	1.7	-1.8	---
Oct	1984	68.8	48.2	58.5	7.2	4.7	1.5	---
Nov	1984	49.2	27.1	38.2	-2.5	4.6	1.0	13.0
Dec	1984	50.7	32.3	41.5	8.8	3.1	-1.4	5.0
Jan	1985	28.8	14.5	21.6	-5.4	0.4	-3.2	41.0
Feb	1985	38.3	18.0	28.1	-0.4	0.7	-2.0	26.0
Mar	1985	53.4	31.5	42.5	3.2	6.3	2.5	1.0
Apr	1985	66.6	39.0	52.8	4.0	1.4	-2.5	4.5
Period	1984-1985					22.9		90.5

* Max=maximum, Min=minimum, Dep=departure from Normal

Table 5. Climatological data for the experimental area at Point Pleasant, West Virginia 1984-1985*.

Month	Year	Temperature (°F)**				Rainfall (in)		Snow (in)
		Max	Min	Mean	Dep.	Total	Dep.	
Sept	1984	76.4	48.6	62.5	-4.3	1.9	-1.2	---
Oct	1984	71.3	52.2	61.8	7.0	5.8	3.4	---
Nov	1984	53.7	31.0	42.4	-1.5	4.3	1.5	---
Dec	1984	53.2	32.2	42.7	7.9	3.8	0.7	2
Jan	1985	33.1	16.5	24.8	-5.8	2.5	---	8
Feb	1985	40.9	16.9	28.9	-3.8	2.9	---	10
Mar	1985	57.7	32.5	45.1	3.1	3.9	-0.1	---
Apr	1985	70.8	42.6	56.7	4.3	1.4	-2.1	---
Period						26.6		20

*Data taken from the Hogsett Gallipolis Dam station (NOAA bulletins)

**Max=maximum, Min=minimum, Dep=departure from normal

and locations were obtained with Abruzzi rye. When planted in mid-September 1983 and harvested in April 1984 at Reedsville and Morgantown, Abruzzi rye yielded 17% and 84% more, respectively, than Wheeler rye. During the second year of the study, Abruzzi rye yielded 26% less than Wheeler rye at Reedsville. The yield reduction was caused by a snow mold fungus disease (*Gerlachia* spp.) that resulted in the death of some plants. Winter rye planted early was more susceptible than that planted later. Wheeler was more resistant to the snow mold fungus than Abruzzi. The cooler-than-normal temperatures that prevailed during January and February 1985 and a heavier annual snowfall, 32.7 inches above normal (Table 4), that covered the ground for about 6 weeks at Reedsville created ideal environmental conditions for development of snow mold disease.

Yields of the winter ryes were significantly higher than those of the winter legumes. These results are contrary to the findings of Hargrove (1986), who worked in Georgia with Wrens Abruzzi rye and common hairy vetch planted in the fall (mid-October and mid-November) on a sandy loam soil and harvested the second half of May. He found that both crops produced about the same amount of dry matter (4.4 tons per acre).

Yields of field pea and hairy vetch were similar for the late-April harvests except when the crops were planted in mid-September 1983 at Morgantown. Late planting (early or mid-October) did not allow

Table 6. Dry matter yields of field pea, hairy vetch and winter rye forages harvested in April 1984 and April 1985.

Crop	Seeding date		
	Mid-September	Early October	Mid-October
-----lb. per acre-----			
Morgantown-1984			
Field pea	877 c*	---	---
Hairy vetch	1,591 ab	---	---
Abruzzi rye	1,935 a	1,603 a	498 a
Wheeler rye	1,051 bc	815 b	248 a
Reedsville-1984			
Field pea	652 b	---	---
Hairy vetch	897 b	---	---
Abruzzi rye	3,383 a	3,099 a	1,306 a
Wheeler rye	2,903 a	2,620 a	1,163 a
Morgantown-1985			
Field pea	591 c	671 c	343 c
Hairy vetch	655 c	526 c	254 c
Abruzzi rye	4,153 a	4,006 a	3,491 a
Wheeler rye	3,481 b	2,987 b	2,785 b
Reedsville-1985			
Field pea	558 c	607 b	150 c
Hairy vetch	593 c	252 b	---
Abruzzi rye	2,399 b	3,216 a	2,603 a
Wheeler rye	3,244 a	2,890 a	1,950 b
Point Pleasant-1985			
Field pea	---	---	---
Hairy vetch	---	---	---
Abruzzi rye	3,273 a	3,255 a	2,480 a
Wheeler rye	2,256 b	2,091 b	1,295 b

*Within columns at each location, values followed by the same letter did not differ at the 0.05 probability level using Duncan's New Multiple Range Test.

for the development of significant amounts of forage above the mower height at either Morgantown or Reedsville in 1983-1984.

Forage yields of legumes during the 1984-1985 period (Table 6) were low for all locations and seeding dates studied. Few legume plants survived the winter at Point Pleasant. Contributing to this lack of survival was a dry, sandy soil that resulted in slow and erratic germination. The later planting dates seemed to have more of an adverse effect on hairy vetch yields than on field pea yields, although this trend was not significantly different at the 0.05 level according to Duncan's New Multiple Range Test. There was a significant interaction between crop and seeding date at Reedsville in 1985.

Dry matter yields of winter rye harvested in early winter and spring

Dry matter yields of winter rye cultivars that were planted in mid-September 1983 and harvested in December 1983 and April 1984 are shown in Table 7. Yields of winter rye cultivars harvested in December 1983 were similar at each location. Average forage yields were 953 and 574 lb. per acre at Reedsville and Morgantown, respectively. Harvesting at this time (December) reduced April yields compared with those of winter rye cultivars harvested only in April 1984 (Table 6). However, the total yields of rye forage harvested in December 1983 and April 1984 at Morgantown and Reedsville were similar to those of rye cultivars harvested in April 1984 only.

Dry matter yields of winter forages harvested in late spring

Dry matter yields of Abruzzi rye planted either in early October (5,573 lb. per acre) or mid-October (3,048 lb. per acre) at Morgantown (Table 8) and harvested for the first time 22 May 1984 were higher than those obtained with Wheeler rye. However, yield of Wheeler rye planted in mid-September was 3.3% higher than that of Abruzzi rye. Total dry matter yields of the winter rye cultivars harvested for the first time in May, and those harvested in December and May, were about two times greater than the yields of rye cultivars harvested in April and May, and those harvested in December, April, and May. Yields of Wheeler rye regrowth from April and second regrowth from April of plots also harvested in December were 59 and 57%, respectively, higher than those of Abruzzi rye. This difference may be in part attributed to more tillering of Wheeler rye during regrowth periods.

Comparison of the two winter legumes shows that dry matter yield of hairy vetch planted in mid-September and harvested in May 1984

Table 7. Dry matter yields of winter rye forages harvested in December 1983 and April 1984.

Crop	Harvest date	Yield
(lb. per acre)		
Morgantown		
Abruzzi rye	2 December 1983	532 c*
Wheeler rye	2 December 1983	615 c <i>average = 574</i>
Abruzzi rye	20 April 1984	1,284 a
Wheeler rye	20 April 1984	822 b
Reedsville		
Abruzzi rye	1 December 1983	1,026 b <i>2953 avg</i>
Wheeler rye	1 December 1983	880 b
Abruzzi rye	27 April 1984	2,353 a <i>2339</i>
Wheeler rye	27 April 1984	2,325 a

*Within columns at each location, values followed by the same letter did not differ at the 0.05 probability level using Duncan's New Multiple Range Test.

(3,481 lb. per acre) was 43% higher than that of field pea. Dry matter yields of the legume crops were similar when planted later (early and mid-October).

Forage digestibility

In general, IVDMD increased as seeding date was delayed from mid-September to mid-October during both years and at all locations (Table 9). Differences among crops appeared to be associated with stage of maturity at harvest time. Abruzzi rye matured earlier than Wheeler rye, and hairy vetch matured earlier than field pea.

During 1983-1984, the highest IVDMD values at Morgantown (85.3%) and Reedsville (88.4%) were obtained with Wheeler rye planted in mid-October, and field pea planted in mid-September, respectively. During the second year the highest IVDMD values at Reedsville (86%) and Morgantown (81.6%) were obtained when field pea was planted in mid-October. The highest IVDMD value at Point Pleasant was obtained when Wheeler rye was planted in mid-October.

Table 8. Dry matter yields of field pea, hairy vetch and winter rye at Morgantown during the 1983-1984 period.

Crop	Harvest dates			Total Yield
	December 1983	April 1984	May 1984	
----- lb. per acre -----				
Planted mid-September				
Abruzzi rye	---	---	5,813	5,813
	---	1,935	807	2,742
	532	---	4,614	5,146
	532	1,284	780	2,596
Wheeler rye	---	---	6,002	6,002
	---	1,051	1,281	2,332
	615	---	4,786	5,401
	615	822	1,227	2,665
Field pea	---	877	---	877
	---	---	2,442	2,442
Hairy vetch	---	1,591	---	1,591
	---	---	3,481	3,481
Planted early October				
Abruzzi rye	---	---	5,573	5,573
	---	1,603	914	2,517
Wheeler rye	---	---	4,158	4,158
	---	815	1,569	2,384
Field pea	---	---	1,839	1,839
Hairy vetch	---	---	1,985	1,985
Planted mid-October				
Abruzzi rye	---	---	3,048	3,048
	---	498	896	1,394
Wheeler rye	---	---	2,921	2,921
	---	248	1,396	1,644
Field pea	---	---	897	897
Hairy vetch	---	---	761	761

Table 9. In vitro dry matter digestibility of field pea, hairy vetch and winter rye forages harvested in April 1984 and April 1985.

Crop	Seeding date		
	Mid-September	Early October	Mid-October
----- (%) -----			
Morgantown-1984			
Field pea	84.8 a*	---	---
Hairy vetch	81.8 b	---	---
Abruzzi rye	73.8 c	75.8 b	79.9 b
Wheeler rye	82.6 b	82.3 a	85.3 a
Reedsville-1984			
Field pea	88.4 a	---	---
Hairy vetch	83.4 b	---	---
Abruzzi rye	72.7 d	76.5 b	79.0 b
Wheeler rye	80.2 c	80.5 a	83.7 a
Morgantown-1985			
Field pea	76.6 a	78.1 a	81.6 a
Hairy vetch	75.4 a	72.9 b	73.4 b
Abruzzi rye	61.1 c	61.3 c	62.3 c
Wheeler rye	72.3 b	73.0 b	73.7 b
Reedsville-1985			
Field pea	83.2 a	84.6 a	86.0 a
Hairy vetch	78.7 b	81.0 b	---
Abruzzi rye	65.1 d	69.2 d	72.3 c
Wheeler rye	75.3 c	78.3 c	80.7 b
Point Pleasant-1985			
Field pea	---	---	---
Hairy vetch	---	---	---
Abruzzi rye	63.9 b	65.5 b	68.1 b
Wheeler rye	78.2 a	78.0 a	80.3 a

*Within columns at each location, values followed by the same letter did not differ at the 0.05 probability level using Duncan's New Multiple Range Test.

The IVDMD values of field pea were always higher than those of hairy vetch, for all seeding dates, locations and years. Similarly, IVDMD values of Wheeler rye were always higher than those of the earlier-maturing Abruzzi rye. The IVDMD values of winter rye cultivars planted in mid-September 1983 and harvested in December 1983 and April 1984 ranged from 75.8 to 83.1 (Table 10).

Delaying harvest of winter rye cultivars from late April to late May resulted in a decrease of *in vitro* dry matter digestibility (Table 11) from 73.8% to 47.2% for Abruzzi rye, and from 82.6 to 52.8% for Wheeler rye. Rye forage harvested at an advanced stage of maturity in late May would not meet the maintenance requirements of most livestock. The IVDMD values of winter legumes harvested for the first time in May 1984 at Morgantown ranged from 69.6 to 73.3% (Table 11).

Crude protein of winter forages harvested in early spring

Crude protein concentrations of the forage legumes were greater than those of the winter ryes at all locations (Table 12). Results obtained for the rye cultivars in 1983-1984 generally showed increasing crude protein concentrations as seeding date was delayed. Results were not consistent in 1984-1985.

During 1983-1984, the highest crude protein concentrations at Morgantown (27.0%) and Reedsville (26.6%) were obtained when field peas were planted in mid-September. These values were significantly higher than those of other forages. Comparison of the two rye cultivars (Table 12) shows that the average crude protein concentrations of Wheeler rye at Morgantown (14.8%) and Reedsville (18.0%) for the first year were significantly higher than the average crude protein values of Abruzzi rye at Morgantown (11.0%) and Reedsville (15.3%), respectively. However, at all these levels, the protein requirements in the total ration for a beef cow (dry pregnant or lactating) or ewe would be met, as established by NRC standards (NRC 1984, 1985).

During the 1984-1985 year, the highest crude protein concentrations at Morgantown (22.6%) and Reedsville (26.0%) were for hairy vetch planted in mid-September. The crude protein concentrations (Table 12) were not affected by seeding dates as in the previous year, with the exception of crude protein in hairy vetch at Morgantown which decreased from 22.6% when planted in mid-September to 17.3% when planted later. Again, at all locations, the crude protein concentrations of Wheeler rye were significantly higher than those of Abruzzi rye. The lower crude protein concentrations of Abruzzi rye may be associated with a "dilution effect." Numerous reports in the literature have

Table 10. Crude protein and in vitro dry matter digestibility values of winter rye harvested in December 1983 and April 1984.

Crop	Harvest date	Crude protein (%)	IVDMD (%)
Morgantown			
Abruzzi rye	2 December 1983	21.4 a*	79.2 b
Wheeler rye	2 December 1983	21.8 a	79.3 b
Abruzzi rye	20 April 1984	10.5 c	77.7 c
Wheeler rye	20 April 1984	13.1 b	83.1 a
Reedsville			
Abruzzi rye	1 December 1983	28.8 a	80.9 a
Wheeler rye	1 December 1983	27.1 a	81.6 a
Abruzzi rye	27 April 1984	14.8 c	75.8 b
Wheeler rye	27 April 1984	17.4 b	81.0 a

*Within columns at each location, values followed by the same letter did not differ at the 0.05 probability level using Duncan's New Multiple Range Test.

indicated a negative correlation between crude protein and dry matter yield and between crude protein and maturity.

The relatively low crude protein values of winter ryes in this experiment are similar to those obtained by Brown and Almodares (1976) with three winter rye cultivars planted 21 September and harvested 10 April in Georgia. In general, the crude protein percentages obtained for rye forage harvested in mid to late April met the minimum requirements for many classes of livestock.

Crude protein of winter rye harvested in early winter and spring

Crude protein concentrations of winter rye cultivars that were planted in mid-September 1983 and harvested in December 1983 and April 1984 are shown in Table 10. Crude protein concentrations of winter rye cultivars harvested in December 1983 ranged from 21.4 to 28.8%. These values are comparable to those of winter legumes planted in mid-September 1983 and harvested in April 1984 at Morgantown and Reedsville, and to those of legumes planted in September 1984 and harvested in April 1985 at Reedsville. Crude protein concentrations of winter rye cultivars at the second harvest (April 1984)

Table 11. In vitro dry matter digestibility of field pea, hairy vetch and winter rye forages at Morgantown during the 1983-1984 period.

Crop	Harvest dates		
	December 1983	April 1984	May 1984
----- (%) -----			
Planted mid-September			
Abruzzi rye	---	---	47.2
	---	73.8	61.1**
	79.2	---	43.8
	79.2	77.7*	60.9***
Wheeler rye	---	---	52.8
	---	82.6	64.7**
	79.2	---	54.5
	79.2	83.1*	64.7***
Field pea	---	84.8	---
	---	---	71.9
Hairy vetch	---	81.8	---
	---	---	69.6
Planted early October			
Abruzzi rye	---	---	44.8
	---	75.8	59.1**
Wheeler rye	---	---	52.9
	---	82.3	62.9**
Field pea	---	---	72.4
Hairy vetch	---	---	71.3
Planted mid-October			
Abruzzi rye	---	---	50.2
	---	79.9	61.3**
Wheeler rye	---	---	55.2
	---	85.3	61.2**
Field pea	---	---	73.3
Hairy vetch	---	---	70.3

* Digestibility of regrowth from plots harvested in December.

** Digestibility of regrowth from plots harvested in April.

*** Digestibility of second regrowth from plots harvested in December and April.

Table 12. Crude protein concentrations of field pea, hairy vetch and winter rye forages harvested in April 1984 and April 1985.

Crop	Seeding date		
	Mid-September	Early October	Mid-October
----- (%) -----			
Morgantown-1984			
Field pea	27.0 a*	---	---
Hairy vetch	25.1 b	---	---
Abruzzi rye	10.4 d	10.2 b	12.3 b
Wheeler rye	13.0 c	13.9 a	17.4 a
Reedsville-1984			
Field pea	26.6 a	---	---
Hairy vetch	24.8 b	---	---
Abruzzi rye	14.9 d	15.4 b	15.6 b
Wheeler rye	17.1 c	18.2 a	18.6 a
Morgantown-1985			
Field pea	16.3 b	15.7 a	16.4 a
Hairy vetch	22.6 a	17.4 a	17.2 a
Abruzzi rye	9.3 d	8.4 c	8.5 c
Wheeler rye	12.6 c	11.6 b	10.7 b
Reedsville-1985			
Field pea	25.0 a	25.4 a	23.6 a
Hairy vetch	26.0 a	22.4 b	---
Abruzzi rye	11.0 c	10.8 d	13.3 c
Wheeler rye	13.9 b	14.4 c	15.2 b
Point Pleasant-1985			
Field pea	---	---	---
Hairy vetch	---	---	---
Abruzzi rye	7.2 b	6.9 b	7.1 b
Wheeler rye	10.0 a	10.5 a	9.9 a

*Within columns at each location, values followed by the same letter did not differ at the 0.05 probability level using Duncan's New Multiple Range Test.

were significantly lower than those of the first harvest (December 1983) and similar to those of winter rye cultivars (Table 11) planted on the same date (mid- September 1983) and harvested only in April 1984.

Crude protein of winter forages harvested in late spring

Crude protein concentrations of the winter rye cultivars harvested for the first time in May 1984, or with a previous harvest in December 1983, at Morgantown, ranged from 3.9 to 6.9% (Table 13). At these levels, rye forage would have to be supplemented to meet the protein requirements in the total ration of growing heifers (11% and 9.2% for heifers weighting 730 and 1,460 lb., respectively) or lactating beef cow (9.2% total protein) of average milking ability as established by NRC standards (NRC 1984). Crude protein concentrations of winter rye cultivars harvested in May were significantly lower than those obtained with the same cultivars harvested in April. Crude protein concentrations of the winter legumes harvested in May ranged from 20.4 to 24.1%. These legume forages may be used to meet the protein requirements (16-18% total protein) for high-producing dairy cattle (NRC, 1988). Hairy vetch planted in mid-September or a mixture of hairy vetch and winter rye may be a better alternative than winter rye alone when the forage is to be used in late spring.

Ground cover

Seeding date had a significant effect on ground cover provided by the winter legumes during 1983-1984 (Table 14). Ground cover decreased as seeding date was delayed, which was a reflection of the effect of cold temperature on plant development. The highest percentages of ground cover at Morgantown (92.5%) and Reedsville (48%) were obtained when hairy vetch was planted in mid-September.

During the second year of the study, ground cover was estimated visually at monthly intervals during the growing period for each crop. Data obtained are presented in Tables 15 to 17. As in 1983-1984, seeding date had a significant effect on ground cover and there were some significant interactions between seeding date and cultivars. In general, the winter rye cultivars provided more ground cover than the legumes during the winter season. The percentage of ground cover at harvest time at Morgantown was always higher than 80% with both winter rye cultivars, and higher than 74% at Reedsville and Point Pleasant when planted in either mid-September or in early October.

Table 13. Crude protein of field pea, hairy vetch and winter rye forages at Morgantown during the 1983-1984 period.

Crop	Harvest dates		
	December 1983	April 1984	May 1984
----- (%) -----			
Planted mid-September			
Abruzzi rye	---	---	4.6
	---	10.4	12.0
	21.4	---	3.9
	21.4	10.5	10.9
Wheeler rye	---	---	5.0
	---	13.0	12.5
	21.8	---	6.5
	21.8	13.1	11.1
Field pea	---	27.0	---
Hairy vetch	---	---	22.9
	---	25.1	---
	---	---	20.9
Planted early October			
Abruzzi rye	---	---	4.1
	---	10.2	10.9
Wheeler rye	---	---	6.3
	---	13.9	11.7
Field pea	---	---	20.4
Hairy vetch	---	---	22.8
Planted mid-October			
Abruzzi rye	---	---	5.1
	---	12.3	10.4
Wheeler rye	---	---	6.9
	---	17.4	9.4
Field pea	---	---	22.8
Hairy vetch	---	---	24.1

Table 14. Ground cover provided by field pea and hairy vetch in April 1984.

Crop	Seeding date					
	Mid-Sept	Early Oct	Mid-Oct	Mid-Sept	Early Oct	Mid-Oct
	----- (%) -----					
	Morgantown			Reedsville		
Field pea	69.5 b*	37.3 a	16.5 a	41.8 b	16.8 a	4.3 a
Hairy vetch	92.5 a	43.5 a	13.5 a	48.0 a	17.5 a	3.0 a

*Means within a column followed by the same letter are not significantly different at the 0.05 probability level using Duncan's New Multiple Range Test.

Table 15. Ground cover of field pea, hairy vetch and winter rye at Morgantown during the 1984-1985 period.

Crop	Seeding date	One month	Two months	March 1985	April 1985
		after seeding 1984	after seeding 1984		
		----- (%) -----			
Field pea	Mid-Sept	25 e*	54 f	53 ef	49 ef
	Early Oct	49 bc	64 de	60 de	60 cde
	Mid-Oct	43 cd	48 f	43 fg	39 f
Hairy vetch	Mid-Sept	22 e	56 ef	63 cde	65 cd
	Early Oct	47 c	62 de	55 e	68 bc
	Mid-Oct	41 cd	45 f	41 g	52 def
Abruzzi rye	Mid-Sept	37 d	89 ab	89 a	82 ab
	Early Oct	55 ab	78 c	76 bc	84 a
	Mid-Oct	56 a	71 cd	70 bcd	80 ab
Wheeler rye	Mid-Sept	39 d	92 a	92 a	87 a
	Early Oct	59 a	80 bc	83 ab	82 ab
	Mid-Oct	58 a	71 cd	72 bcd	83 ab

*Means within a column followed by the same letter are not significantly different at the 0.05 probability level using Duncan's New Multiple Range Test.

Table 16. Ground cover of field pea, hairy vetch and winter rye at Reedsville during the 1984-1985 period.

Crop	Seeding date	One month after seeding 1984	Two months after seeding 1984	March 1985	April 1985
		----- (%) -----			
Field pea	Mid-Sept	22 g*	49 cd	31 ef	49 e
	Early Oct	43 b	58 b	37 de	74 bc
	Mid-Oct	29 f	33 e	8 g	30 f
Hairy vetch	Mid-Sept	22 g	49 cd	34 e	72 c
	Early Oct	42 bc	56 bc	21 f	60 d
	Mid-Oct	19 g	24 f	2 g	9 g
Abruzzi rye	Mid-Sept	37 de	90 a	74 b	74 bc
	Early Oct	72 a	88 a	82 ab	81 abc
	Mid-Oct	33 e	43 d	45 d	59 de
Wheeler rye	Mid-Sept	39 c	94 a	88 a	84 ab
	Early Oct	74 a	88 a	92 a	87 a
	Mid-Oct	40 bcd	53 b	58 c	71 c

*Means within a column followed by the same letter are not significantly different at the 0.05 probability level using Duncan's New Multiple Range Test.

Table 17. Ground cover of field pea, hairy vetch and winter rye at Point Pleasant during the 1984-1985 period.

Crop	Seeding dates	One month after seeding 1984	Two months after seeding 1984	March 1985	April 1985
		----- (%) -----			
Field pea	Mid-Sept	30 h*	80 c	35 c	1 e
	Early Oct	38 g	66 d	18 d	2 e
	Mid-Oct	48 f	--	19 d	2 e
Hairy vetch	Mid-Sept	28 h	84 bc	34 c	12 d
	Early Oct	36 g	69 d	18 d	4 e
	Mid-Oct	48 f	--	19 d	6 de
Abruzzi rye	Mid-Sept	60 e	93 a	77 ab	81 a
	Early Oct	67 cd	90 ab	75 b	78 ab
	Mid-Oct	77 ab	--	72 b	72 b
Wheeler rye	Mid-Sept	66 d	94 a	86 a	82 a
	Early Oct	72 bc	93 a	87 a	80 a
	Mid-Oct	79 a	--	72 b	64 c

*Means within a column followed by the same letter are not significantly different at the 0.05 probability level using Duncan's New Multiple Range Test.

Summary and Conclusions

The effects of mid-September, early October, and mid-October seeding dates and early December, late April and late May harvest dates on dry matter yield, crude protein, *in vitro* dry matter digestibility, and ground cover of Wrens Abruzzi and Wheeler rye cultivars, common hairy vetch, and Austrian winter field pea were determined in field experiments at Morgantown and Reedsville, West Virginia, from September 1983 to April 1985 and at Point Pleasant, West Virginia, from September 1984 to April 1985. The experimental design for the fall seeding date experiments was a split plot with four replications. Treatments included a factorial arrangement of three seeding dates and four cultivars. Rye cultivars were harvested between boot and early flowering stages and legumes were harvested between bud and early flowering in late April. Rye cultivars planted in mid-September 1983 were also harvested in December 1983. An additional harvest of all crops was made in late May 1984 at Morgantown.

Dry matter yields of Abruzzi rye harvested in early spring (late April) were higher than those of Wheeler rye at all seeding dates and locations with one exception. At Reedsville in 1984-1985 Abruzzi rye seeded in mid-September was severely infected with the snow mold fungus (*Gerlachia* spp.). As a result, its yield for that seeding date was lower than the yield of Wheeler rye. The cooler than normal temperatures that occurred during January and February 1985 and snowfall that covered the ground for almost six weeks at Reedsville created ideal environmental conditions for development of snow mold disease. Yields of Abruzzi rye, when planted in mid-September 1983 and harvested in April 1984, at Reedsville and Morgantown, were 17% and 84%, respectively, higher than those of Wheeler rye. However, yield of Wheeler rye (6,002 lb. per acre) was 3.3% higher than that of Abruzzi rye when harvested in late May at Morgantown. Mean yields of rye forage for the mid-September, early October, and late October planting dates were 2,808, 2,658, and 1,782 lb. per acre, respectively, when harvested in late April. Both winter rye varieties provided good ground cover ($\geq 78\%$) during winter and early spring when they were planted by early October.

Yields of the winter rye cultivars were significantly higher than those of the winter legumes. At Morgantown and Reedsville, mean yields for field pea and hairy vetch planted in mid-September and harvested in late April were 670 and 934 lb. per acre, respectively. Yields of hairy vetch and field pea planted in mid-September 1983 and harvested in late May 1984 were 3,481 and 2,442 lb. per acre, respec-

tively. Field pea yielded more than hairy vetch when plantings were made in early or late October although yields were low for these planting times. Forage yields of legumes during the second year (April harvest) were low for all locations and seeding dates. Few legume plants established and survived the winter at Point Pleasant because of the excessively drained sandy soil and lack of snow cover.

In general, IVDMD of forage increased as seeding date was delayed from mid-September to mid-October during both years and at all locations. Differences among crops appeared to be associated with stage of maturity at harvest time. Abruzzi rye matured earlier than Wheeler rye, and hairy vetch matured earlier than field pea. The IVDMD values of winter rye cultivars that were planted in mid-September 1983 and harvested in December 1983 and April 1984 ranged from 75.8 to 83.1%. Delaying harvest of winter rye cultivars from late April to late May at Morgantown resulted in a decrease in IVDMD from 73.8% to 47.2% for Abruzzi rye, and from 82.6% to 52.8% for Wheeler rye. The IVDMD values of winter legumes harvested for the first time in May 1984 at Morgantown ranged from 69.6 to 73.3%.

As expected, crude protein concentrations of the forage legumes were greater than those of the winter ryes harvested in early spring at all locations. Crude protein for the rye cultivars in 1983-1984 increased as seeding date was delayed. Crude protein values obtained with the rye cultivars harvested for the first time in May or with a previous harvest in December 1983, at Morgantown, ranged from 3.9 to 6.9%. At these levels, the protein requirements in the total ration of growing heifers or lactating beef cow of average milking ability would not be met. Crude protein concentrations of winter legumes harvested in late May ranged from 20.4 to 24.1%. These legumes may be used to meet protein requirements of high-producing dairy cattle.

Our studies indicate that Abruzzi rye should be harvested earlier than Wheeler rye to optimize yield of quality forage. Forage quality of both varieties decreases rapidly once heading begins. Rye varieties should be chosen based upon expected planting date of corn. Abruzzi rye should be the choice if corn is planted early in spring and Wheeler rye the choice if corn or other row crops are planted later. Rye should be planted on or before October 1. Winter annual legumes should be planted in mid-September or earlier because they establish slowly. Hairy vetch and field pea provided a significant amount of quality forage when harvest was delayed until late May.

References

- Agriculture Institute of Canada 1982. *Effect of cattle grazing on subsequent grain yield of fall rye (Secale cereale L.) in Southwestern Saskatchewan*. Can. J. Plant Sci. 62:795- 796.
- Albrecht, H. R. 1942. **Vetch Varieties for Soil Improvement and Seed Production in Alabama**. Agric. Exp. Stn. Alabama Polytechnic Inst. Bull. 253. 15 pp.
- Association of Official Analytical Chemists. 1970. **Official Methods of Analysis**. AOAC. Washington, D. C. 1015 pp.
- Auld, D. L., Adams, K. J., Swensen, J. B., and Murray, G. A. 1983a. *Diallel analyses of winterhardiness in peas*. Crop Sci. 23:763-766.
- _____, Bettis, B. L., Dial, M. J., and Murray, G. A. 1982. *Austrian winter and spring peas as green manure crops in northern Idaho*. Agron. J. 74:1047-1050.
- _____, Ditterline, R. L., Murray, G. A., and Swensen, J. B. 1983b. *Screening peas for winterhardiness under field and laboratory conditions*. Crop Sci. 23:85-88.
- Barnes, R. F. 1966. *The development and application of in vitro rumen fermentation techniques*. in **Proc. X International Grassland Congress**. Helsinki, Finland. pp. 434-438.
- Bishnoi, U. R. 1980. *Effect of seeding rates and row spacing on forage and grain production of triticale, wheat, and rye*. Crop Sci. 20:107-108.
- _____, and Hughes, J. L. 1979. *Agronomic performance and protein content of fall-planted triticale, wheat, and rye*. Agron. J. 71:359-360.
- Briggle, L. W. 1959. **Growing Rye**. Farmer's Bull. No. 2145. USDA. 16 pp.

- Brown, A. R., and Almodares, A. 1976. *Quantity and quality of triticale forage compared to other small grains*. Agron. J. 68:264-266.
- Carter, J. F., and Larson, K. L. 1964. *Oats, Peas and Vetch for Hay and Silage in North Dakota*. North Dakota State Univ., Agric. Exp. Stn. Bull. 447. 8 pp.
- Dowdy, S. and Wearden, S., Eds. 1983. *Statistics for Research*. John Wiley and Sons, New York, New York. 537 pp.
- Dowker, B. D. 1969. *Field methods of assessing winter hardiness in peas*. Euphytica 18:398-402.
- Dunavin, L. S. 1982. *Vetch and clover overseeded on a bahiagrass sod*. Agron. J. 74:793-796.
- Ebelhar, S. A., Frye, W. W., and Blevins, R. L. 1984. *Nitrogen from legume cover crops for no-tillage corn*. Agron. J. 76:51-55.
- Fribourg, H. A., Carlisle, R. J., and Reich, V. H. 1982. *Yields of warm- and cool-season forages on adjacent soils*. Agron. J. 74:663-667.
- Funchess, M. J. 1928. *Time of seeding and turning vetch for cotton and corn*. J. Am. Soc. Agron. 20:294-297.
- Gusta, L. V., and Fowler, D. B. 1976. *Effects of temperature on dehardening and rehardening of winter cereals*. Can. J. Plant Sci. 56:673-678.
- Hagerdon, D. J., (ed.) 1984. *Compendium of Pea Diseases*. Am. Phytopathological Soc. St Paul, Minnesota. 57 pp.
- Hargrove, W. L. 1986. *Winter legumes as a nitrogen source for no-till grain sorghum*. Agron. J. 78:70-74.
- Harter, L. L. 1968. *Pea Diseases and Their Control*. USDA Farmer's Bull. No. 1735.
- Heath, M. E., Barnes, R. F., and Metcalfe, D. S., (ed.) 1985. *Forages*. The Iowa State University Press. Ames, Iowa. 643 pp.

- Henderson, J. L., and Davies, R. O. 1955. *The yield and composition of mixed cereal-legume crops at different stages of growth.* Part I., Empire J. Exp. Agric. 23:131-144.
- Heneise, H. K., and Murray, G. A. 1980. *Effect of rowspacing on yield of spring planted Austrian winter field pea.* Agron. J. 72:369-371.
- Hermann, F. J. 1960. **Vetches in the United States - Native, Naturalized and Cultivated.** USDA. Agric.Handbook No. 168. 84 pp.
- Kimber, D. S. 1965. *Trials of rye varieties for forage 1957-63.* J. Nat. Inst. Agric. Bot. 10:195-203.
- Klages, K. H. 1928. *Comparative winterhardiness of species and varieties of vetches and peas in relation to their yielding ability.* J. Am. Soc. Agron. 20:982-987.
- Klebesadel, L. J. 1969. *Winter survival and spring forage yield of winter rye varieties in subarctic Alaska as influenced by date of planting.* Agron. J. 61:708-712.
- Markarian, D., and Andersen, R. L. 1966. *The inheritance of winter hardiness in Pisum I.* Euphytica 15:102-110.
- Martin, J. H., Leonard, W. H., and Stamp, D. L. 1976. **Principles of Field Crop Production.** Macmillan Publishing Co., Inc. New York. pp. 755-764.
- McKee, R., and Schoth, H. A. 1961. **Vetch Culture and Uses.** USDA Farmer's Bull. No. 1740. 22 pp.
- Morey, D. D. 1972. *Rye, southern style.* Crops and Soils 24(7):12-14.
- Murray, G. A., Auld, D. L., Kraft, J. M., Lee, G. A., Muehlbauer, F. J., and O'Keefe, L. E. 1978. **Dry Pea and Lentil Production in the Pacific Northwest.** Univ. of Idaho, Agric. Exp. Stn. Bull. 578. 8 pp.

- National Academy of Science. 1971. **Atlas of Nutritional Data on United States and Canadian Feeds**. NAS, Washington, D.C. 772 pp.
- National Research Council. 1988. **Nutrient Requirements of Dairy Cattle**. 6th Revised Ed. NRC, Washington, D. C. 157 pp.
- _____. 1984. **Nutrient Requirements of Beef Cattle**. 6th Revised Ed. NRC, Washington, D. C. 90 pp.
- _____. 1985. **Nutrient Requirements of Sheep**. 6th Revised Ed. NRC, Washington, D. C. 99 pp.
- Nyvall, R. F. 1979. **Field Crop Diseases Handbook**. AVI Publishing Company, Inc., Connecticut. pp. 155-156.
- Slinkard, A. E., and Murray, G. A. 1979. *Austrian winter field pea industry in Idaho*. Econ. Bot. 33:280-283.
- Smiley, R. W., (ed.) 1984. **Compendium of Turfgrass Diseases**. Am. Phytopathological Soc. St Paul, Minnesota. 102 pp.
- Smith, C. W. 1974. *Yield and laboratory evaluations of five triticale cultivars, Arthur wheat, and Balbo rye in East Tennessee*. Ph. D. thesis. The University of Tennessee. 142 pp.
- Sprague, M. A. 1954. *The effect of grazing management on forage and grain production from rye, wheat, and oats*. Agron. J. 46:29-33.
- Swensen, J. B., and Murray, G. A. 1983. *Cold acclimation of field peas in a controlled environment*. Crop Sci. 23:27-30.
- Thorlacius, S. O., and Beacom, S. E. 1981. *Feeding value for lambs of fababean, field pea, corn, and oat silages*. Can. J. Animal Sci. 61:663-668.
- United States Department of Agriculture. 1948. **Grass. Yearbook of Agriculture**. USDA. Washington. p. 719.
- Westcott, C. 1971. **Plant Disease Handbook**. Van Nostrand Reinhold Company, New York, New York. p. 404.

**West Virginia Department of Agriculture, West Virginia Crop
Reporting Service. 1987. West Virginia Agric. Statistics.
C.R. Bull. No. 18. 64 pp.**

Blank Page in Original Bulletin

Blank Page in Original Bulletin

Blank Page in Original Bulletin

Blank Page in Original Bulletin