1-1-1991

Macrolepidopterous moths : collected by blacklight trap at Cooper's Rock State Forest, West Virginia : a baseline study

Linda Butler

Vicki Kondo

Follow this and additional works at: https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins

Digital Commons Citation

Butler, Linda and Kondo, Vicki, "Macrolepidopterous moths : collected by blacklight trap at Cooper's Rock State Forest, West Virginia : a baseline study" (1991). *West Virginia Agricultural and Forestry Experiment Station Bulletins*. 705.
https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins/599

This Bulletin is brought to you for free and open access by the Davis College of Agriculture, Natural Resources And Design at The Research Repository @ WVU. It has been accepted for inclusion in West Virginia Agricultural and Forestry Experiment Station Bulletins by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

Macrolepidopterous Moths

*Collected by Blacklight Trap
at Cooper's Rock State Forest, West Virginia:
A Baseline Study*

Bulletin 705

April 1991

Agricultural and Forestry Experiment Station
West Virginia University

Authors

Linda Butler is Professor of Entomology and Vicki Kondo is Entomology Research Assistant, Division of Plant and Soil Sciences, College of Agriculture and Forestry, West Virginia University.

This research was supported in part by a grant from the USDA Forest Service and in part from funds of the West Virginia Agricultural and Forestry Experiment Station. We would like to acknowledge the assistance of Dr. Edwin Townsend in data management.

**West Virginia University
Agricultural and Forestry Experiment Station
College of Agriculture and Forestry
Robert H. Maxwell, Director
Morgantown**

Macrolepidopterous Moths Collected by Blacklight Trap at Cooper's Rock State Forest, West Virginia: A Baseline Study

Linda Butler and Vicki Kondo

Introduction

Lepidoptera, particularly Macrolepidoptera, have been the subject of considerable interest in recent years as indicators of bio-diversity and environmental balance. A number of Macrolepidoptera (primarily butterflies) are considered endangered or threatened, especially because of habitat destruction.

As the gypsy moth (*Lymantria dispar* L., Lymantriidae) moves into the central and southern Appalachians, concern grows for the possible impact of gypsy moth defoliation and suppression programs on non-target Lepidoptera. To determine such impacts, baseline data collected by appropriate sampling methods are essential.

Various types of light traps have long been used for control, survey or ecological studies of Macrolepidoptera. We report here the results of a three-year study of macrolepidopterous moths found at the West Virginia University Forest at Cooper's Rock State Forest, conducted to establish baseline data prior to build-up of gypsy moth populations. We also discuss use of blacklight trapping for baseline and non-target impact studies.

Literature Review

As early as 1879, light traps were being used to collect insects (Comstock 1879). Williams (1939) listed three objectives in the use of light traps: (1) the collection of insect specimens to study distribution or taxonomy; (2) the study of insect ecology by evaluating numbers of captured specimens active at particular localities or dates and changes in populations due to climatic variation; and (3) a means of control for certain insects. Williams (1939) referred to numerous reports of light traps used for these purposes.

An excellent summary of light-trap studies was presented by Hienton (1974), with a review of trap design changes over the years. Design standards for blacklight traps for insect surveys were given by Harding et al. (1966) and Smith et al. (1974). Light traps of many types have been compared for efficiency in capturing moths. Cantelo (1990) compared blacklight and Robinson mercury vapor traps for catch of agricultural pests. Frost (1957) compared insect captures taken with the Pennsylvania light trap with traps of other designs. Other comparative studies have been reported by Brether-ton (1954), Vaishampayan and Verma (1983), Williams et al. (1955), Smith et al. (1959), Labanowski (1980), Waring (1980), Heath (1966) and Belton and Kempster (1963).

Light traps fitted with polarizing filters were more attractive to certain moths than traps without filters (Davidson et al. 1973). Light-trap collections of economically important moths increased when light traps were baited with phenylacetaldehyde (Cantelo and Jacobson 1979). Novak (1983) described a mercury vapor light trap incorporating an electrified wire grid for sampling moths of agricultural importance. A Texas-type trap equipped with electrodes was found to increase catch numbers and species numbers over traps without electrodes (Mizutani et al. 1982). Hollingsworth et al. (1961) showed that light-trap catches could be altered by enclosing traps with cylinders of hardware cloth of different mesh sizes. They also found that artificial wind breaks to shield traps from effects of prevailing wind increased trap catches. Ammar (1975) increased efficiency of light traps for catching small Lepidoptera and other insects by using wire mesh. Robinson and Robinson (1950) discussed the behavior of moths near light traps of various designs. A recent discussion of light-trap designs and their relationship to capture efficiency was given by Muirhead-Thomson (1991).

Light traps have been evaluated as control agents for various agricultural pests, particularly tomato and tobacco hornworm, European corn borer, cabbage looper, corn earworm and pink bollworm. A review of the subject was given by Cantelo (1974). Results of light trapping for control of economically important species have been mixed. In a three-and-one-half year study of mass trapping of moths isolated on St. Croix, U.S. Virgin Islands, Cantelo et al. (1974) noted that populations of several pest species of Sphingidae and Noctuidae declined. Williams (1939), however, noted no consistent decline in 150 species of Noctuidae and Geometridae, many of which were agricultural pests, that he collected in four years at Rothamsted, U.K. In further collections at the same location over a nine-year period, Taylor et al. (1978) recorded wide fluctuations, but no downward trends in species numbers.

Some arrangements or densities of light traps might increase pest numbers within an area. Debolt et al. (1979) stated that up to 14 times as many cabbage looper eggs were found in lightly trapped as in heavily trapped areas.

Smith (1962) noted that a light trap at one location in Wisconsin appeared to stimulate an outbreak of armyworm by attracting moths into the area.

Extensive literature reports the use of light traps in survey of moths, particularly those of agricultural importance. Attempts have been made to use data to predict pest outbreaks, to forecast population trends, to time pesticide applications and to monitor effectiveness of controls. Light-trap surveys throughout the world have been instrumental in detecting new regional and national records from immigrant species of moths.

Light traps are valuable in many types of ecological and bionomical studies including flight phenology, migration patterns, long-distance flight capability, nocturnal flight time, sex ratios, voltinism, winter survival, reproductive potential and mating status. Morphological studies such as determination of melanism and insect wing length variations use light-trap collections.

The influence of weather on light-trap catches has been the subject of many studies; impact of moonphase, temperature, humidity, rainfall, wind and barometric pressure have been discussed. Light-trap captures are generally higher on warm, humid nights with no moonlight (McGeachie 1989).

A moth may reach a light trap by either chance or attraction; the distance over which attraction operates is highly dependent on light intensity and light wavelength (Hartstack 1979). Most researchers agree that the distance of positive attraction is no more than a few hundred feet. Obviously, this distance is also dependent on sex and species of moth, among other factors. Some species may disperse more widely during some years (as discussed by Sargent, 1976, for *Catocala* during periods of great abundance), and hence by chance may come within attraction distance of light traps. Baker and Sadovy (1978) used mark-release/recapture methods for estimating the distance of response of moths to a 125-watt mercury vapor light trap. They concluded that the response range of the two moth species studied was only about three meters. In tests with 15-watt blacklight traps, *Manduca sexta* (Johannson) (Sphingidae) exhibited attraction within 4.6 m of the radiation source, and for *Heliothis zea* (Boddie) (Noctuidae) the distance was 6.1 m (Stewart et al. 1969). In contrast, *Spodoptera littoralis* Boisduval (Noctuidae) was attracted to a 6-watt blacklight from at least 200 m (Plaut 1971).

Williams (1940) used light-trap catches of insects as a tool for estimating and forecasting changes in insect populations. He stated that it is possible to correct mean insect catch for each month for its variation from the normal due to changes in temperature, wind, and other factors. The result would be a value for what the catch would have been if all conditions had been normal for the month. The differences that remain, according to Williams, are due to population effects. King and Hind (1960) presented a method for the calculation of the theoretical catch level of insects taken in light traps. They

suggested that departures from the theoretical catch are due to high or low activity and must be accounted for by weather. Williams (1940) discussed the concept that short-term (daily) variations in insect catch are functions of activity rather than population changes, and that long-term (seasonal or annual) variations in catch are functions of changing population rather than of activity.

Light trapping is a relative, not absolute, method of population estimation (Southwood 1978). In criticizing the use of light traps for population estimation, Taylor and Carter (1961) stated that the number of any species caught depends not only on their commonness but also on their attractiveness to light. Taylor and French (1974) noted further that the relation between the catch and the parent population is not known, hence the catch cannot be treated as a true sample. They stressed the importance of selecting sampling sites as alike as possible if catches from different traps are to be compared.

Taylor and Brown (1972) noted that a light-trap catch represents a congregation at a point in space. They distinguished three different functions relating catch and space that may help to envisage the catch as a sample. They distinguished between local and migrant activity and discussed the concept of "area of influence" of the trap. White (1989) noted that the trappable area itself is rarely definable. Hartstack et al. (1968) used a series of oil-and-water-filled pans displayed around light traps to measure efficiency of traps. They concluded that large numbers of insects attracted to light traps are not caught. McGeachie (1988) used a video camera to estimate light-trap efficiency in the field. He divided insect flight tracks into three categories: (1) new arrivals, (2) passers by and (3) local flights. While trap efficiency varied with wind speed, he recorded a maximum of 39% efficiency in virtually calm conditions.

Sargent (1976) noted that the relative abundance of a species may vary considerably from year to year at a given location; the overall pattern of species abundance also varies over the years. Sargent's comments were made particularly regarding records of S.A. Hessel who light trapped *Catocala* spp. (Noctuidae) on virtually every night of twelve consecutive *Catocala* seasons. The variation in Hessel's data led Sargent to suggest the futility of making long-term assessments of populations based on records from a single season at any locality no matter how extensive collecting may be that season. Rejmanek and Spitzer (1982) reported results of a 12-year study of fluctuation in populations of noctuid moths in Bohemia. They discussed the relationship between species diversity and species stability in Lepidoptera communities monitored by light trapping.

Spitzer and Leps (1988) studied year-to-year variation in abundance of noctuid moths in three Central European habitats differing in successional status and degree of disturbance. Highest variation was found in moth species with a high population growth rate. Climax habitats supported

populations of moths with low variation in abundance and low population growth rate.

White (1989) discussed the concept of "dilution ratios" of moths as their numbers are reduced due to frequent trapping. Thomas (1989) noted that the optimum frequency for light trapping at a given site was every third to seventh night. He stated that with care, light trapping and live release can provide valid population indices for comparison between generations and years for a given moth species.

Placement of light traps greatly influences moth catch. Hausmann (1990a) found that two light traps located only 45 m apart showed great differences in species composition. He noted that the method of using single light traps has to be regarded as site specific and must be considered in ecological interpretations of data. Williams et al. (1955) discussed the importance of rotating traps among designated trapping sites to minimize effects of differences among traps.

In a study comparing light-trap catches in deciduous and coniferous woodland habitats, Waring (1989) warned that woodland studies are especially challenging because of the amount of shading and screening of light traps by vegetation. He stressed the need for carefully choosing light-trap sites. Magurran (1985) studied differences in *Macrolepidoptera* in Ireland and noted that insect diversity can be correlated with site factors, but as the relationship is a very complex one, such correlations should be interpreted with great care.

Mizutani (1982) arranged 10 light traps in a line at regular intervals of 28.6 m between two forests about 200 m apart. He found that catches of moths were greatest in numbers of species and individuals within about 30 m from the forest edge and concluded that 30 m from the forest edge is the most suitable placement for a light trap.

A number of studies have been conducted on the use of light traps to evaluate the *Macrolepidoptera* community at a single site over a period of several years. Dirks (1937) recorded 344 species of *Macrolepidoptera* collected over a four-year period at one site in Orono, Maine. During his studies at Rothamsted, Williams (1939) collected 356 species over four years. The Rothamsted insect survey, which includes moth monitoring, has continued since 1969. A history of the survey was given by Taylor (1979). During six years of blacklight-trap collections at Archbold Biological Station in Florida, Frost (1964) captured 330 *Macrolepidoptera* species. In a recent study of a New Jersey oak forest, Moulding and Madenjian (1979) recorded 410 species over a five-year period.

Rings et al. (1987) conducted a nine-year study of the *Lepidoptera* of the Wilderness Center, Stark County, Ohio, and recorded 426 species of *macrolepidopterous* moths. In similar studies at Atwood Lake Park, Mohican State Forest and State Park and Fowler Woods, Ohio, Rings and

Metzler (1988, 1989, 1990) listed 382, 417 and 374 forms and species of macrolepidopterous moths, respectively. Sampling methods for these studies included ultraviolet light traps, mercury vapor light, bait traps, sugaring and netting.

Holloway (1984) advocated the use of light-trap sampling of moths as a means of indicating response of rain forest insect communities to disturbance. He concluded that this simple and quick approach uses elements of the fauna that are easily sampled, widely distributed, diverse, show specificity to vegetation type, and are sensitive to change.

Recent and ongoing studies are being conducted in the eastern United States using light trapping to establish baseline data and to evaluate impact of gypsy-moth defoliation or sprays applied as a part of gypsy-moth suppression programs. Peacock and Bullington (1989) recently conducted a study on effects of *Bacillus thuringiensis* Berliner var. *kurstaki* on native Lepidoptera in the Shenandoah National Park, Virginia, using blacklight traps to monitor moth populations. They noted that total season capture of selected species in treated and untreated areas was extremely variable, prohibiting any legitimate analysis of the data.

Sample (1991) recently completed a study of diflubenzuron impact on non-target insects as food of the Virginia big-eared bat in West Virginia. Blacklight traps operated in seven pairs of treated and untreated plots were the primary sampling tools. Sample found that total species richness and species richness in four of eight Lepidoptera families was reduced. Of the 83 macrolepidopterous species in sufficient abundance for statistical analysis, 47 were significantly reduced.

In an ongoing study in the Great Smoky Mountains National Park, Watson and Lambdin (1989) are collecting baseline data on Lepidoptera prior to buildup of gypsy-moth populations. Blacklight traps are being used to collect samples from 10 permanent sites, five each in North Carolina and Tennessee.

Light trapping of insects has provided valuable information on insect distribution, taxonomy, ecology, and control for more than a century. Currently, light traps are particularly useful tools for establishing baseline data and for measuring impacts due to forest disturbance by pesticide spraying.

Materials and Methods

This study was conducted at the West Virginia University Forest at Cooper's Rock State Forest, located in Preston and Monongalia counties about 32 km east of Morgantown, West Virginia. The area consists of a 50- to 60-year old even-aged mixed mesophytic forest and has a mean elevation of 561 m (Carvell 1983). Near the light trapping site the canopy is white and red oak (*Quercus alba* L., *Q. rubra* L., Fagaceae) and the understory is red

maple (*Acer rubrum* L., Aceraceae), black cherry (*Prunus serotina* Ehrh., Rosaceae) and black birch (*Betula lenta* L., Betulaceae). The light-trap site was approximately 200 m from the nearest significant clearing, Sand Spring Road, and was at coordinates 39°40' North latitude by 79°47' West longitude.

From mid-March to late October 1984–1986, a single blacklight trap was operated one night each week. The trap was a 15-watt, photo-cell-controlled UV type (Ellisco Co., Philadelphia) set on tripod legs. Trap nights were chosen to avoid unseasonably low temperatures when possible, but no attempt was made to avoid rainy nights.

Macrolepidopterous moths were identified by comparison with the West Virginia University Arthropod Collection; counts were made for each species. Voucher specimens from the study are deposited in the WVU Arthropod Collection. Species are named and listed after the checklist given by Hodges et al. (1983).

Results

From 1984 to 1986, 400 species of macrolepidopterous moths in 13 families were captured in one blacklight trap operated at a single site at Cooper's Rock State Forest (Tables 1 and 2). The Noctuidae contributed 220 species and the Geometridae 102 species. Considerable variation in frequency and abundance of species occurred among the three years. This variation was especially evident in the Noctuidae; for example, while 220 species were trapped during the three-year study, only 81 of those species were trapped all three years. Sixty-two species were trapped only two of the three years, and 77 were collected only one year during the study (Table 1).

Table 2 lists the 400 species, giving total catch for each year. Each species is ranked with regard to relative abundance each year. The inclusive flight range for all three years from date of first capture to date of last capture is included. The most abundant species were the geometrids *Itame pustularia*, *Probole amicaria*, *Lomographa glomeraria*, *Melanolophia canadaria* and *Campaea perlata*. Other abundant species were *Halysidota tessellaris* and *Clemensia albata* (Arctiidae); *Malacosoma americanum* (Lasiocampidae); *Orthodes cynica*, *Idia aemula* and *Zanclognatha ochreipennis* (Noctuidae) and *Nadata gibbosa*, *Heterocampa guttivitta*, and *Peridea angulosa* (Notodontidae).

Considerable fluctuation in abundance of many species occurred from year to year. Counts for *Hyphantria cunea* (Arctiidae) were 153 and 169 for 1984 and 1985, respectively; only 13 specimens were captured in 1986. *Halysidota tessellaris* (Arctiidae) counts dropped from 698 in 1984 to 149 in 1985. *Pero honestaria* (Geometridae) declined from 89 (1984) to 23 (1985) to 0 (1986). *Besma quercivoraria* (Geometridae) captures dropped from 279 to 114 to 25. In 1984, 173 *Orygia leucostigma* (Lymantriidae) were captured,

dropping to 37 in 1985 and rebounding to 163 in 1986. *Idia aemula* (Noctuidae) catches declined from 346 in 1984 to 38 in 1985. *Zanclognatha laevigata* and *Z. ochreipennis* (Noctuidae) both showed a three-fold decline in counts from 1984 to 1985. Counts for *Morrisonia confusa* (Noctuidae) were 79 (1984), 119 (1985) and 0 (1986). *Peridea angulosa* (Notodontidae) counts dropped from 192 in 1984 to 59 in 1985.

While many species showed dramatic declines in trapped numbers from 1984 to 1986, some species increased. Counts for *Probole amicaria* (Geometridae) increased from 133 (1984) to 1073 (1985). Only 61 *Heterocampa guttivitta* (Notodontidae) were trapped in 1984, while 297 were taken in 1986. During that time, *Nadata gibbosa* (Notodontidae) counts also increased from 77 to 240. Counts of some species, such as *Melanolophia canadaria* (Geometridae) and *Lomographa glomeraria* (Geometridae) remained rather stable during the three-year period.

Dates on flight range (Table 2) are given as earliest catch and latest catch for the three years combined. Variation in length of flight season among species reflects varying degrees of synchrony of adult emergence, differences in adult longevity, effect of multivoltine versus univoltine species and population levels. In general, longer flight seasons are shown for species at higher populations since greater numbers would give greater variation in emergence and longevity.

Examples of multivoltine species are *Melanolophia canadaria* (Geometridae), *Campaea perlata* (Geometridae) and *Lomographa glomeraria* (Geometridae), all of which showed relatively high populations and long flight season. *Polia latex* (Noctuidae) is an example of a univoltine species with a long flight season reflecting high populations and relative lack of synchrony of adult emergence. *Lithophane hemina* (Noctuidae) is an example of a species that emerges in the autumn and overwinters as an adult. Moths of different generations are picked up in light traps operated from March through October, but the species is univoltine.

Discussion

The study site at Cooper's Rock State Forest has a highly diverse population of Macrolepidoptera. We collected 400 species of moths at a single blacklight trap site, a higher number than reported in other studies. Where more species were recorded, several different light trap sites were used and/or sampling methods in addition to blacklight traps were employed.

As indicated in Table 1, species richness fluctuated dramatically over the three years of the study. Hausmann (1990b) noted that in a two-year study of light-trapped moths near Munich, Germany, the "species spectra" covered by one light trap showed a year-to-year turnover of 35%–55%. The yearly turnover in our study falls within this expected range.

The results of our study and reports of others cited in this paper can give us some guidance on use of light traps in conducting baseline studies or in determining non-target impacts of insecticides or some other disturbance. We recommend consideration of the following items:

- Collect as many years of baseline data as possible to gain insight into expected "normal fluctuation" of light-trapped moths from year to year. Operate as many light traps as possible over the study area so that variation will even out among sites and over years. Disturbed sites, by their nature, are likely to show wider fluctuations in populations from year to year (Spitzer and Leps 1988).
- Do not operate traps too frequently at each site. Thomas (1989) suggested that trapping every three to seven nights was optimum. If traps are operated less frequently than every two weeks, some moth species, especially those in low populations, may not be captured.
- Carefully choose light-trap sites if data from traps on different sites are to be compared. Waring (1989) noted that woodland studies are especially challenging because of shading and screening by vegetation. Conducting vegetation analysis and pairing sites with similar vegetational composition for light-trap comparison is desirable. Using this method Sample (1991) strengthened his study in use of light traps to determine Dimilin impact on moths as food for the Virginia big-eared bat.
- Rotate traps among sites at each sampling period to avoid bias due to trap differences. Operate light traps simultaneously on as many study sites as possible to avoid the additional variable of different weather conditions on trapping results.
- Use sufficiently large study plots to avoid attracting moths from outside plots. Plaut (1971) noted that some noctuids may be attracted from 200 m. Small plots also allow immigration and emigration of more actively flying moths. Once moths move into a plot from the surrounding matrix, they then may be attracted to traps, thus obscuring treatment effects in the study plots.

With proper precautions, light traps are a useful tool for studying non-target disturbance impacts on moths. We do not use traps to define populations, but need to be concerned only with relative differences among trap sites as affected by treatments.

Summary

A single blacklight trap was operated weekly at Cooper's Rock State Forest, West Virginia from March to October 1984-1986. Four hundred species of Macrolepidoptera were taken. Yearly fluctuation was considerable among many of the species. A review of literature on blacklight trapping studies is presented.

Literature Cited

- Ammar, E.D. 1975. A light trap modified for catching only smaller insects and its efficiency in catching certain groups compared with a sweeping net. *Zeits. Angew. Entomol.* 79:104-109.
- Baker, R.R. and Y.S. Sadovy. 1978. The distance and nature of the light trap response of moths. *Nature*, London: 276:818-821.
- Belton, P. and R.H. Kempster. 1963. Some factors affecting the catches of Lepidoptera in light traps. *Can. Entomol.* 95:832-837.
- Bretherton, R.F. 1954. Moth traps and their lamps: an attempt at comparative analyses. *Entomol. Gaz.* 5:145-154.
- Cantelo, W.W. 1974. Blacklight traps as control agents: an appraisal. *Bull. Entomol. Soc. Amer.* 20: 279-282.
- Cantelo, W.W. 1990. Comparative efficiency of a blacklight trap and a Robinson trap in trapping moths. *Southwest. Entomol.* 15:159-162.
- Cantelo, W.W., J.L. Goodenough, A.H. Baumhover, J.S. Smith, Jr., J.M. Stanley and T.J. Henneberry. 1974. Mass trapping with blacklight: effects on isolated populations of insects. *Environ. Entomol.* 3:389-395.
- Cantelo, W.W. and M. Jacobson. 1979. Phenylacetaldehyde attracts moths to bladder flower and to blacklight traps. *Environ. Entomol.* 8:444-447.
- Carvell, K.L. 1983. A summary of 1973-1982 weather data from the West Virginia University Forest. *W.Va. Forestry Notes* 10:13-16.
- Comstock, J.H. 1879. Fires, trap-lanterns, etc. pp. 262-275. In *Report on Cotton Insects*. USDA Dir. Entomol. Washington, D.C.
- Davidson, A., J.E. Gallagher and H.S. Hsiao. 1973. Reactions of some moths and aquatic insects to light traps fitted with polarizing filters. *Ann. Entomol. Soc. Amer.* 66:232-233.
- Debolt, J.W., W.W. Wolf, T.J. Henneberry and P.V. Vail. 1979. Evaluation of light traps and sex pheromones for control of cabbage looper and other lepidopterous insect pests of lettuce. *USDA Tech. Bull. No. 1606*, 39 pp.
- Dirks, C.O. 1937. Biological studies of Maine moths by light trap methods. *Maine Agric. Expt. Stn. Bull.* 389.
- Frost, S.W. 1957. The Pennsylvania insect light trap. *J. Econ. Entomol.* 50:287-292.
- Frost, S.W. 1964. Insects taken in light traps at the Archbold Biological Station, Highlands County, Florida. *Fla. Entomol.* 47:129-161.

Harding, W.C., Jr., J.G. Hartsock and G.G. Rohwer. 1966. Blacklight trap standards for general insect surveys. *Bull. Entomol. Soc. Amer.* 12:31-32.

Hartstack, A.W. 1979. Light sources, trap design and other factors affecting moth catch. Chap. 15. *In* Movement of Highly Mobile Insects: Concepts and Methodology in Research. Ralb, R.L. and G.G. Kenedy (eds.), 456 pp.

Hartstack, A.W., Jr., J.P. Hollingsworth and D.A. Lindquist. 1968. A technique for measuring trapping efficiency of electric insect traps. *J. Econ. Entomol.* 61:546-552.

Hausmann, A. 1990a. Die Bedeutung des genauen Lichtfallen-Standortes für die Aussagekraft des Fangergebnisses. *Atalanta* 21:301-312.

Hausmann, A. 1990b. Zur Dynamik von Nachfalter Artenspektren Verhalten als Elemente von Verbreitungsstrategien. *Spixiana. Zeits. Zool.* 16, 222 pp.

Heath, J. 1966. A comparison of catches obtained in insect light traps fitted with 16 watt and 6 watt blacklight fluorescent tubes. *Entomol. Record* 78: 222-223.

Hienton, T.E. 1974. Summary of investigations of electric insect traps. USDA tech. Bull 1498, 136 pp.

Hodges, R.W., T. Dominic, D.R. Davis, D.C. Ferguson, J.G. Franclemont, E.G. Monroe and J.A. Powell. 1983. Checklist of the Lepidoptera of America North of Mexico. London. E.W. Classey Ltd. and Washington, D.C. Wedge Entomol. Res. Found. 284 pp.

Hollingsworth, J.P., C.P. Briggs, III, P.A. Glick and H.M. Graham. 1961. Some factors influencing light trap collections. *J. Econ. Entomol.* 54:305-308.

Holloway, J.D. 1984. Moths as indicator organisms for categorizing rain-forest and monitoring changes and regeneration processes. *Tropical Rain Forest: The Leeds Symposium*, pp. 235-242.

King, E.W. and A.T. Hind. 1960. Activity and abundance in insect light trap sampling. *Ann. Entomol. Soc. Amer.* 53:524-529.

Labanowski, G.S. 1980. Catches of moths in the blacklight and mercury-filament traps in an orchard at Skierniewice, Poland. *Pulsk. Pismo Entomol.* 50:473-487.

Magurran, A.E. 1985. The diversity of Macrolepidoptera in two contrasting woodland habitats at Banagher, Northern Ireland. *Proc. Roy. Irish Acad., Sect. B, Biol., Geol. and Chem. Sci.* 85:121-132.

- McGeachie, W.J. 1988. A remote sensing method for the estimation of light trap efficiency. *Bull. Entomol. Res.* 78:379–385.
- McGeachie, W.J. 1989. The effects of moonlight illuminance, temperature and wind speed on light-trap catches of moths. *Bull. Entomol. Res.* 79:185–192.
- Mizutani, M. 1982. Optimum setting position of the light trap for the collection of moths (Lepidoptera). *Jap. J. Appl. Entomol. Zool.* 26:58–62.
- Mizutani, M., J. Mishima and Y. Kobayashi. 1982. The electrode type light trap for the collection of moths. *App. Entomol. and Zool.* 17:172–178.
- Moulding, J.D. and J.J. Madenjian. 1979. Macrolepidopteran moths light-trapped in a New Jersey oak forest (Lepidoptera). *Proc. Entomol. Soc. Wash.* 81:135–144.
- Muirhead-Thomson, R.C. 1991. Trap responses of flying insects. The influence of trap design on capture efficiency. Academic Press, San Diego. 304 pp.
- Novak, I. 1983. An efficient light-trap for catching insects. *Acta Entomol. Bohemoslovica.* 80:29–34.
- Peacock, J.W. and S.W. Bullington. 1989. Evaluation of the impact of *Bacillus thuringiensis* on non-target Lepidoptera in the Shenandoah National Park. *Proc. 1989 Nat. Gypsy Moth Rev.* Nov. 6–9, 1989, Annapolis.
- Plaut, H.N. 1971. Distance of attraction of moths of *Spodoptera littoralis* to BL radiation, and recapture of moths released at different distance of an ESA blacklight standard trap. *J. Econ. Entomol.* 64:1402–1404.
- Rejmanek, M. and K. Spitzer. 1982. Bionomic strategies and long-term fluctuations in abundance of Noctuidae (Lepidoptera). *Acta Entomol. Bohemoslovica* 79:81–96.
- Rings, R.W. and E.H. Metzler. 1988. Preliminary annotated checklist of the Lepidoptera of Atwood Lake Park, Ohio. *Ohio J. Sci.* 88:59–168.
- Rings, R.W. and E.H. Metzler. 1989. A preliminary checklist of the Lepidoptera of Mohican State Forest and Mohican State Park, Ashland County, Ohio. *Ohio J. Sci.* 89:78–88.
- Rings, R.W. and E.H. Metzler. 1990. The Lepidoptera of Fowler Woods State Nature Preserve, Richland County, Ohio. *The Great Lakes Entomol.* 23:43–56.

Rings, R.W., R.M. Rittles, R.W. Hawes and E.H. Metzler. 1987. A nine-year study of the Lepidoptera of the Wilderness Center, Stark County, Ohio. *Ohio J. Sci.* 87:55–61.

Robinson, H.S. and P.J.M. Robinson. 1950. Some notes on the observed behavior of Lepidoptera in flight in the vicinity of light-trap sources together with a description of a light trap designed to take entomological samples. *Ent. Gaz.* 1:3–20.

Sample, Bradley E. 1991. Effects of Dimilin on food of the endangered Virginia big-eared bat. Ph.D. Dissertation, West Virginia University. 201 pp.

Sargent, T.D. 1976. *Legion of Night – The Underwing Moths*. Univ. Massachusetts Press, Amherst. 222 pp.

Smith, J.S., Jr., J.M. Stanley, J.G. Hartsock and L.E. Campbell. 1974. S-1 Black-light insect-survey trap, plans and specifications. USDA, ARS-S-31, 8 pp.

Smith, P.W. 1962. The use of blacklight insect traps as an entomological tool. *Proc. N. Central Branch – E.S.A.* 17:38–39.

Smith, P.W., J.G. Taylor and J.W. Apple. 1959. A comparison of insect traps equipped with 6- and 15-watt blacklight lamps. *J. Econ. Entomol.* 52:1212–1214.

Southwood, T.R. E. 1978. *Ecological methods with particular reference to the study of insect populations*. 2nd Edition. Wiley & Sons, New York. 524 pp.

Spitzer, K. and J. Leps. 1988. Determinants of temporal variation in moth abundance. *Oikos* 53:31–36.

Stewart, P.A., J.J. Lam, Jr. and J.L. Blythe. 1969. Influence of distance on attraction of tobacco hornworm and corn earworm moths to radiations of a blacklight lamp. *J. Econ. Entomol.* 62:58–60.

Taylor, L.R. 1979. The Rothamsted insect survey – an approach to the theory and practice of synoptic pest forecasting in agriculture. *In* *Movement of Highly Mobile Insects: Concepts and Methodology in Research*. R.L. Rabb and G.G. Kennedy, Eds. pp. 148–185.

Taylor, L.R. and E.S. Brown. 1972. Effects of light-trap design and illumination on samples of moths in the Kenya highlands. *Bull. Entomol. Res.* 62:91–112.

Taylor, L.R. and C.I. Carter. 1961. The analysis of numbers and distribution in an aerial population of Macrolepidoptera. Trans. Roy. Entomol. Soc. London 113: 369–386.

Taylor, L.R. and R.A. French. 1974. Effects of light-trap design and illumination on samples of moths in an English woodland. Bull. Entomol. Res. 63:593–594.

Taylor, L.R., R.A. French and I.P. Woiwod. 1978. The Rothamsted insect survey and the urbanization of land in Great Britain, pp. 31–66. In Frankie, G.W. and C.S. Koehler (eds.), perspectives in urban entomology. Academic Press, New York. 417 pp.

Thomas, C.D. 1989. Limits and scope of light-trapping for studying moth population dynamics. New Zealand Entomol. 12:89–90.

Vaishampayan, S.M. and R. Verma. 1983. Comparative efficiency of various light sources in trapping adults of *Heliothis armigera* (Hubn.), *Spodoptera litura* (Boisd.) and *Agrotis ipsilon* (Hufn.) (Lepidoptera: Noctuidae). Indian J. Agric. Sci. 53:163–167.

Waring, P. 1980. A comparison of the Heath and Robinson M.V. traps. Entomol. Record 92:283–289.

Waring, P. 1989. Comparison of light-trap catches in deciduous and coniferous woodland habitats. Entomol. Record 101:1–10.

Watson, J.K. and P.L. Lambdin. 1989. Diversity of lepidopteran fauna collected from selected sites throughout the Great Smoky Mountains National Park. Proc. 1989. Nat. Gypsy Moth Rev. Nov. 6–9, 1989, Annapolis.

White, E.G. 1989. Light-trapping frequency and data analysis – a reply. New Zealand Entomol. 12:91–94.

Williams, C.B. 1939. An analysis of four years of captures of insects in a light trap. Part I. General survey; sex proportion; phenology; and time of flight. Trans. Roy. Entomol. Soc. London 87:79–131.

Williams, C.B. 1940. An analysis of four years captures of insects in a light trap. Part II. The effect of weather conditions on insect population. Trans. Roy. Entomol. Soc. London 90:227–306.

Williams, C.B., R.A. French and M.M. Hosni. 1955. A second experiment on testing the relative efficiency of insect traps. Bull. Entomol. Res. 46:193–204.

TABLE 1.

Number of species of Macrolepidoptera in all families taken by blacklight trap at Cooper's Rock State Forest, West Virginia (1984-1986) during any one, two or three years of the study.

FAMILY	# of YEARS Species Trapped			TOTAL SPECIES
	1	2	3	
THYATIRIDAE			3	3
DREPANIDAE			2	2
GEOMETRIDAE	20	22	60	102
EPIPLEMIDAE			1	1
MIMALLONIDAE		1	1	2
APATELODIDAE	1		1	2
LASIOCAMPIDAE		1	3	4
SATURNIIDAE		2	4	6
SPHINGIDAE	1	3	2	6
NOTODONTIDAE	10	2	19	31
ARCTIIDAE	6	5	6	17
LYMANTRIIDAE	1	1	2	4
NOCTUIDAE	77	62	81	220
TOTAL SPECIES	116	99	185	400

TABLE 2.

Macrolepidoptera species collected by blacklight trap at Cooper's Rock State Forest, West Virginia during 1984-1986. Total catch numbers, rank and inclusive flight range are given.

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
THYATIRIDAE				
Habrosyne scripta (Gosse)	2(215)	14(102)	5(134)	24MAY-15AUG
Pseudothyatira cymatophoroides (Gn.)	4(179)	3(179)	11(95)	29MAY-13SEP
Euthyatira pudens (Gn.)	2(215)	12(111)	2(184)	15APR-12MAY
DREPANIDAE				
Drepana arcuata Wlk.	11(130)	6(151)	22(64)	10MAY-11SEP
Oreta rosea (Wlk.)	18(95)	8(131)	4(147)	30MAY-24SEP
GEOMETRIDAE				
Helimata cycladata G. & R.	4(179)	3(179)	6(124)	05JUN-28JUN
Itame pustularia (Gn.)	949(1)	588(3)	337(5)	12JUN-11SEP
Semiothisa aemulataria (Wlk.)	89(29)	101(27)	30(58)	02MAY-11SEP
Semiothisa ulsterata (Pears.)	44(61)	96(28)	16(76)	10MAY-15AUG
Semiothisa pinistrobata Fgn.	3(191)	0(.)	57(39)	06JUN-14AUG
Semiothisa ocellinata (Gn.)	25(83)	1(224)	8(106)	12MAY-21AUG
Enconista dislocaria (Pack.)	13(115)	8(131)	3(161)	10MAY-13JUN
Glena cribrataria (Gn.)	18(95)	23(78)	65(34)	29MAY-06JUL
Aethalura intertexta (Wlk.)	1(242)	6(151)	6(124)	02MAY-17JUL
Iridopsis larvaria (Gn.)	165(15)	195(9)	119(20)	02MAY-19SEP
Anavitrinella pampinaria (Gn.)	15(104)	7(141)	7(116)	06MAY-21SEP
Cleora sublunaria (Gn.)	2(215)	1(224)	0(.)	10APR-12MAY
Ectropis crepuscularia (D. & S.)	0(.)	73(36)	43(46)	25APR-28AUG
Protoboarmia porcelaria (Gn.)	54(50)	6(151)	0(.)	12MAY-10SEP
Epimecis hortaria (F.)	50(54)	120(21)	127(18)	25APR-24SEP
Melanolophia canadaria (Gn.)	497(4)	624(2)	553(3)	08APR-28OCT
Biston betularia (L.)	70(41)	63(42)	22(64)	19APR-11SEP
Hypagyrtis unipunctata (Haw.)	60(46)	40(55)	37(48)	05JUN-24SEP
Phigalia titea (Cram.)	11(130)	12(111)	15(79)	28MAR-27APR
Paleacrita vernata (Peck)	1(242)	3(179)	5(134)	19MAR-12APR
Paleacrita merriccata Dyar	0(.)	0(.)	3(161)	19MAR-19MAR
Erannis tiliaria (Harr.)	3(191)	2(199)	3(161)	28OCT-06NOV
Lomographa semiclarata (Wlk.)	2(215)	4(163)	1(212)	12MAY-10JUL
Lomographa vestaliata (Gn.)	18(95)	133(18)	236(8)	10MAY-07AUG
Lomographa glomeraria (Grt.)	519(3)	538(4)	584(1)	02APR-25MAY
Cabera erythemaria Gn.	0(.)	15(96)	16(76)	16MAY-30JUL
Cabera quadrifasciaria (Pack.)	0(.)	1(224)	0(.)	10JUL-10JUL
Lytrosis unitaria (H.-S.)	10(136)	3(179)	27(60)	21JUN-18JUL
Euchlaena serrata (Drury)	0(.)	4(163)	5(134)	06JUN-27JUN
Euchlaena obtusaria (Hbn.)	15(104)	1(224)	0(.)	22JUN-10SEP

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Euchlaena johnsonaria</i> (Fitch)	0(.)	1(224)	0(.)	05JUN-05JUN
<i>Euchlaena marginaria</i> (Minot)	5(166)	2(199)	0(.)	10MAY-08JUN
<i>Euchlaena tigrinaria</i> (Gn.)	10(136)	7(141)	3(161)	05JUN-28JUN
<i>Euchlaena irraria</i> (B. & McD.)	10(136)	2(199)	2(184)	12JUN-28JUN
<i>Xanthotype sospeta</i> (Drury)	1(242)	1(224)	2(184)	05JUN-05SEP
<i>Pero honestaria</i> (Wlk.)	89(29)	23(78)	0(.)	25APR-23AUG
<i>Pero morrisonaria</i> (Hy. Edw.)	22(88)	73(36)	29(59)	05JUN-30JUL
<i>Nacophora quernaria</i> (J.E. Smith)	13(115)	16(94)	22(64)	16MAY-27JUL
<i>Campaea perlata</i> (Gn.)	153(16)	121(20)	461(4)	29MAY-02OCT
<i>Ennomos magnaria</i> Gn.	20(91)	45(51)	48(43)	24SEP-22OCT
<i>Homochlodes disconventa</i> (Wlk.)	108(24)	52(47)	64(36)	15APR-01OCT
<i>Selenia kentaria</i> (G. & R.)	3(191)	9(125)	0(.)	19APR-08AUG
<i>Metanema inatomaria</i> (Gn.)	0(.)	0(.)	2(184)	06JUN-10JUL
<i>Metarranthis duaria</i> (Gn.)	7(154)	10(120)	5(134)	19APR-08JUN
<i>Metarranthis angularia</i> (B. & McD.)	0(.)	0(.)	7(116)	06JUN-12JUN
<i>Metarranthis hypocharia</i> (H.-S.)	75(36)	49(49)	1(212)	23MAY-28JUN
<i>Metarranthis homuraria</i> (G. & R.)	0(.)	0(.)	18(73)	06JUN-20JUN
<i>Cepphis armataria</i> (H. & S.)	5(166)	4(163)	12(91)	12JUN-30JUL
<i>Anagoga occiduaria</i> (Wlk.)	73(40)	9(125)	3(161)	25APR-15AUG
<i>Probole alienaria</i> (H.-S.)	80(33)	15(96)	0(.)	10MAY-06JUL
<i>Probole amicarica</i> (H.-S.)	133(19)	1073(1)	580(2)	06MAY-23AUG
<i>Plagodis serinaria</i> H.-S.	74(39)	173(12)	77(32)	16MAY-28JUN
<i>Plagodis phlogosaria</i> (Gn.)	24(84)	5(157)	3(161)	10MAY-08AUG
<i>Plagodis fervidaria</i> (H.S.)	8(149)	2(199)	0(.)	03JUN-08AUG
<i>Plagodis alchoolaria</i> (Gn.)	111(23)	186(11)	32(54)	10MAY-13AUG
<i>Caripeta divisata</i> Wlk.	1(242)	0(.)	1(212)	27JUL-07AUG
<i>Besma endropiaria</i> (G. & R.)	75(36)	90(30)	89(27)	24MAY-14AUG
<i>Besma quercivoraria</i> (Gn.)	279(7)	114(23)	25(62)	29APR-24SEP
<i>Lambdina fiscellaria</i> (Gn.)	15(104)	8(131)	0(.)	06SEP-17OCT
<i>Lambdina pellucidaria</i> (G. & R.)	3(191)	23(78)	13(83)	25APR-13AUG
<i>Lambdina fervidaria</i> (Hbn.)	105(26)	10(120)	0(.)	08MAY-29AUG
<i>Eusarca confusaria</i> Hbn.	1(242)	1(224)	2(184)	06JUN-21SEP
<i>Tetracis crocallata</i> Gn.	1(242)	1(224)	0(.)	05JUN-06JUL
<i>Tetracis cachexiata</i> Gn.	124(21)	217(7)	180(11)	16MAY-06JUL
<i>Eugonobapta nivosaria</i> (Gn.)	12(122)	12(111)	8(106)	02JUL-19JUL
<i>Eutrapela clemataria</i> (J.E. Smith)	29(75)	29(67)	8(106)	08APR-08AUG
<i>Prochoerodes transversata</i> (Drury)	36(67)	13(107)	62(37)	02JUL-22OCT
<i>Antepione thisoaria</i> (Gn.)	12(122)	0(.)	3(161)	25MAY-15AUG
<i>Nematocampa limbata</i> (Haw.)	3(191)	4(163)	5(134)	27JUN-24SEP
<i>Nemoria lixaria</i> (Gn.)	0(.)	1(224)	0(.)	19APR-19APR
<i>Nemoria bistriaria</i> Hbn.	26(80)	24(76)	2(184)	25APR-29AUG
<i>Nemoria mimosaria</i> (Gn.)	0(.)	8(131)	0(.)	13AUG-13AUG
<i>Pleuroprucha insulsaria</i> (Gn.)	4(179)	7(141)	8(106)	20JUN-25OCT
<i>Pleuroprucha asthenaria</i> (Wlk.)	13(115)	0(.)	0(.)	21SEP-19OCT
<i>Cyclophora packardi</i> (Prout)	3(191)	7(141)	3(161)	10MAY-02OCT

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Cyclophora pendulinaria</i> (Gn.)	27(78)	28(69)	9(105)	02MAY-29AUG
<i>Scopula limboundata</i> (Haw.)	91(28)	25(75)	19(71)	02JUL-26SEP
<i>Dysstroma hersiliata</i> (Gn.)	0(.)	0(.)	1(212)	10JUL-10JUL
<i>Eulithis diversilineata</i> (Hbn.)	11(130)	6(151)	5(134)	06JUL-26SEP
<i>Eulithis explanata</i> (Wlk.)	2(215)	0(.)	0(.)	06JUL-27JUL
<i>Hydriomena divisaria</i> (Wlk.)	69(42)	17(92)	0(.)	25APR-03JUN
<i>Hydria prunivorata</i> (Fgn.)	13(115)	11(118)	5(134)	12JUN-19JUL
<i>Mesoleuca ruficillata</i> (Gn.)	0(.)	1(224)	0(.)	05JUN-05JUN
<i>Anticlea vasilata</i> Gn.	2(215)	3(179)	1(212)	25APR-16MAY
<i>Xanthorhoe ferrugata</i> (Cl.)	0(.)	1(224)	0(.)	02MAY-02MAY
<i>Xanthorhoe lacustrata</i> (Gn.)	5(166)	7(141)	0(.)	10MAY-10SEP
<i>Epirrhoe alternata</i> (Mu"ller)	0(.)	0(.)	1(212)	12JUN-12JUN
<i>Euphyia unangulata</i> (Haw.)	0(.)	0(.)	1(212)	21AUG-21AUG
<i>Orthonama obstipata</i> (F.)	2(215)	2(199)	6(124)	06MAY-28AUG
<i>Orthonama centrostrigaria</i> (Woll.)	5(166)	7(141)	11(95)	05JUN-17OCT
<i>Hydrelia inornata</i> (Hulst)	41(64)	135(17)	147(14)	16MAY-09SEP
<i>Venusia cambrica</i> Curt.	0(.)	2(199)	0(.)	05JUN-19JUL
<i>Trichodezia albiovittata</i> (Gn.)	0(.)	3(179)	0(.)	10MAY-30MAY
<i>Eubaphe mendica</i> (Wlk.)	3(191)	1(224)	0(.)	30MAY-15AUG
<i>Horisme intestinata</i> (Gn.)	0(.)	1(224)	1(212)	12JUN-13AUG
<i>Eupithecia miserulata</i> Grt.	208(10)	163(14)	84(29)	28MAR-21SEP
<i>Cladara limitaria</i> (Wlk.)	5(166)	0(.)	0(.)	08MAY-12MAY
<i>Cladara atroliturala</i> (Wlk.)	4(179)	19(84)	4(147)	08APR-16MAY
<i>Lobophora nivigerata</i> Wlk.	0(.)	2(199)	0(.)	10MAY-10MAY
<i>Heterophleps refusaria</i> (Wlk.)	0(.)	0(.)	1(212)	17JUL-17JUL
<i>Heterophleps triguttaria</i> H.-S.	0(.)	1(224)	4(147)	12JUN-06AUG
<i>Dyspteris abortivaria</i> (H.-S.)	4(179)	1(224)	2(184)	08JUN-30JUL
EPIPLEMIDAE				
<i>Calledapteryx dryopterata</i> Grt.	1(242)	1(224)	3(161)	30MAY-07AUG
MIMALLONIDAE				
<i>Lacosoma chiridota</i> Grt.	1(242)	3(179)	0(.)	05JUN-28JUN
<i>Cicinnus melsheimeri</i> (Harr.)	7(154)	27(72)	17(74)	05JUN-28JUN
APATELODIDAE				
<i>Apatelodes torrefacta</i> (J.E. Smith)	23(85)	4(163)	12(91)	12JUN-08AUG
<i>Olceclostera angelica</i> (Grt.)	1(242)	0(.)	0(.)	06JUL-06JUL
LASIOCAMPIDAE				
<i>Tolyte velleda</i> (Stoll)	36(67)	14(102)	2(184)	10SEP-01OCT
<i>Phyllodesma americana</i> (Harr.)	2(215)	5(157)	0(.)	19APR-12MAY
<i>Malacosoma distria</i> Hbn.	108(24)	36(60)	146(15)	20JUN-18JUL
<i>Malacosoma americanum</i> (F.)	407(5)	162(15)	193(10)	05JUN-27JUL

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
SATURNIIDAE				
Eacles imperialis (Drury)	1(242)	0(.)	3(161)	17JUL-03AUG
Dryocampa rubicunda (F.)	23(85)	4(163)	6(124)	16MAY-08AUG
Automeris io (F.)	16(101)	13(107)	7(116)	24MAY-10JUL
Antheraea polyphemus (Cram.)	43(62)	13(107)	7(116)	10MAY-13AUG
Actias luna (L.)	3(191)	6(151)	5(134)	02MAY-08AUG
Callosamia angulifera (Wlk.)	2(215)	0(.)	6(124)	10JUL-08AUG
SPHINGIDAE				
Manduca jaminearum (Guer.)	1(242)	1(224)	0(.)	10JUL-03AUG
Ceratomia undulosa (Wlk.)	1(242)	0(.)	2(184)	10JUL-07AUG
Paonias excaecatus (J.E. Smith)	31(74)	57(45)	78(31)	30MAY-08AUG
Paonias myops (J.E. Smith)	64(44)	44(52)	88(28)	16MAY-15AUG
Laothoe juglandis (J.E. Smith)	1(242)	0(.)	4(147)	02JUL-19JUL
Pachysphinx modesta (Harr.)	0(.)	0(.)	1(212)	10JUL-10JUL
NOTODONTIDAE				
Clostera albosigma Fitch	1(242)	0(.)	0(.)	03JUN-03JUN
Clostera inclusa (Hbn.)	3(191)	0(.)	0(.)	03JUN-03AUG
Datana ministra (Drury)	9(142)	24(76)	11(95)	05JUN-03AUG
Datana angusii G.& R.	0(.)	0(.)	5(134)	06JUN-17JUL
Datana drexelii Hy. Edw.	1(242)	0(.)	0(.)	27JUL-27JUL
Datana integerrima G.& R.	1(242)	0(.)	0(.)	22JUN-22JUN
Nadata gibbosa (J.E. Smith)	77(35)	84(33)	240(7)	06MAY-26SEP
Hyperaeschra georgica (H.-S.)	10(136)	14(102)	1(212)	02MAY-27JUL
Peridea basitriens (Wlk.)	1(242)	2(199)	2(184)	10JUL-27JUL
Peridea angulosa (J.E. Smith)	192(12)	59(44)	118(22)	12JUN-28AUG
Peridea ferruginea (Pack.)	5(166)	1(224)	0(.)	05JUN-08AUG
Pheosia rimosa Pack.	1(242)	0(.)	0(.)	08AUG-08AUG
Odontosis elegans (Stkr.)	1(242)	3(179)	5(134)	12JUN-03AUG
Nerice bidentata Wlk.	2(215)	0(.)	0(.)	08AUG-23AUG
Ellida caniplaga (Wlk.)	0(.)	5(157)	0(.)	19APR-10MAY
Gluphisia septentrionis Wlk.	2(215)	0(.)	0(.)	08AUG-08AUG
Furcula borealis (Guer.-Meneville)	1(242)	1(224)	1(212)	10MAY-30JUL
Symmerista leucitys Franc.	48(58)	73(36)	119(20)	29MAY-03AUG
Dasylophia anguina (J.E. Smith)	9(142)	4(163)	3(161)	10MAY-17SEP
Macrurocampa marthesia (Cram.)	52(52)	89(31)	110(24)	12JUN-14AUG
Heterocampa obliqua Pack.	9(142)	7(141)	10(103)	30MAY-15AUG
Heterocampa umbrata Wlk.	3(191)	4(163)	4(147)	10MAY-13AUG
Heterocampa guttivitta (Wlk.)	61(45)	201(8)	297(6)	02MAY-13AUG
Heterocampa biundata Wlk.	23(85)	43(53)	113(23)	24MAY-05SEP
Lochmaeus manteo Doubleday	20(91)	37(58)	60(38)	05JUN-14AUG
Lochmaeus bilineata (Pack.)	0(.)	0(.)	1(212)	10JUL-10JUL
Schizura ipomoeae Doubleday	7(154)	17(92)	35(49)	06JUN-05SEP
Schizura unicornis (J.E. Smith)	6(160)	7(141)	0(.)	05JUN-15AUG

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Schizura leptinoides</i> (Grt.)	1(242)	1(224)	1(212)	05JUN-27JUL
<i>Oligocentria semirufescens</i> (Wlk.)	49(56)	18(90)	32(54)	05JUN-09SEP
<i>Oligocentria lignicolor</i> (Wlk.)	12(122)	2(199)	1(212)	28JUN-23AUG

ARCTIIDAE

<i>Hypoprepia miniata</i> (Kby.)	14(109)	0(.)	13(83)	10JUL-23JUL
<i>Hypoprepia fucosa</i> Hbn.	21(90)	14(102)	0(.)	10JUL-15AUG
<i>Clemensia albata</i> Pack.	83(32)	312(5)	200(9)	29MAY-11OCT
<i>Haploa clymene</i> (Brown)	1(242)	0(.)	0(.)	08AUG-08AUG
<i>Holomelina opella</i> (Grt.)	0(.)	0(.)	1(212)	20JUN-20JUN
<i>Pyrrharctia isabella</i> (J.E. Smith)	3(191)	0(.)	0(.)	28JUN-28JUN
<i>Spilosoma latipennis</i> Stretch	1(242)	0(.)	1(212)	12JUN-12JUN
<i>Spilosoma virginica</i> (F.)	3(191)	2(199)	1(212)	10MAY-21AUG
<i>Hyphantria cunea</i> (Drury)	153(16)	169(13)	13(83)	10MAY-18JUL
<i>Apantesis phalerata</i> (Harr.)	5(166)	1(224)	0(.)	10MAY-28JUN
<i>Apantesis nais</i> (Drury)	0(.)	0(.)	1(212)	16MAY-16MAY
<i>Apantesis figurata</i> (Drury)	1(242)	0(.)	0(.)	28JUN-28JUN
<i>Halysidota tessellaris</i> (J.E. Smith)	698(2)	149(16)	144(16)	30MAY-15AUG
<i>Lophocampa caryae</i> Harr.	2(215)	63(42)	34(51)	29MAY-28JUN
<i>Cycnia tenera</i> Hbn.	1(242)	2(199)	2(184)	12JUN-15AUG
<i>Euchaetes egle</i> (Drury)	3(191)	0(.)	1(212)	28JUN-17JUL
<i>Cisseps fulvicollis</i> (Hbn.)	1(242)	0(.)	0(.)	28JUN-28JUN

LYMANTRIIDAE

<i>Dasychira dorsipennata</i> (B. & McD.)	0(.)	0(.)	67(33)	20JUN-23JUL
<i>Dasychira obliquata</i> (G. & R.)	0(.)	4(163)	1(212)	23JUL-13AUG
<i>Dasychira</i> spp.	75(36)	47(50)	53(40)	12JUN-24SEP
<i>Orgyia leucostigma</i> (J.E. Smith)	173(14)	37(58)	163(13)	28JUN-30OCT

NOCTUIDAE

<i>Idia americalis</i> (Gn.)	15(104)	12(111)	32(54)	16MAY-19OCT
<i>Idia aemula</i> Hbn.	346(6)	38(57)	45(45)	12JUN-30OCT
<i>Idia rotundalis</i> (Wlk.)	205(11)	231(6)	65(34)	10JUL-21SEP
<i>Idia diminuentis</i> (B. & McD.)	0(.)	3(179)	2(184)	23JUL-13AUG
<i>Idia scobialis</i> (Grt.)	4(179)	3(179)	4(147)	10JUL-21SEP
<i>Idia lubricalis</i> (Gey.)	15(104)	0(.)	2(184)	13JUL-29AUG
<i>Phalaenophana pyramusalis</i> (Wlk.)	9(142)	4(163)	13(83)	30MAY-11SEP
<i>Zanclognatha lituralis</i> (Hbn.)	8(149)	39(56)	19(71)	27JUN-24JUL
<i>Zanclognatha laevigata</i> (Grt.)	144(18)	41(54)	48(43)	20JUN-06SEP
<i>Zanclognatha pedipilalis</i> (Gn.)	13(115)	0(.)	3(161)	20JUN-13SEP
<i>Zanclognatha protumnusalis</i> (Wlk.)	7(154)	0(.)	0(.)	03AUG-15AUG
<i>Zanclognatha cruralis</i> (Gn.)	1(242)	12(111)	0(.)	05JUN-19JUL
<i>Zanclognatha jacchusalis</i> (Wlk.)	5(166)	0(.)	0(.)	28JUN-08AUG
<i>Zanclognatha ochreipennis</i> (Grt.)	214(9)	70(39)	42(47)	20JUN-02OCT
<i>Chytolita morbidalis</i> (Gn.)	46(60)	126(19)	31(57)	29MAY-13JUL

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Phalaenostola metonalis</i> (Wlk.)	3(191)	0(.)	3(161)	12JUN-24SEP
<i>Phalaenostola larentioides</i> Grt.	9(142)	1(224)	3(161)	03JUN-23AUG
<i>Tetanolita mynesalis</i> (Wlk.)	0(.)	0(.)	4(147)	17JUL-30JUL
<i>Bleptina caradrinalis</i> Gn.	2(215)	0(.)	0(.)	03JUN-08JUN
<i>Renia factiosalis</i> (Wlk.)	43(62)	0(.)	0(.)	03AUG-15AUG
<i>Renia nemoralis</i> B. & McD.	0(.)	0(.)	1(212)	23JUL-23JUL
<i>Renia discoloralis</i> Gn.	1(242)	31(66)	5(134)	06JUL-06SEP
<i>Renia sobrialis</i> (Wlk.)	4(179)	3(179)	0(.)	06AUG-15AUG
<i>Palthis angularis</i> (Hbn.)	8(149)	0(.)	2(184)	08JUN-15AUG
<i>Palthis asopialis</i> (Gn.)	0(.)	0(.)	1(212)	24SEP-24SEP
<i>Rivula propinqualis</i> Gn.	0(.)	3(179)	1(212)	05JUN-06SEP
<i>Bomolocha baltimoralis</i> (Gn.)	116(22)	104(25)	107(25)	29APR-26SEP
<i>Bomolocha palparia</i> (Wlk.)	1(242)	0(.)	0(.)	13JUL-13JUL
<i>Bomolocha abalienalis</i> (Wlk.)	2(215)	0(.)	0(.)	06JUL-03AUG
<i>Bomolocha deceptalis</i> (Wlk.)	0(.)	5(157)	1(212)	06MAY-05JUN
<i>Bomolocha madefactalis</i> (Gn.)	0(.)	1(224)	0(.)	05JUN-05JUN
<i>Bomolocha edictalis</i> (Wlk.)	7(154)	1(224)	2(184)	03JUL-08AUG
<i>Hypena humuli</i> Harr.	1(242)	2(199)	0(.)	25APR-03AUG
<i>Plathypena scabra</i> (F.)	12(122)	8(131)	7(116)	23AUG-30OCT
<i>Pangrapta decoralis</i> Hbn.	20(91)	28(69)	21(68)	23MAY-09SEP
<i>Metalectra discalis</i> (Grt.)	0(.)	2(199)	2(184)	12JUN-13AUG
<i>Scolecocampa liburna</i> (Gey.)	2(215)	0(.)	3(161)	17JUL-23AUG
<i>Scoliopteryx libatrix</i> (L.)	0(.)	1(224)	0(.)	10JUL-10JUL
<i>Anticarsia gemmatilis</i> Hbn.	0(.)	0(.)	2(184)	02OCT-02OCT
<i>Panopoda rufimargo</i> (Hbn.)	26(80)	2(199)	6(124)	28JUN-08AUG
<i>Phoberia atomaris</i> Hbn.	33(71)	13(107)	3(161)	08APR-25MAY
<i>Cissusa spadix</i> (Cram.)	7(154)	8(131)	0(.)	04MAY-25MAY
<i>Zale lunata</i> (Drury)	1(242)	0(.)	1(212)	10OCT-28OCT
<i>Zale galbanata</i> (Morr.)	1(242)	0(.)	0(.)	08MAY-08MAY
<i>Zale aeruginosa</i> (Gn.)	1(242)	0(.)	0(.)	22JUN-22JUN
<i>Zale undularis</i> (Drury)	1(242)	0(.)	0(.)	22JUN-22JUN
<i>Zale minerea</i> (Gn.)	53(51)	85(32)	13(83)	08APR-08AUG
<i>Zale lunifera</i> (Hbn.)	12(122)	82(34)	13(83)	02APR-03JUN
<i>Zale unilineata</i> (Grt.)	4(179)	8(131)	0(.)	10MAY-25MAY
<i>Euparthenos nubilis</i> (Hbn.)	5(166)	0(.)	2(184)	16MAY-29AUG
<i>Allotria elonympha</i> (Hbn.)	14(109)	4(163)	4(147)	05JUN-26SEP
<i>Parallelia bistriaris</i> Hbn.	101(27)	19(84)	21(68)	10MAY-10SEP
<i>Euclidia cuspidata</i> (Hbn.)	0(.)	1(224)	0(.)	05JUN-05JUN
<i>Caenurgina erechthea</i> (Cram.)	1(242)	0(.)	0(.)	28JUN-28JUN
<i>Catocala resecta</i> Grt.	4(179)	1(224)	4(147)	28AUG-02OCT
<i>Catocala vidua</i> (J.E. Smith)	1(242)	0(.)	0(.)	29AUG-29AUG
<i>Catocala palaeogama</i> Gn.	2(215)	0(.)	0(.)	23AUG-23AUG
<i>Catocala subnata</i> Grt.	0(.)	1(224)	0(.)	23AUG-23AUG
<i>Catocala ilia</i> (Cram.)	6(160)	19(84)	10(103)	03JUL-26SEP
<i>Catocala relictata</i> Wlk.	1(242)	0(.)	1(212)	28AUG-26SEP

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
Catocala sordida Grt.	6(160)	0(.)	0(.)	03AUG-08AUG
Catocala gracilis Edw.	0(.)	2(199)	0(.)	19JUL-13AUG
Catocala andromedae Gn.	3(191)	0(.)	1(212)	30JUL-08AUG
Catocala ultronia (Hbn.)	12(122)	27(72)	11(95)	17JUL-01OCT
Catocala grynea (Cram.)	1(242)	1(224)	0(.)	15AUG-06SEP
Catocala blandula Hulst	2(215)	0(.)	2(184)	06JUL-08AUG
Catocala similis Edw.	1(242)	0(.)	1(212)	17JUL-03AUG
Catocala micronympha Gn.	2(215)	1(224)	0(.)	27JUL-15AUG
Catocala amica (Hbn.)	26(80)	0(.)	2(184)	23JUL-10SEP
Allagrapha aerea (Hbn.)	1(242)	0(.)	0(.)	03AUG-03AUG
Chrysanypha formosa (Grt.)	3(191)	6(151)	1(212)	05JUN-18JUL
Eosporopteryx thyatroides (Gn.)	1(242)	1(224)	1(212)	03AUG-24SEP
Autographa precatonis (Gn.)	3(191)	2(199)	6(124)	03JUN-24SEP
Autographa ampla (Wlk.)	0(.)	1(224)	0(.)	06AUG-06AUG
Anagrapha falcifera (Kby.)	1(242)	3(179)	0(.)	05JUN-28JUN
Paectes oculatrix (Gn.)	1(242)	0(.)	0(.)	15AUG-15AUG
Baileya doubledayi (Gn.)	1(242)	0(.)	0(.)	03AUG-03AUG
Baileya ophthalmica (Gn.)	0(.)	14(102)	2(184)	10MAY-06JUN
Baileya levitans (Sm.)	14(109)	0(.)	4(147)	08MAY-27JUL
Meganola minuscula (Zell.)	22(88)	7(141)	0(.)	25APR-25MAY
Nola triquetra (Fitch)	0(.)	0(.)	11(95)	02APR-30JUL
Hyperstrotia pervertens (B. & McD.)	0(.)	1(224)	2(184)	17JUL-06AUG
Hyperstrotia secta (Grt.)	1(242)	15(96)	51(42)	20JUN-03AUG
Lithacodia muscosa (Gn.)	3(191)	0(.)	3(161)	12JUN-29AUG
Lithacodia albidula (Gn.)	1(242)	0(.)	0(.)	08AUG-08AUG
Lithacodia synochitis (G. & R.)	0(.)	0(.)	1(212)	02JUL-02JUL
Lithacodia musta (G. & R.)	0(.)	0(.)	4(147)	27JUN-17JUL
Lithacodia carneola (Gn.)	5(166)	1(224)	1(212)	05JUN-03AUG
Cerma cerintha (Tr.)	18(95)	3(179)	3(161)	12JUN-08AUG
Tarachidia candefacta (Hbn.)	1(242)	0(.)	0(.)	28JUN-28JUN
Colocasia propinquinella (Grt.)	13(115)	15(96)	5(134)	19APR-15AUG
Charadra deridens (Gn.)	2(215)	0(.)	0(.)	25MAY-27JUL
Raphia frater Grt.	3(191)	0(.)	0(.)	22JUN-13JUL
Acronicta americana (Harr.)	12(122)	5(157)	12(91)	12JUN-07AUG
Acronicta dactylina Grt.	1(242)	0(.)	0(.)	22JUN-22JUN
Acronicta innotata Gn.	5(166)	0(.)	6(124)	27JUN-08AUG
Acronicta radcliffei (Harv.)	0(.)	2(199)	1(212)	25APR-23AUG
Acronicta funeralis G. & R.	0(.)	1(224)	0(.)	10MAY-10MAY
Acronicta hasta Gn.	47(59)	19(84)	8(106)	02MAY-23AUG
Acronicta morula G. & R.	0(.)	1(224)	0(.)	02MAY-02MAY
Acronicta interrupta Gn.	1(242)	1(224)	0(.)	16MAY-23AUG
Acronicta lobeliae Gn.	2(215)	4(163)	0(.)	10MAY-23AUG
Acronicta fragilis Gn.	85(31)	23(78)	52(41)	10MAY-06AUG
Acronicta ovata Grt.	177(13)	57(45)	177(12)	16MAY-05SEP
Acronicta modica Wlk.	49(56)	10(120)	4(147)	05JUN-15AUG

continuea

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Acronicta haesitata</i> (Grt.)	3(191)	9(125)	35(49)	29MAY-13AUG
<i>Acronicta tristis</i> Sm.	27(78)	0(.)	0(.)	22JUN-03AUG
<i>Acronicta increta</i> Morr.	60(46)	10(120)	12(91)	29MAY-23AUG
<i>Acronicta inclara</i> Sm.	18(95)	12(111)	34(51)	05JUN-03AUG
<i>Acronicta retardata</i> (Wlk.)	33(71)	10(120)	13(83)	05JUN-14AUG
<i>Acronicta caesarea</i> Sm.	0(.)	0(.)	22(64)	12JUN-30JUL
<i>Acronicta impleta</i> Wlk.	1(242)	5(157)	1(212)	02MAY-15AUG
<i>Acronicta lithospila</i> Grt.	1(242)	0(.)	0(.)	13JUL-13JUL
<i>Polygrammate hebraeicum</i> Hbn.	50(54)	19(84)	27(60)	05JUN-29AUG
<i>Eudryas grata</i> (F.)	6(160)	7(141)	0(.)	28JUN-15AUG
<i>Apamea amputatrix</i> (Fitch)	1(242)	0(.)	0(.)	28JUN-28JUN
<i>Agroperina dubitans</i> (Wlk.)	1(242)	0(.)	0(.)	23AUG-23AUG
<i>Agroperina helva</i> (Grt.)	0(.)	2(199)	0(.)	28AUG-28AUG
<i>Oligia modica</i> (Gn.)	0(.)	0(.)	1(212)	21AUG-21AUG
<i>Oligia crytora</i> (Franc.)	2(215)	0(.)	0(.)	13JUL-13JUL
<i>Oligia mactata</i> (Gn.)	1(242)	0(.)	1(212)	26SEP-02OCT
<i>Oligia illocata</i> (Wlk.)	17(100)	19(84)	7(116)	05SEP-17OCT
<i>Amphipoea americana</i> (Speyer)	1(242)	0(.)	0(.)	03AUG-03AUG
<i>Papaipema lysimachiae</i> Bird	0(.)	1(224)	0(.)	24SEP-24SEP
<i>Papaipema inquaesita</i> (G.& R.)	1(242)	0(.)	0(.)	26SEP-26SEP
<i>Papaipema baptisiae</i> (Bird)	0(.)	2(199)	0(.)	10SEP-24SEP
<i>Papaipema marginidens</i> (Gn.)	0(.)	1(224)	0(.)	11OCT-11OCT
<i>Papaipema cerussata</i> (Grt.)	1(242)	0(.)	0(.)	17OCT-17OCT
<i>Papaipema unimoda</i> (Sm.)	1(242)	0(.)	0(.)	26SEP-26SEP
<i>Achatodes zaeae</i> (Harr.)	0(.)	0(.)	1(212)	21AUG-21AUG
<i>Iodopepla u-album</i> (Gn.)	1(242)	0(.)	0(.)	03AUG-03AUG
<i>Euplexia benesimilis</i> McD.	28(76)	36(60)	15(79)	10MAY-26SEP
<i>Phlogophora iris</i> Gn.	9(142)	32(65)	8(106)	16MAY-28AUG
<i>Phlogophora periculosa</i> Gn.	39(65)	23(78)	7(116)	15AUG-21SEP
<i>Chytonix palliatricula</i> (Gn.)	34(69)	29(67)	14(82)	16MAY-15AUG
<i>Hyppa xylinoides</i> (Gn.)	1(242)	1(224)	0(.)	03AUG-06AUG
<i>Nedra ramosula</i> (Gn.)	4(179)	0(.)	0(.)	27APR-26SEP
<i>Phosphila turbulenta</i> Hbn.	34(69)	52(47)	83(30)	12JUN-03AUG
<i>Phosphila miselioides</i> (Gn.)	14(109)	11(118)	17(74)	05JUN-19SEP
<i>Callopietria mollissima</i> (Gn.)	52(52)	36(60)	33(53)	10MAY-09SEP
<i>Amphipyra pyramidoides</i> Gn.	14(109)	26(74)	16(76)	23JUL-30OCT
<i>Amphipyra tragopoginis</i> (Cl.)	0(.)	1(224)	0(.)	10JUL-10JUL
<i>Anorthodes tarda</i> (Gn.)	19(94)	67(40)	8(106)	10MAY-26SEP
<i>Balsa labecula</i> (Grt.)	1(242)	3(179)	0(.)	30MAY-13JUL
<i>Spodoptera frugiperda</i> (J.E. Smith)	3(191)	0(.)	3(161)	28AUG-17OCT
<i>Spodoptera ornithogalli</i> (Gn.)	1(242)	1(224)	0(.)	26SEP-11OCT
<i>Elaphria versicolor</i> (Grt.)	0(.)	2(199)	0(.)	12JUN-12JUN
<i>Elaphria festivoides</i> (Gn.)	11(130)	102(26)	8(106)	10MAY-06JUL
<i>Elaphria grata</i> Hbn.	1(242)	2(199)	6(124)	20JUN-09SEP
<i>Galgula partita</i> Gn.	1(242)	1(224)	2(184)	12JUN-22OCT

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Perigea xanthioides</i> Gn.	2(215)	0(.)	0(.)	28JUN-03AUG
<i>Platysenta vecors</i> (Gn.)	60(46)	4(163)	4(147)	02MAY-26SEP
<i>Platysenta sutor</i> (Gn.)	1(242)	0(.)	1(212)	22JUN-05SEP
<i>Lithophane semiusta</i> Grt.	1(242)	0(.)	1(212)	08APR-21SEP
<i>Lithophane innominata</i> (Sm.)	0(.)	1(224)	1(212)	28MAR-02APR
<i>Lithophane hemina</i> Grt.	4(179)	9(125)	3(161)	19MAR-17SEP
<i>Lithophane signosa</i> (Wlk.)	1(242)	0(.)	0(.)	26SEP-26SEP
<i>Lithophane antennata</i> (Wlk.)	1(242)	1(224)	0(.)	28MAR-04MAY
<i>Lithophane grotei</i> Riley	0(.)	2(199)	1(212)	28MAR-12APR
<i>Lithophane unimoda</i> (Lint.)	0(.)	0(.)	2(184)	02APR-08APR
<i>Pyreferra hesperidago</i> (Gn.)	0(.)	4(163)	8(106)	19MAR-29APR
<i>Pyreferra citrombra</i> Franc.	1(242)	9(125)	6(124)	19MAR-12MAY
<i>Eupsilia n. sp.</i>	1(242)	0(.)	0(.)	27APR-27APR
<i>Eupsilia morrisoni</i> (Grt.)	1(242)	4(163)	1(212)	19MAR-11OCT
<i>Epiglaea decliva</i> (Grt.)	10(136)	1(224)	11(95)	12OCT-30OCT
<i>Chaetaglaea sericea</i> (Morr.)	1(242)	0(.)	0(.)	17OCT-17OCT
<i>Eucirroedia pampina</i> (Gn.)	0(.)	0(.)	1(212)	10OCT-10OCT
<i>Sunira bicolorago</i> (Gn.)	32(73)	65(41)	21(68)	28AUG-31OCT
<i>Anathix ralla</i> (G. & R.)	38(66)	0(.)	0(.)	23AUG-17OCT
<i>Eutolype rolandi</i> Grt.	0(.)	2(199)	1(212)	08APR-19APR
<i>Copipanolis styracis</i> (Gn.)	8(149)	8(131)	11(95)	27MAR-27APR
<i>Psaphida resumens</i> Wlk.	10(136)	33(64)	1(212)	15APR-27APR
<i>Psaphida thaxteriana</i> (Grt.)	0(.)	8(131)	0(.)	28MAR-28MAR
<i>Polia nimbosea</i> (Gn.)	55(49)	35(63)	126(19)	12JUN-03AUG
<i>Polia imbrifera</i> (Gn.)	1(242)	3(179)	3(161)	05JUN-02JUL
<i>Polia detracta</i> (Wlk.)	0(.)	106(24)	25(62)	05JUN-17JUL
<i>Polia goodelli</i> (Grt.)	0(.)	1(224)	0(.)	12JUN-12JUN
<i>Polia latex</i> (Gn.)	131(20)	79(35)	94(26)	24MAY-29AUG
<i>Lacanobia legitima</i> (Grt.)	1(242)	0(.)	0(.)	13JUL-13JUL
<i>Hadena ectypa</i> (Morr.)	0(.)	0(.)	1(212)	10JUL-10JUL
<i>Lacinipolia renigera</i> (Steph.)	1(242)	1(224)	0(.)	28JUN-01OCT
<i>Lacinipolia lorea</i> (Gn.)	1(242)	0(.)	0(.)	08JUN-08JUN
<i>Pseudaletia unipuncta</i> (Haw.)	9(142)	15(96)	5(134)	25APR-26SEP
<i>Leucania phragmatidicola</i> Gn.	12(122)	0(.)	1(212)	06JUN-28JUN
<i>Leucania multilinea</i> Wlk.	1(242)	0(.)	0(.)	22JUN-22JUN
<i>Leucania lapidaria</i> (Grt.)	5(166)	0(.)	0(.)	22JUN-28JUN
<i>Leucania ursula</i> (Fbs.)	1(242)	1(224)	1(212)	28JUN-10JUL
<i>Orthosia rubescens</i> (Wlk.)	11(130)	93(29)	13(83)	08APR-25MAY
<i>Orthosia hibisci</i> (Gn.)	3(191)	15(96)	15(79)	02APR-27APR
<i>Crocigrapha normani</i> (Grt.)	16(101)	23(78)	0(.)	19APR-25MAY
<i>Himella intractata</i> (Morr.)	1(242)	0(.)	0(.)	08MAY-08MAY
<i>Egira alternans</i> (Wlk.)	2(215)	4(163)	3(161)	08APR-16MAY
<i>Achatia distincta</i> Hbn.	3(191)	12(111)	0(.)	19APR-18MAY
<i>Morrisonia evicta</i> (Grt.)	1(242)	0(.)	0(.)	18MAY-18MAY
<i>Morrisonia confusa</i> (Hbn.)	79(34)	119(22)	0(.)	02MAY-12JUN

continued

SPECIES	TOTAL CATCH (RANK)			INCLUSIVE FLIGHT RANGE
	1984	1985	1986	
<i>Nephelodes minians</i> Gn.	28(76)	28(69)	7(116)	06SEP-26SEP
<i>Homorthodes furfurata</i> (Grt.)	6(160)	3(179)	1(212)	06JUL-27JUL
<i>Protorthodes oviduca</i> (Gn.)	1(242)	0(.)	0(.)	25MAY-25MAY
<i>Pseudorthodes vecors</i> (Gn.)	11(130)	3(179)	11(95)	16MAY-13AUG
<i>Orthodes crenulata</i> (Butler)	2(215)	0(.)	0(.)	18JUL-23AUG
<i>Orthodes cynica</i> (Gn.)	252(8)	188(10)	137(17)	16MAY-18JUL
<i>Agrotis venerabilis</i> Wlk.	2(215)	1(224)	0(.)	10SEP-26SEP
<i>Agrotis ipsilon</i> (Hufn.)	4(179)	3(179)	3(161)	23JUL-28OCT
<i>Ochropleura plecta</i> (L.)	0(.)	1(224)	1(212)	29MAY-06SEP
<i>Peridroma saucia</i> (Hbn.)	2(215)	1(224)	4(147)	12MAY-17SEP
<i>Diarsia jucunda</i> (Wlk.)	6(160)	0(.)	1(212)	10JUL-03AUG
<i>Spaelotis clandestina</i> (Harr.)	0(.)	2(199)	1(212)	12JUN-10JUL
<i>Graphiphora haruspica</i> (Grt.)	1(242)	0(.)	0(.)	03AUG-03AUG
<i>Xestia dolosa</i> Franc.	14(109)	18(90)	8(106)	16MAY-26SEP
<i>Xestia normaniana</i> (Grt.)	13(115)	0(.)	1(212)	23AUG-09SEP
<i>Xestia smithii</i> (Snell.)	2(215)	0(.)	1(212)	05SEP-09SEP
<i>Xestia bicarnea</i> (Gn.)	16(101)	9(125)	2(184)	06AUG-26SEP
<i>Xestia badinodis</i> (Grt.)	0(.)	3(179)	0(.)	24SEP-01OCT
<i>Cerastis tenebrifera</i> (Wlk.)	1(242)	1(224)	0(.)	14APR-19APR
<i>Anaplectoides prasina</i> (D.& S.)	0(.)	0(.)	1(212)	17JUL-17JUL
<i>Eueretagrotis perattenta</i> (Grt.)	1(242)	8(131)	2(184)	03JUL-24SEP
<i>Eueretagrotis attenta</i> (Grt.)	8(149)	0(.)	1(212)	10JUL-08AUG
<i>Heptagrotis phyllophora</i> (Grt.)	5(166)	2(199)	2(184)	06JUL-03AUG
<i>Abagrotis alternata</i> (Grt.)	66(43)	16(94)	0(.)	18JUL-26SEP
<i>Heliothis zea</i> (Boddie)	3(191)	0(.)	2(184)	21SEP-28OCT