1-1-1995

Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain section.

Linda Butler

Follow this and additional works at: [https://researchrepository.wvu.edu/
wv_agricultural_and_forestry_experiment_station_bulletins](https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins)

Digital Commons Citation

Butler, Linda, "Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain section." (1995). *West Virginia Agricultural and Forestry Experiment Station Bulletins*. 712.
https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins/603

This Bulletin is brought to you for free and open access by the Davis College of Agriculture, Natural Resources And Design at The Research Repository @ WVU. It has been accepted for inclusion in West Virginia Agricultural and Forestry Experiment Station Bulletins by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

Canopy Arthropods at Fernow Experimental Forest in
West Virginia's Allegheny Mountain Section: I.
Macrolepidopterous Moths Collected
by Blacklight Trap During a Dimilin Impact Study

RECEIVED
JAN 29 1996

EVANSDALE LIBRARY
WEST VIRGINIA UNIVERSITY

S127
E1
712
Nov. 1995

Bulletin 712 • November 1995
Agricultural and Forestry Experiment Station
West Virginia University

Authors

Linda Butler is Professor of Entomology and Vicki Kondo and Gregory Chrislip are Entomology Research Assistants, Division of Plant and Soil Sciences, College of Agriculture and Forestry, West Virginia University.

This research was supported by a grant from the USDA Forest Service. We would like to acknowledge the assistance and cooperation of the USDA Forest Service Appalachian Integrated Pest Management Program personnel, our undergraduate field and laboratory assistants and the data management assistance of Dr. Edwin Townsend and David Partsch.

West Virginia University
Agricultural and Forestry Experiment Station
College of Agriculture and Forestry
Rosemary Haggett, Director
Morgantown, WV

Canopy Arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain Section: I. Macrolepidopterous Moths Collected by Blacklight Trap During a Dimilin Impact Study

Linda Butler, Vicki Kondo and Gregory Chrislip

INTRODUCTION

Gypsy moth, *Lymantria dispar* (L.) (Lepidoptera: Lymantriidae) was introduced into North America in 1869 and has now spread over much of the eastern half of the United States. Larval infestations were first found in West Virginia in 1975. Gypsy moth larvae prefer oak foliage. Their capacity to cause defoliation, which may result in tree mortality, has caused gypsy moth to be considered the most serious hardwood defoliator in the eastern United States (Doane and McManus, 1981).

Among the management methods employed against gypsy moth have been aerial applications of insecticides. In recent years the insect growth regulator Dimilin (1-(4 chlorophenyl)-3-(2, b-diflubenzoyl) urea) or diflubenzuron has been applied to thousands of acres of forest land for gypsy moth suppression. It was first used in West Virginia in 1979. While Dimilin is considered to have minimal direct effects on vertebrates, a number of studies have shown impact of Dimilin on non-target terrestrial forest arthropods (Martinat et al., 1988, 1993; Sample et al., 1993; Butler and Kondo, 1993).

In 1989, we began a study to evaluate Dimilin impact on canopy arthropods. The study was conducted at the Fernow Experimental Forest near Parsons, West Virginia, in the Allegheny Mountain section of the state. During the next six years we sampled by means of blacklight traps, pole pruning of foliage and bur-lap-banded trees. Results of the Dimilin impact study and arthropod diversity in foliage will be presented elsewhere. We present here an evaluation of richness and abundance of macrolepidopterous moths as collected by blacklight trap on four closed watersheds from 1990–1994.

MATERIALS AND METHODS

Study Sites

The Fernow Experimental Forest is located on the unglaciated Allegheny Plateau (latitude 39° 05'N, longitude 79° 41'W). Elevations range from 533 to 1,112 m with rugged topography and slopes of 10 to 60 percent. The Fernow has a mean of 145 frost-free days, a mean annual precipitation of 147 cm and mean temperature of 9° C (Northeastern Forest Experimental Station, 1987). The experimental forest encompasses 1,902 ha and has been divided into a series of watersheds for research purposes (Adams et al., 1993). This study was conducted on watersheds 1, 4, 7 and 13. Data on predominant tree species on watersheds are from Stephenson 1992 (unpublished).

Watershed 1 is 30.11 ha with a NE aspect. It was clearcut to 6 inches DBH (diameter at breast height) in 1957 and 1958. Primary tree species with regard to basal area include sugar maple (*Acer saccharum* Marsh.), white basswood (*Tilia heterophylla* Vent.), beech (*Fagus grandifolia* Ehrh.), black locust (*Robinia pseudoacacia* L.), cucumber magnolia (*Magnolia acuminata* L.) and red oak (*Quercus rubra* L.). The forest floor is covered by herbaceous growth, primarily nettles (*Urtica* spp.).

Watershed 4 is 38.73 ha with an ESE aspect. It was cut heavily from 1905 to 1910 but has not been timbered since that time. Woody shrubs and vines are common, but little herbaceous ground cover is present. Primary tree species include beech, sugar maple, moosewood (*Acer pensylvanicum* L.), tulip-poplar (*Liriodendron tulipifera* L.), red oak, white basswood and cucumber magnolia. In parts of the watershed are dense understories of young black cherry (*Prunus serotina* Ehrh.).

Watershed 7 is 24.23 ha with an E aspect. It is a new growth forest; in places the canopy is dense with little understory vegetation except hayscented fern (*Dennstaedtia punctinlobula* (Michx.) Moore). The watershed was clearcut and maintained barren with herbicides between 1964 and 1969. Tree species include black birch (*Betula lenta* L.), sugar maple, red oak, black cherry, tulip-poplar and blackgum (*Nyssa sylvatica* Marsh.).

Watershed 13 is 14.23 ha with a NNE aspect. It is a relatively mature forest with little understory vegetation. The watershed was cut early in the century and had light selective cutting up to 1960, but no cutting subsequently. Predominant tree species include sugar maple, red oak, tulip-poplar, black cherry, black birch and white basswood.

Arthropod sampling on the Fernow began in 1989 with pole pruning of foliage and banding of trees. Light traps were employed beginning in 1990. On May 16, 1992, Dimilin 4L was applied to watersheds 1 and 13 at a rate of 0.5 oz AI/acre. Light trap sampling continued through mid-August 1994.

One night each week from early or mid-May to early or mid-August (May 7–August 13, 1990; May 1–August 12, 1991; May 11–August 10, 1992; May 10–August 9, 1993; and May 9–August 8, 1994) four blacklight traps were operated on the Fernow, one on each watershed. Trapping occurred for all watersheds on the same nights. Traps were the 8-watt Universal Light Trap (Model 2851 K) from BioQuip Products (Gardena, CA) and were battery-operated and equipped with photoelectric eyes. Traps were operated the same night each week with no attempt being made to avoid nights that were suboptimal for light trapping (cool, full moon, etc.).

Macrolepidopterous moths were identified by comparison with the West Virginia University Arthropod Collection; counts were made for each species. Voucher specimens from the study are deposited in the WVU Arthropod Collection. Species are listed after the checklist given by Hodges et al. (1983).

RESULTS

Over the five years of blacklight trapping 376 species of macrolepidopterous moths representing 13 families were taken from the four Fernow watersheds (Table 1). Families with the most species were the Noctuidae with 196 and Geometridae with 103. Considerable variation occurred in species captured from year to year (Table 1). For example, of the 196 species of Noctuidae taken over five years, only 42 of the species appeared in the traps all five years. Fifty-seven of the 196 noctuids were represented in only one of the five years. Somewhat less variability was seen for the Geometridae where 42 of 103 species were collected every year and only 17 species were represented in a single year (Table 1).

Table 2 lists the 376 species taken during the study. Species are arranged from most to least abundant over the total five-year period; inclusive flight range is given from earliest date of capture to latest date of capture over the five years. The most abundant species were the geometrids *Itame pustularia* (Gn.), *Hydrelia inornata* (Hulst), *Lomographa glomeraria* (Grt.), *Probole amicaria* (H.-S.), *Melanolophia canadaria* (Gn.), *Iridopsis larvaria* (Gn.) and *Euphyia unangulata* (Haw.), the arctiids *Halysidota tessellaris* (J.E. Smith) and *Hypoprepia fucosa* Hbn. and the notodontid *Peridea basitriens* (Wlk.) (Table 2). Despite accounting for 52 percent of all species taken during the study, the noctuids did not make up the most abundant species. *Euplexia benesimilis* McD. was the most abundant noctuid, but its numbers over five years were 10 percent of those of the most abundant geometrid. Forty of the 196 noctuid species collected in the study were represented by a single individual (Table 2).

Other relatively abundant species were the lasiocampid *Malacosoma americanum* (F.), the arctiid *Clemensia albata* Pack., the geometrids *Heterophleps refusaria* (Wlk.), *Campaea perlata* (Gn.), *Hydria prunivorata* (Fgn.) and *Eupithecia herefordaria* C. & S., the drepanid *Drepana arcuata* Wlk., noc-

tuids *Idia rotundalis* (Wlk.), *Polia detracta* (Wlk.), *Bomolocha edictalis* (Wlk.), and *Acronicta ovata* Grt. and the notodontid *Peridea angulosa* (J.E. Smith).

Abundance of many species fluctuated strongly over the years of the study. The most abundant species, *Itame pustularia*, ranged from 452 to 504 in 1991–1992, increased to 1,709 in 1993 and to 4,223 in 1994. The peak catch of *Hydrelia inornata* was in 1991 with 3,532 individuals, with 1994 being the second highest year with 1,354. While two of the *Lomographa* were almost equally abundant during the study (2,876 *L. glomeraria* and 2,301 *L. vestaliata*), there was no clear pattern each year to indicate that they were cycling together or were interfering with each other. *L. glomeraria* was most abundant in 1993 (1,634), while *L. vestaliata* was most abundant in 1994 (841).

No pattern seemed evident regarding the year of greatest abundance for the most abundant species. More *Idia rotundalis* were trapped in 1990 (222) than in any other study year. *Hydrelia inornata*, *Halysidota tessellaris*, *Melanolophia canadaria*, *Iridopsis larvaria*, *Clemensia albata*, *Heterophleps refusaria*, *Drepana arcuata*, *Euplexia benesimilis*, and *Bomolocha edictalis* were captured in greatest numbers in 1991. While 1992 saw the lowest abundance of trapped moths for many of the most abundant species, *Hydria prunivorata* was most abundant in traps that year (291). The year 1993 was the peak catch year for *Lomographa glomeraria*, *Peridea basitriens*, *Malacosoma americanum*, *Campaea perlata*, *Polia detracta* and *Eupethecia herefordaria*. For 1994, *Itame pustularia*, *Lomographa vestaliata*, *Probole amicaria*, *Euphyia unangulata*, *Hypoprepia fucosa*, *Acronicta ovata* and *Peridea angulosa* were trapped in greatest abundance.

The 21 most abundant species were trapped each year of the study. The geometrid *Eupithecia herefordaria*, the 22nd most abundant species, was represented by 590 individuals in traps during the study, but was not captured at all in 1990.

Relatively little year-to-year fluctuation in light-trap capture was seen in abundance of some moth species, for example, *Peridea basitriens*, *Idia rotundalis* and the notodontid *Nadata gibbosa* (J.E. Smith).

The inclusive flight range given for each species reflects the limitation of the current study. Traps were operated only from early or mid-May to early or mid-August each year. Many species began flying before and/or continued flying after those dates.

Variation in length of flight season among species reflects varying degrees of synchrony of adult emergence, differences in adult longevity, effect of multivoltine versus univoltine species and population levels of the different species. Bi- or multivoltine species include *Melanolophia canadaria*, *Lomographa vestaliata*, *L. glomeraria*, *Campaea perlata*, *Probole amicaria*, *Iridopsis larvaria*, *Euphyia unangulata*, *Peridea basitriens* and *Drepana arcuata*. Flight ranges for *Halysidota tessellaris* and *Malacosoma americanum* are examples of univoltine

species. *Halysidota tessellaris* shows a relative lack of synchrony in adult emergence as reflected in the broad flight ranges, while adult emergence in *M. americanum* is relatively synchronized.

DISCUSSION

Trapping of 376 species of macrolepidopterous moths from early or mid-May through early or mid-August over a period of five years indicates a high level of diversity in the Fernow Experimental Forest. While four traps were operated simultaneously on four different watersheds, all watersheds were relatively small, ranging in size from 14 to 30 ha, and they were relatively similar in vegetation composition. Had traps been operated over a long portion of the season, for example, from March to November, additional moth species may have been collected.

During a simultaneous study of macrolepidopterous larvae on foliage of four tree species groups, commonly collected larvae whose adults fly at times not sampled by the light traps included *Erannis tiliaria* Harr. and *Alsophila pometaria* Harr., whose females are wingless and whose males fly in the fall, *Epiglaea decliva* Grt., a fall flying moth, and *Phigalia titea*, whose females are flightless and whose males fly in early spring (Butler et al. 1995).

Species of moths that develop as larvae on host plants unrepresented or poorly represented in the study watersheds were not collected or were in very low abundance in this study. For example, *Catocala palaeogama* Gn. and *C. residua* Grt. feed exclusively on hickory as larvae. Virtually no hickory was located in the study sites; only one individual of each species was taken during the five-year study.

In an earlier baseline study of macrolepidopterous moths blacklight-trapped at Coopers Rock State Forest, West Virginia (Butler and Kondo 1991), 400 species were collected in a single trap at one location from 1984 to 1986. There were several differences between the two studies. At Coopers Rock, a single 15-watt trap was used at a single site rather than 8-watt traps at four adjacent watersheds. The Coopers Rock trap was operated for approximately 30 weeks each year for three years, whereas the Fernow traps were in operation for five years, but only for 15 weeks each year. In addition, the Fernow traps were operated the same night each week (Monday) regardless of expected weather conditions, whereas the Coopers Rock trap nights, although about at 7-day intervals, were chosen to avoid unseasonably low temperatures when possible, thus likely increasing catch abundance and diversity.

Many similarities were seen between the results of the Coopers Rock and current studies. In the earlier study the most abundant species were also the geometrids *Itame pustularia*, *Probole amicaria*, *Lomographa glomeraria* and *Melanolophia canadaria*; *Campaea perlata* was also relatively abundant but

Hydrelia inornata (second in abundance at Fernow) was relatively less abundant. Other abundant species at Coopers Rock included *Halysidota tessellaris*, *Clemensia albata* and *Malacosoma americanum*, again, similar to current results. However, some noctuids and notodontids were relatively more abundant at Coopers Rock than at Fernow; these include *Orthodes cynica* (Gn.), *Idia aemula* Hbn., *Zanclognatha ochreipennis* (Grt.), *Nadata gibbosa* (J.E. Smith), *Heterocampa guttivitta* (Wlk.) and *Peridea angulosa* (J.E. Smith).

The degree of year-to-year fluctuation in moth species richness was similar between the two studies; again the greatest fluctuation occurred for the Noctuidae.

Most of the differences in species richness between the two studies is attributable to the longer trapping season at Coopers Rock providing species not captured at Fernow; for example, *Paleacrita vernata* (Peck), *P. merriccata* Dyar, *Phigalia titea* (Cram.), *Pyreferra citrombra* Franc., *Eutolyte rolandi* Grt., *Copipanolis styrcis* (Gn.), and *Psaphida thaxteriana* (Grt.), which fly in early spring (March, April). Species such as *Erannis tiliaria*, *Ennomos magnaria* Gn., *Tolyte vellea* (Stoll), *Anticarsia gemmatalis* Hbn., *Papaipema* spp., *Epiglaea decliva*, *Chaetagnalea sericea* (Morr.) and *Eucirroedia pampina* (Gn.) fly in October and November and thus were not taken at Fernow. Some species, particularly *Lithophane* spp., fly from the fall and winter into early spring and were poorly represented in the Fernow traps.

Several species were taken at Fernow but not at Coopers Rock. *Meganola spodia* Franc. is common in more eastern areas of West Virginia (Butler 1989). Only seven *M. spodia* were taken at Fernow, all in 1991; the species has not been taken at Coopers Rock. *Eupithecia herefordaria* was not recorded at Coopers Rock, possibly because of misidentification with *E. miserulata* Grt. *Sicya macularia* (Harr.), *Itame subcessaria* (Wlk.), *Cosmia calami* (Harr.), *Pyrrhia umbra* (Hufn.), *Pyrrhia exprimens* (Wlk.), *Oruza albocostaliata* (Pack.), *Lithophane querquera* Grt., *Eupsilia vinulenta* (Grt.) and *Orthosia revicta* (Morr.) are examples of Fernow species not taken at Coopers Rock.

Light traps have once again shown their value in facilitating a study of richness and abundance of forest macrolepidoptera.

SUMMARY

Macrolepidopterous moths were sampled from four adjacent watersheds at Fernow Experimental Forest from 1990 to 1994. Each year, four 8-watt black-light traps were operated from early or mid-May to early or mid-August and resulted in the catch of 376 species of macrolepidopterous moths from 13 families over the five-year study. Yearly fluctuation was considerable for many species.

LITERATURE CITED

- Adams, M.B., J.N. Kochenderfer, F. Wood, T.R. Angradi, and P. Edwards. 1993. Forty years of hydrometeorological data from the Fernow Experimental Forest, West Virginia. USDA Forest Service. Gen. Tech. Rpt. NE-184. 24 pp.
- Butler, L. 1989. Observations on *Meganola spodia* Franclemont (Lepidoptera: Noctuidae) with a description of the mature larva. Proc. Entomol. Soc. Wash. 91:615-619.
- Butler, L. and V. Kondo. 1991. Macrolepidopterous moths collected by black-light trap at Coopers Rock State Forest, West Virginia: a baseline study. WV Agric. and For. Exp. Sta. Bull. 705, 25 pp.
- Butler, L. and V. Kondo. 1993. Impact of Dimilin on non-target Lepidoptera: Results of an operational gypsy moth suppression program at Coopers Rock State Forest, West Virginia. WV Agric. and For. Exp. Sta. Bull. 710. 21 pp.
- Butler, L., G. Chrislip, and V. Kondo. 1995. Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain Section: II. Macrolepidopterous larvae on foliage and under burlap bands. WV Agric. and For. Exp. Sta. Bull. 713. 16 pp.
- Doane, C.C. and M.L. McManus. 1981. The gypsy moth: research toward integrated pest management. USDA Forest Service Technical Bull. 1584. 757 pp.
- Hodges, R.W., T. Dominic, D.R. Davis, D.C. Ferguson, J.G. Franclemont, E.G. Munroe, and J.A. Powell. 1983. Checklist of the Lepidoptera of America North of Mexico. London. E.W. Classey Ltd. and Washington, D.C. Wedge Entomol. Res. Found. 284 pp.
- Martinat, P.J., D.T. Jennings, and R.C. Whitmore. 1993. Effects of diflubenzuron on the litter spider and orthopteroid community in a central appalachian forest infested with gypsy moth (Lepidoptera: Lymantriidae). Environ. Entomol. 22:1003-1008.
- Martinat, P.J., C.C. Coffman, K. Dodge, R.J. Cooper, and R.C. Whitmore. 1988. Effect of diflubenzuron on the canopy arthropod community in a central Appalachian forest. J. Econ. Entomol. 81:261-267.
- Northeastern Forest Experiment Station. 1987. Forest Research: Fernow Experimental Forest. USDA. 12 pp.
- Sample, B.E., L. Butler, and R.C. Whitmore. 1993. Effects of an operational application of Dimilin on non-target insects. Can. Entomol. 125:173-179.
- Stephenson, S.L. 1992. Department of Biology, Fairmont State College, Fairmont, West Virginia. Personal Communication.

TABLE 1.

Number of species of Macrolepidoptera in all families taken by blacklight traps at the Fernow Experimental Forest, West Virginia during any one, two, three, four and five years of the study (1990–1994).

FAMILY	Number of Species Trapped During a Given Number of Years					TOTAL SPECIES
	1	2	3	4	5	
THYATIRIDAE	0	0	0	1	2	3
DREPANIDAE	0	0	1	0	1	2
GEOMETRIDAE	17	8	14	22	42	103
EPIPLEMIDAE	0	1	0	0	0	1
MIMALLONIDAE	1	0	0	1	0	2
APATELODIDAE	0	0	1	0	1	2
LASIOCAMPIDAE	0	1	0	1	2	4
SATURNIIDAE	0	1	0	1	3	5
SPHINGIDAE	2	0	0	1	5	8
NOTODONTIDAE	4	2	4	5	13	28
ARCTIIDAE	3	3	2	3	6	17
LYMANTRIIDAE	0	1	0	1	3	5
NOCTUIDAE	57	38	30	29	42	196
TOTAL SPECIES	84	55	52	65	120	376

TABLE 2.

Macrolepidoptera species collected by blacklight trap at the Fernow Experimental Forest, West Virginia during 1990–1994. Species are listed from most to least abundant; inclusive flight range during the five-year period is given.

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Itame pustularia</i> (Gn.)	GEO	452	467	504	1709	4223	7355	17JUN–13AUG
<i>Hydrelia inornata</i> (Hulst)	GEO	421	3532	106	994	1354	6407	01MAY–12AUG
<i>Lomographa glomeraria</i> (Grt.)	GEO	596	437	207	1634	2	2876	01MAY–13JUL
<i>Lomographa vestaliata</i> (Gn.)	GEO	139	564	67	690	841	2301	10MAY–16JUL
<i>Halysidota tessellaris</i> (J.E. Smith)	ARC	275	1172	232	163	404	2246	28MAY–10AUG
<i>Probole amicarica</i> (H.-S.)	GEO	85	421	99	592	836	2033	01MAY–13AUG
<i>Melanophia canadaria</i> (Gn.)	GEO	206	1085	147	188	336	1962	01MAY–13AUG
<i>Iridopsis larvaria</i> (Gn.)	GEO	303	939	9	62	263	1576	01MAY–13AUG
<i>Euphyia unangulata</i> (Haw.)	GEO	263	302	74	275	323	1237	01MAY–13AUG
<i>Hypoprepia fucosa</i> Hbn.	ARC	318	211	76	136	350	1091	01JUL–13AUG

continued

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Peridea basitriens</i> (Wlk.)	NOT	233	134	182	315	158	1022	13MAY–13AUG
<i>Malacosoma americanum</i> (F.)	LAS	73	61	32	212	495	873	06JUN–20JUL
<i>Clemensia albata</i> Pack.	ARC	51	552	62	73	35	773	11MAY–13AUG
<i>Heterophleps refusaria</i> (Wlk.)	GEO	107	445	74	65	76	767	06MAY–12AUG
<i>Campaea perlata</i> (Gn.)	GEO	103	158	42	248	201	752	20MAY–13AUG
<i>Drepana arcuata</i> Wlk.	DRE	115	368	5	84	148	720	01MAY–13AUG
<i>Euplexia benesimilis</i> McD.	NOC	100	498	22	39	41	700	10MAY–13AUG
<i>Idia rotundalis</i> (Wlk.)	NOC	222	73	92	118	150	655	10JUN–13AUG
<i>Polia detracta</i> (Wlk.)	NOC	80	171	88	199	94	632	28MAY–13JUL
<i>Hydria prunivora</i> (Fgn.)	GEO	5	229	291	59	16	600	11MAY–13AUG
<i>Bomolocha edictalis</i> (Wlk.)	NOC	46	217	42	183	107	595	17JUN–13AUG
<i>Eupithecia herefordaria</i> C. & S.	GEO	0	117	14	380	79	590	10MAY–12AUG
<i>Acronicta ovata</i> Grt.	NOC	30	90	8	123	290	541	28MAY–13AUG
<i>Peridea angulosa</i> (J.E. Smith)	NOT	148	35	103	50	199	535	11JUN–13AUG
<i>Chytolita morbidalis</i> (Gn.)	NOC	52	96	79	151	102	480	20MAY–13JUL
<i>Hyphantria cunea</i> (Drury)	ARC	4	60	39	81	284	468	10MAY–06JUL
<i>Heterocampa guttivitta</i> (Wlk.)	NOT	102	178	0	78	82	440	10MAY–09AUG
<i>Semiothisa aemulataria</i> (Wlk.)	GEO	22	370	8	25	8	433	01MAY–12AUG
<i>Orthodes cynica</i> (Gn.)	NOC	0	137	45	80	153	415	20MAY–09AUG
<i>Dryocampa rubicunda</i> (F.)	SAT	15	129	20	134	104	402	10MAY–10AUG
<i>Zanclognatha laevigata</i> (Grt.)	NOC	161	65	26	84	53	389	07JUN–13AUG
<i>Semiothisa ocellinata</i> (Gn.)	GEO	115	130	15	63	64	387	10MAY–13AUG
<i>Nadata gibbosa</i> (J.E. Smith)	NOT	97	65	40	95	85	382	13MAY–13AUG
<i>Idia aemula</i> Hbn.	NOC	38	66	17	100	158	379	24MAY–13AUG
<i>Paonias myops</i> (J.E. Smith)	SPH	36	105	34	126	73	374	10MAY–13AUG
<i>Zale unilineata</i> (Grt.)	NOC	15	30	17	93	183	338	01MAY–15JUN
<i>Zanclognatha ochreipennis</i> (Grt.)	NOC	81	23	11	139	67	321	14JUN–13AUG
<i>Besma endropiaria</i> (G. & R.)	GEO	69	97	0	79	75	320	20MAY–12AUG
<i>Paonias excaecatus</i> (J.E. Smith)	SPH	24	54	37	133	62	310	20MAY–10AUG
<i>Malacosoma distria</i> Hbn.	LAS	84	58	25	51	90	308	17JUN–20JUL
<i>Morrisonia confusa</i> (Hbn.)	NOC	15	32	13	151	85	296	01MAY–14JUN
<i>Oligocentria semirufescens</i> (Wlk.)	NOT	33	67	18	124	51	293	03JUN–13AUG
<i>Scopula limboundata</i> (Haw.)	GEO	53	73	50	60	53	289	03JUN–08AUG
<i>Semiothisa pinistrobata</i> Fgn.	GEO	107	162	0	6	0	275	13MAY–13AUG
<i>Zale minerea</i> (Gn.)	NOC	17	23	34	129	72	275	01MAY–06AUG
<i>Hypagyrtis unipunctata</i> (Haw.)	GEO	10	116	13	99	30	268	28MAY–12AUG
<i>Aethalura intertexta</i> (Wlk.)	GEO	2	81	121	49	14	267	11MAY–03AUG
<i>Meganola minuscula</i> (Zell.)	NOC	30	76	1	159	0	266	01MAY–12AUG
<i>Eugonobapta nivosaria</i> (Gn.)	GEO	17	51	35	94	36	233	17JUN–08AUG
<i>Symmerista leucitys</i> Franc.	NOT	0	0	12	91	123	226	06JUN–27JUL
<i>Acronicta hasta</i> Gn.	NOC	44	41	6	79	55	225	01MAY–13AUG

continued

*GEO=Geometridae, ARC=Arctiidae, NOT=Notodontidae, LAS=Lasiocampidae, DRE=Drepanidae, NOC=Noctuidae, SAT=Saturniidae, SPH=Sphingidae, LYM=Lymantriidae, EPT=Epipleminae, THY=Thyatiridae, APA=Apatelodidae, MIM=Mimallonidae

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Heliomata cycladata</i> G. & R.	GEO	7	37	1	70	109	224	11MAY-12JUL
<i>Plagodis serinaria</i> H.-S.	GEO	13	56	1	40	109	219	13MAY-29JUN
<i>Bomolocha sordidula</i> (Grt.)	NOC	4	187	1	14	8	214	01MAY-12AUG
<i>Besma quercivoraria</i> (Gn.)	GEO	26	129	6	20	27	208	01MAY-13AUG
<i>Renia sobrialis</i> (Wlk.)	NOC	67	87	0	39	12	205	17JUN-12AUG
<i>Homochlodes disconventa</i> (Wlk.)	GEO	22	48	10	51	59	190	09MAY-12AUG
<i>Cabera erythemaria</i> Gn.	GEO	9	69	16	43	45	182	10MAY-12AUG
<i>Lochmaeus manteo</i> Doubleday	NOT	37	65	28	27	25	182	18MAY-13AUG
<i>Bomolocha baltimoralis</i> (Gn.)	NOC	51	70	1	39	21	182	07MAY-13AUG
<i>Eudryas grata</i> (F.)	NOC	34	72	16	23	23	168	03JUN-10AUG
<i>Polia latex</i> (Gn.)	NOC	33	64	6	39	18	160	10MAY-02AUG
<i>Heterocampa biundata</i> Wlk.	NOT	35	63	10	20	28	156	28MAY-13AUG
<i>Lymantria dispar</i> (L.)	LYM	14	16	37	38	45	150	04JUL-13AUG
<i>Pangrapta decoralis</i> Hbn.	NOC	4	85	18	29	14	150	28MAY-09AUG
<i>Symmerista albifrons</i> (J.E. Smith)	NOT	13	133	0	0	0	146	28MAY-23JUL
<i>Heterophleps triguttaria</i> H.-S.	GEO	9	92	7	29	8	145	13MAY-13AUG
<i>Zanclognatha lituralis</i> (Hbn.)	NOC	65	8	39	17	14	143	07JUN-03AUG
<i>Lithacodia carneola</i> (Gn.)	NOC	13	65	0	32	28	138	13MAY-12AUG
<i>Callopietria mollissima</i> (Gn.)	NOC	24	64	2	17	30	137	13MAY-13AUG
<i>Hydriomena divisaria</i> (Wlk.)	GEO	8	30	0	88	10	136	01MAY-04JUL
<i>Acronicta fragilis</i> Gn.	NOC	41	82	0	2	8	133	10MAY-23JUL
<i>Pero honestaria</i> (Wlk.)	GEO	31	11	11	28	44	125	01MAY-13AUG
<i>Biston betularia</i> (L.)	GEO	22	53	1	5	41	122	13MAY-13AUG
<i>Lambdina pellucidaria</i> (G. & R.)	GEO	17	72	10	19	4	122	06MAY-08AUG
<i>Acronicta haesitata</i> (Grt.)	NOC	7	44	1	14	56	122	20MAY-08AUG
<i>Anavitrinella pampinaria</i> (Gn.)	GEO	37	2	0	20	58	117	17JUN-09AUG
<i>Tetracis cachexiata</i> Gn.	GEO	6	61	6	19	20	112	13MAY-20JUL
<i>Lytrosis unitaria</i> (H.-S.)	GEO	4	41	10	15	40	110	10JUN-03AUG
<i>Polia nimbose</i> (Gn.)	NOC	3	22	3	42	40	110	17JUN-01AUG
<i>Lophocampa caryae</i> Harr.	ARC	13	66	2	8	17	106	13MAY-25JUN
<i>Idia americalis</i> (Gn.)	NOC	5	19	24	46	11	105	24MAY-13AUG
<i>Dasylophia thyatiroides</i> (Wlk.)	NOT	3	98	0	0	0	101	20MAY-13AUG
<i>Epimecis hortaria</i> (F.)	GEO	10	28	8	49	4	99	10MAY-09AUG
<i>Acronicta innotata</i> Gn.	NOC	16	38	1	12	31	98	21MAY-08AUG
<i>Cladara atroliturata</i> (Wlk.)	GEO	37	44	0	16	0	97	01MAY-03JUN
<i>Anorthodes tarda</i> (Gn.)	NOC	17	7	3	60	10	97	13MAY-20JUL
<i>Macrurocampa marthesia</i> (Cram.)	NOT	14	34	20	16	12	96	24JUN-09AUG
<i>Eulithis diversilineata</i> (Hbn.)	GEO	13	26	3	38	14	94	17JUN-13AUG
<i>Anagoga occiduaria</i> (Wlk.)	GEO	7	32	3	40	11	93	01MAY-13AUG
<i>Pero morrisonaria</i> (Hy. Edw.)	GEO	14	47	1	26	0	88	14MAY-12AUG
<i>Dasylophia anguina</i> (J.E. Smith)	NOT	8	11	2	9	55	85	11MAY-09AUG

continued

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Orthonama centrostrigaria</i> (Woll.)	GEO	9	36	16	20	2	83	11MAY-13AUG
<i>Acronicta inclara</i> Sm.	NOC	19	27	1	17	19	83	28MAY-27JUL
<i>Xestia dolosa</i> Franc.	NOC	30	22	2	17	12	83	10MAY-30JUL
<i>Trichodezia albovittata</i> (Gn.)	GEO	0	44	4	32	2	82	01MAY-12AUG
<i>Dasychira basiflava</i> (Pack.)	LYM	33	13	0	18	16	80	17JUN-13AUG
<i>Orthodes crenulata</i> (Butler)	NOC	61	15	0	0	0	76	21MAY-30JUL
<i>Dasychira dorsipennata</i> (B. & McD.)	LYM	9	36	4	23	2	74	10JUN-13AUG
<i>Orthosia rubescens</i> (Wlk.)	NOC	26	17	6	25	0	74	07MAY-14MAY
<i>Bomolocha deceptalis</i> (Wlk.)	NOC	26	33	1	4	7	71	01MAY-13AUG
<i>Antheraea polyphemus</i> (Cram.)	SAT	5	31	4	21	9	70	24MAY-30JUL
<i>Lithacodia muscosa</i> (Gn.)	NOC	0	65	0	1	1	67	03JUN-22JUL
<i>Crociographa normani</i> (Grt.)	NOC	19	7	1	27	13	67	06MAY-06JUN
<i>Eupithecia miserulata</i> Grt.	GEO	59	5	0	0	0	64	14MAY-06AUG
<i>Orgyia leucostigma</i> (J.E. Smith)	LYM	17	15	2	2	28	64	17JUN-13AUG
<i>Platysenta vecors</i> (Gn.)	NOC	16	29	2	10	5	62	01MAY-13AUG
<i>Dysstroma hersiliata</i> (Gn.)	GEO	5	26	9	15	4	59	28MAY-06JUL
<i>Calledapteryx dryopterata</i> Grt.	EPI	4	55	0	0	0	59	28MAY-13AUG
<i>Idia scobialis</i> (Grt.)	NOC	26	5	14	14	0	59	28MAY-13AUG
<i>Cissusa spadix</i> (Cram.)	NOC	4	1	2	50	0	57	01MAY-17MAY
<i>Polygrammate hebraeicum</i> Hbn.	NOC	3	15	0	1	37	56	28MAY-18JUL
<i>Acronicta americana</i> (Harr.)	NOC	12	3	7	21	12	55	17MAY-26JUL
<i>Nemoria bistriaria</i> Hbn.	GEO	0	26	5	19	4	54	10MAY-09AUG
<i>Catocala ultronia</i> (Hbn.)	NOC	6	16	1	21	10	54	08JUL-13AUG
<i>Baileya ophthalmica</i> (Gn.)	NOC	10	8	2	18	16	54	10MAY-09JUL
<i>Acronicta increta</i> Morr.	NOC	19	34	0	0	1	54	10JUN-15JUL
<i>Euparthenos nubilis</i> (Hbn.)	NOC	12	8	4	11	17	52	13MAY-12AUG
<i>Acronicta modica</i> Wlk.	NOC	3	49	0	0	0	52	03JUN-30JUL
<i>Plagodis alcoolaria</i> (Gn.)	GEO	8	10	2	14	17	51	10MAY-15JUL
<i>Cladara limitaria</i> (Wlk.)	GEO	0	11	11	29	0	51	01MAY-18MAY
<i>Pseudothyatira</i> <i>cymatophoroides</i> (Gn.)	THY	3	21	3	12	10	49	28MAY-12AUG
<i>Phalaenostola larentioides</i> Grt.	NOC	15	34	0	0	0	49	13MAY-13AUG
<i>Automeris io</i> (F.)	SAT	1	38	1	6	2	48	28MAY-04JUL
<i>Schizura ipomoeae</i> Doubleday	NOT	4	7	3	5	29	48	24JUN-08AUG
<i>Colocasia propinquilinea</i> (Grt.)	NOC	0	11	3	16	18	48	10MAY-08AUG
<i>Haploa lecontei</i> (Guer.- Meneville)	ARC	0	8	1	20	18	47	10JUN-19JUL
<i>Hypera humuli</i> Harr.	NOC	9	25	0	8	1	43	10MAY-13AUG
<i>Diarsia jucunda</i> (Wlk.)	NOC	6	5	14	12	6	43	28JUN-10AUG
<i>Nola triquetra</i> (Fitch)	NOC	0	38	2	0	2	42	01MAY-01AUG

continued

*GEO=Geometridae, ARC=Arctiidae, NOT=Notodontidae, LAS=Lasiocampidae,
DRE=Drepanidae, NOC=Noctuidae, SAT=Saturniidae, SPH=Sphingidae, LYM=Lymantriidae,
EPT=Epiplemidae, THY=Thyatiridae, APA=Apatelodidae, MIM=Mimallonidae

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Eutrapela clemataria</i> (J.E. Smith)	GEO	3	2	5	30	1	41	10MAY-06AUG
<i>Pyrrharctia isabella</i> (J.E. Smith)	ARC	10	26	0	2	2	40	14MAY-06AUG
<i>Hyperaeschra georgica</i> (H.-S.)	NOT	6	7	0	19	6	38	07MAY-08AUG
<i>Phalaenophana pyramusalis</i> (Wlk.)	NOC	20	18	0	0	0	38	03JUN-06AUG
<i>Zanclognatha pedipilalis</i> (Gn.)	NOC	2	1	16	19	0	38	16JUL-09AUG
<i>Balsa labecula</i> (Grt.)	NOC	8	15	4	0	11	38	13MAY-30JUL
<i>Plagodis phlogosaria</i> (Gn.)	GEO	7	5	2	20	3	37	10MAY-10AUG
<i>Prochoerodes transversata</i> (Drury)	GEO	14	5	10	0	8	37	24JUN-13AUG
<i>Cyclophora pendulinaria</i> (Gn.)	GEO	0	7	4	25	1	37	10MAY-12AUG
<i>Heterocampa umbrata</i> Wlk.	NOT	7	22	1	1	6	37	13MAY-08AUG
<i>Peridea ferruginea</i> (Pack.)	NOT	5	1	1	14	15	36	30MAY-08AUG
<i>Plathypena scabra</i> (F.)	NOC	6	10	1	10	9	36	30MAY-12AUG
<i>Ennomos subsignaria</i> (Hbn.)	GEO	19	9	7	0	0	35	15JUN-13AUG
<i>Epirrhoe alternata</i> (Müller)	GEO	5	6	0	0	24	35	07MAY-30JUL
<i>Habrosyne scripta</i> (Gosse)	THY	2	13	1	12	6	34	13MAY-12AUG
<i>Ectropis crepuscularia</i> (D. & S.)	GEO	15	17	0	2	0	34	01MAY-12JUL
<i>Metarranthis hypocharia</i> (H.-S.)	GEO	0	3	3	10	18	34	24MAY-06JUL
<i>Eubaphe mendica</i> (Wlk.)	GEO	0	11	8	4	11	34	03JUN-20JUL
<i>Idia diminuendis</i> (B. & McD.)	NOC	13	9	4	8	0	34	08JUL-09AUG
<i>Apatelodes torrefacta</i> (J.E. Smith)	APA	3	13	4	6	5	31	10JUN-03AUG
<i>Achatia distincta</i> Hbn.	NOC	2	1	2	21	5	31	07MAY-30MAY
<i>Pleuroprucha insularia</i> (Gn.)	GEO	0	30	0	0	0	30	03JUN-12AUG
<i>Dyspteris abortivaria</i> (H.-S.)	GEO	7	10	2	0	11	30	14MAY-06AUG
<i>Parallelia bistriaris</i> Hbn.	NOC	3	8	2	15	2	30	24MAY-09AUG
<i>Acronicta tristis</i> Sm.	NOC	0	3	0	27	0	30	03JUN-12JUL
<i>Cerma cerintha</i> (Tr.)	NOC	6	6	0	1	16	29	03JUN-16JUL
<i>Euchlaena obtusaria</i> (Hbn.)	GEO	17	7	0	3	1	28	10JUN-21JUN
<i>Oligocentria lignicolor</i> (Wlk.)	NOT	4	8	1	12	3	28	10JUN-09AUG
<i>Catocala blandula</i> Hulst	NOC	0	5	4	19	0	28	24JUN-10AUG
<i>Balsa malana</i> (Fitch)	NOC	6	0	0	22	0	28	10MAY-02AUG
<i>Glena cribrataria</i> (Gn.)	GEO	2	24	1	0	0	27	20MAY-08JUL
<i>Acronicta spinigera</i> Gn.	NOC	1	22	0	4	0	27	17MAY-12AUG
<i>Lithophane semiusta</i> Grt.	NOC	27	0	0	0	0	27	07MAY-21MAY
<i>Heterocampa obliqua</i> Pack.	NOT	0	1	0	6	19	26	06JUN-05AUG
<i>Nematocampa limbata</i> (Haw.)	GEO	2	6	1	8	8	25	17JUN-09AUG
<i>Lithophane hemina</i> Grt.	NOC	0	15	0	10	0	25	10MAY-13MAY
<i>Agrotis ipsilon</i> (Hufn.)	NOC	4	5	0	7	9	25	13MAY-09AUG
<i>Olceclostera angelica</i> (Grt.)	APA	5	18	0	1	0	24	17JUN-16JUL
<i>Dasychira obliquata</i> (G. & R.)	LYM	0	13	0	0	11	24	04JUL-12AUG
<i>Abagrotis alternata</i> (Grt.)	NOC	9	11	4	0	0	24	01JUL-12AUG

continues

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Protoaboarmia porcelaria</i> (Gn.)	GEO	3	19	0	0	1	23	13JUN–13AUG
<i>Catocala micronympha</i> Gn.	NOC	13	3	0	0	7	23	04JUL–13AUG
<i>Acronicta morula</i> G. & R.	NOC	0	3	0	1	19	23	10MAY–01AUG
<i>Pseudorthodes vecors</i> (Gn.)	NOC	18	1	3	1	0	23	09JUL–06AUG
<i>Lomographa semiclarata</i> (Wlk.)	GEO	2	2	4	14	0	22	01MAY–07JUN
<i>Nacophora quernaria</i> (J.E. Smith)	GEO	3	8	2	3	5	21	10MAY–16JUL
<i>Sicya macularia</i> (Harr.)	GEO	2	7	3	6	3	21	21JUN–20JUL
<i>Euthyatira pudens</i> (Gn.)	THY	13	5	1	1	0	20	01MAY–21MAY
<i>Orthonama obstipata</i> (F.)	GEO	1	13	1	5	0	20	24MAY–27JUL
<i>Amphipyra pyramidoides</i> Gn.	NOC	2	0	0	5	12	19	26JUL–13AUG
<i>Pseudaletia unipuncta</i> (Haw.)	NOC	6	0	6	4	3	19	29MAY–13AUG
<i>Caripeta divisata</i> Wlk.	GEO	0	4	5	8	1	18	05JUL–12AUG
<i>Thioptera nigrofimbria</i> (Gn.)	NOC	1	16	0	1	0	18	03JUN–13AUG
<i>Chytonix palliatricula</i> (Gn.)	NOC	1	3	0	5	9	18	28MAY–23JUL
<i>Itame subcessaria</i> (Wlk.)	GEO	4	1	0	7	5	17	28JUN–18JUL
<i>Actias luna</i> (L.)	SAT	0	7	1	4	5	17	01MAY–01AUG
<i>Esophropteryx thyatyroides</i> (Gn.)	NOC	6	3	1	5	2	17	28JUN–13AUG
<i>Elaphria festivoides</i> (Gn.)	NOC	2	7	1	4	3	17	24MAY–02JUL
<i>Ochropleura plecta</i> (L.)	NOC	0	10	0	5	2	17	06MAY–12AUG
<i>Euchlaena tigrinaria</i> (Gn.)	GEO	3	12	0	1	0	16	03JUN–18JUN
<i>Callosamia angulifera</i> (Wlk.)	SAT	1	15	0	0	0	16	17JUN–16JUL
<i>Ecliptopera atricolorata</i> (G. & R.)	GEO	5	6	1	2	1	15	01JUL–30JUL
<i>Anagrapha falcifera</i> (Kby.)	NOC	1	6	2	6	0	15	13MAY–26JUL
<i>Hyperstrotia pervertens</i> (B. & McD.)	NOC	1	14	0	0	0	15	10JUN–23JUL
<i>Oligia modica</i> (Gn.)	NOC	6	9	0	0	0	15	08JUL–13AUG
<i>Phosphila miselioides</i> (Gn.)	NOC	0	2	0	6	7	15	10MAY–12AUG
<i>Egira alternans</i> (Wlk.)	NOC	4	4	0	2	5	15	01MAY–30MAY
<i>Laothoe juglandis</i> (J.E. Smith)	SPH	2	1	1	8	2	14	28MAY–20JUL
<i>Lithacodia synochitis</i> (G. & R.)	NOC	1	13	0	0	0	14	10JUN–22JUL
<i>Acronicta impleta</i> Wlk.	NOC	0	1	1	0	12	14	23MAY–08AUG
<i>Plagodis fervidaria</i> (H.-S.)	GEO	6	1	0	0	6	13	14MAY–08AUG
<i>Tetracis crocallata</i> Gn.	GEO	2	6	1	2	2	13	13JUN–20JUL
<i>Xanthorhoe ferrugata</i> (Cl.)	GEO	1	12	0	0	0	13	13MAY–09JUL
<i>Darapsa myron</i> (Cram.)	SPH	3	4	2	2	2	13	10JUN–27JUL
<i>Zale lunifera</i> (Hbn.)	NOC	2	11	0	0	0	13	01MAY–14MAY
<i>Phlogophora iris</i> Gn.	NOC	1	4	1	3	4	13	21MAY–05JUL
<i>Feralia comstocki</i> (Grt.)	NOC	0	5	2	6	0	13	10MAY–13MAY
<i>Orthosia hibisci</i> (Gn.)	NOC	0	6	7	0	0	13	01MAY–18MAY
<i>Semiothisa ulsterata</i> (Pears.)	GEO	0	0	0	0	12	12	30MAY–08AUG

continued

*GEO=Geometridae, ARC=Arctiidae, NOT=Notodontidae, LAS=Lasiocampidae, DRE=Drepanidae, NOC=Noctuidae, SAT=Saturniidae, SPH=Sphingidae, LYM=Lymantriidae, EPT=Epiplemidae, THY=Thyatiridae, APA=Apatelodidae, MIM=Mimallonidae

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Ceratonia undulosa</i> (Wlk.)	SPH	1	1	2	6	2	12	17MAY-01AUG
<i>Acronicta interrupta</i> Gn.	NOC	11	1	0	0	0	12	17JUN-13AUG
<i>Elaphria chalconia</i> (Hbn.)	NOC	1	11	0	0	0	12	28MAY-13AUG
<i>Anticlea vasiliata</i> Gn.	GEO	0	1	0	9	1	11	01MAY-30MAY
<i>Deidamia inscripta</i> (Harr.)	SPH	2	0	1	3	5	11	11MAY-06JUN
<i>Scolecocampa liburna</i> (Gey.)	NOC	0	0	4	7	0	11	19JUL-27JUL
<i>Panopoda rufimargo</i> (Hbn.)	NOC	1	5	0	2	3	11	14JUN-12JUL
<i>Cicinnus melsheimeri</i> (Harr.)	MIM	0	2	1	2	5	10	28MAY-29JUN
<i>Haploa contigua</i> (Wlk.)	ARC	0	0	6	4	0	10	12JUL-20JUL
<i>Renia discoloralis</i> Gn.	NOC	0	3	4	3	0	10	17JUN-09AUG
<i>Palthis angulalis</i> (Hbn.)	NOC	0	3	1	5	1	10	13MAY-09AUG
<i>Catocala ilia</i> (Cram.)	NOC	1	0	0	7	2	10	12JUL-09AUG
<i>Charadra deridens</i> (Gn.)	NOC	5	3	0	1	1	10	13MAY-13AUG
<i>Achatodes zaeae</i> (Harr.)	NOC	4	1	2	2	1	10	08JUL-08AUG
<i>Lacinipolia renigera</i> (Steph.)	NOC	5	1	2	2	0	10	01JUN-30JUL
<i>Homorthodes furfurata</i> (Grt.)	NOC	0	3	0	5	2	10	24JUN-09AUG
<i>Metanema inatommata</i> (Gn.)	GEO	2	1	1	0	5	9	30MAY-01AUG
<i>Cepphis armataria</i> (H. & S.)	GEO	0	4	1	1	3	9	03JUN-20JUL
<i>Lambdina fervidaria</i> (Hbn.)	GEO	3	0	0	6	0	9	26JUL-13AUG
<i>Xanthorhoe lacustrata</i> (Gn.)	GEO	0	9	0	0	0	9	28MAY-10JUN
<i>Spilosoma latipennis</i> Strech	ARC	2	2	1	3	1	9	29MAY-23JUL
<i>Catocala amica</i> (Hbn.)	NOC	0	4	0	3	2	9	08JUL-12AUG
<i>Baileya levitans</i> (Sm.)	NOC	4	3	0	2	0	9	13MAY-26JUL
<i>Hyppa xylinoides</i> (Gn.)	NOC	1	1	0	7	0	9	21MAY-09AUG
<i>Xanthotype urticaria</i> Swett	GEO	1	1	1	4	1	8	28MAY-09AUG
<i>Xanthotype sospeta</i> (Drury)	GEO	1	3	0	0	4	8	30MAY-17JUN
<i>Antepione thisoaria</i> (Gn.)	GEO	2	4	2	0	0	8	13MAY-06AUG
<i>Eulithis explanata</i> (Wlk.)	GEO	3	3	1	1	0	8	17JUN-20JUL
<i>Horisme intestinata</i> (Gn.)	GEO	2	0	1	3	2	8	24MAY-06AUG
<i>Phyllodesma americana</i> (Harr.)	LAS	2	4	0	1	1	8	01MAY-23MAY
<i>Gluphisia septentrionis</i> Wlk.	NOT	2	1	0	2	3	8	13MAY-06AUG
<i>Lascoria ambigua</i> Wlk.	NOC	0	5	0	2	1	8	13MAY-02AUG
<i>Allotria elonympha</i> (Hbn.)	NOC	1	2	1	4	0	8	07JUN-26JUL
<i>Acronicta retardata</i> (Wlk.)	NOC	6	1	0	1	0	8	03JUN-12JUL
<i>Cosmia calami</i> (Harv.)	NOC	0	1	1	2	4	8	11JUL-03AUG
<i>Furcula borealis</i> (Guer.-Meneville)	NOT	2	2	0	1	2	7	01MAY-24MAY
<i>Euchaetes egle</i> (Drury)	ARC	1	3	0	0	3	7	10JUN-25JUL
<i>Idia lubricalis</i> (Gey.)	NOC	3	0	2	1	1	7	02JUL-02AUG
<i>Bomolocha madefactalis</i> (Gn.)	NOC	7	0	0	0	0	7	14MAY-06AUG
<i>Meganola spodia</i> Franclemont	NOC	0	7	0	0	0	7	17JUN-15JUL
<i>Acronicta pruni</i> Harr.	NOC	4	3	0	0	0	7	14MAY-24JUN
<i>Acronicta caesarea</i> Sm.	NOC	0	5	0	0	2	7	01JUL-04JUL
<i>Nedra ramosula</i> (Gn.)	NOC	0	2	0	5	0	7	10MAY-13MAY
<i>Elaphria versicolor</i> (Grt.)	NOC	4	0	0	3	0	7	14JUN-23JUL
<i>Oreta rosea</i> (Wlk.)	DRE	0	2	0	2	2	6	28MAY-01AUG

continued

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Melanolophia signataria</i> (Wlk.)	GEO	0	6	0	0	0	6	01MAY–15JUL
<i>Metarranthis duaria</i> (Gn.)	GEO	0	6	0	0	0	6	20MAY–28MAY
<i>Venusia cambrica</i> Curt.	GEO	3	3	0	0	0	6	06MAY–11JUN
<i>Datana ministra</i> (Drury)	NOT	3	2	0	1	0	6	17JUN–30JUL
<i>Haploa confusa</i> (Lyman)	ARC	0	0	0	6	0	6	19JUL–26JUL
<i>Holomelina opella</i> (Grt.)	ARC	3	1	1	1	0	6	28MAY–28JUN
<i>Lithacodia concinnimacula</i> (Gn.)	NOC	3	1	1	1	0	6	17MAY–08JUN
<i>Protorthodes oviduca</i> (Gn.)	NOC	0	3	1	2	0	6	28MAY–07JUN
<i>Euchlaena irraria</i> (B. & McD.)	GEO	0	5	0	0	0	5	01JUL–01JUL
<i>Zanclognatha jaccusalis</i> (Wlk.)	NOC	0	5	0	0	0	5	03JUN–01JUL
<i>Phoberia atomaris</i> Hbn.	NOC	0	1	2	2	0	5	01MAY–11MAY
<i>Zale lunata</i> (Drury)	NOC	0	5	0	0	0	5	13MAY–15JUL
<i>Caenurgina erechtea</i> (Cram.)	NOC	1	0	0	0	4	5	04JUL–13AUG
<i>Catocala neogama</i> (J.E. Smith)	NOC	0	5	0	0	0	5	12AUG–12AUG
<i>Melanchra adjuncta</i> (Gn.)	NOC	0	0	0	0	5	5	13JUN–27JUN
<i>Spaelotis clandestina</i> (Harr.)	NOC	0	0	2	2	1	5	20JUN–27JUL
<i>Eueretagrotis perattenta</i> (Grt.)	NOC	3	1	0	1	0	5	08JUL–23JUL
<i>Pyrrhia umbra</i> (Hufn.)	NOC	0	0	0	3	2	5	21JUN–25JUL
<i>Semiothisa granitata</i> (Gn.)	GEO	2	1	0	0	1	4	14MAY–01AUG
<i>Eufidonia notataria</i> (Wlk.)	GEO	0	1	0	0	3	4	13MAY–30MAY
<i>Euchlaena serrata</i> (Drury)	GEO	0	1	1	1	1	4	03JUN–06JUL
<i>Euchlaena pectinaria</i> (D. & S.)	GEO	0	0	4	0	0	4	15JUN–29JUN
<i>Tolype notialis</i> Franc.	LAS	0	0	0	2	2	4	01AUG–09AUG
<i>Clostera inclusa</i> (Hbn.)	NOT	0	1	1	1	1	4	10MAY–13JUN
<i>Spilosoma virginica</i> (F.)	ARC	1	1	0	2	0	4	28MAY–13AUG
<i>Palthis asopialis</i> (Gn.)	NOC	2	2	0	0	0	4	11JUN–13AUG
<i>Catocala resecta</i> Grt.	NOC	0	1	0	1	2	4	26JUL–12AUG
<i>Catocala andromedae</i> Gn.	NOC	1	2	0	1	0	4	22JUL–13AUG
<i>Baileya dormitans</i> (Gn.)	NOC	0	2	0	0	2	4	30MAY–01AUG
<i>Lithacodia musta</i> (G. & R.)	NOC	4	0	0	0	0	4	09JUL–30JUL
<i>Leuconycta diphteroides</i> (Gn.)	NOC	1	1	0	0	2	4	28MAY–16JUL
<i>Oligia crytora</i> (Franc.)	NOC	0	4	0	0	0	4	10JUN–24JUN
<i>Lacinipolia lorea</i> (Gn.)	NOC	1	2	0	0	1	4	03JUN–09JUL
<i>Hydriomena pluviata</i> (Gn.)	GEO	0	0	3	0	0	3	11MAY–11MAY
<i>Pheosia rimosa</i> Pack.	NOT	1	0	0	1	1	3	24MAY–08AUG
<i>Schizura leptinoides</i> (Grt.)	NOT	3	0	0	0	0	3	18JUN–16JUL
<i>Cisseps fulvicollis</i> (Hbn.)	ARC	0	3	0	0	0	3	01JUL–08JUL
<i>Phalaenostola metonalis</i> (Wlk.)	NOC	0	3	0	0	0	3	13MAY–13MAY
<i>Bomolocha abalienalis</i> (Wlk.)	NOC	3	0	0	0	0	3	09JUL–16JUL
<i>Catocala sordida</i> Grt.	NOC	0	0	1	0	2	3	18JUL–10AUG
<i>Allagrapha aerea</i> (Hbn.)	NOC	1	2	0	0	0	3	10JUN–18JUN

continued

*GEO=Geometridae, ARC=Arctiidae, NOT=Notodontidae, LAS=Lasiocampidae, DRE=Drepanidae, NOC=Noctuidae, SAT=Saturniidae, SPH=Sphingidae, LYM=Lymantriidae, EPT=Epiplemidae, THY=Thyatiridae, APA=Apatelodidae, MIM=Mimallonidae

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Marathyssa basalis</i> Wlk.	NOC	0	2	0	1	0	3	10MAY–13MAY
<i>Acronicta lobeliae</i> Gn.	NOC	1	0	0	2	0	3	17MAY–04JUN
<i>Polia imbrifera</i> (Gn.)	NOC	1	2	0	0	0	3	24JUN–09JUL
<i>Semiothisa minorata</i> (Pack.)	GEO	1	0	0	1	0	2	09JUL–09AUG
<i>Eusarca confusaria</i> Hbn.	GEO	0	1	0	1	0	2	08JUL–12JUL
<i>Nemoria lixaria</i> (Gn.)	GEO	1	0	1	0	0	2	16JUL–20JUL
<i>Nemoria mimosaria</i> (Gn.)	GEO	2	0	0	0	0	2	21MAY–21MAY
<i>Venusia comptaria</i> (Wlk.)	GEO	0	2	0	0	0	2	01MAY–01MAY
<i>Schizura unicornis</i> (J.E. Smith)	NOT	2	0	0	0	0	2	18JUN–18JUN
<i>Haploa clymene</i> (Brown)	ARC	1	0	0	1	0	2	26JUL–30JUL
<i>Ecpantheria scribonia</i> (Stoll)	ARC	0	2	0	0	0	2	17JUN–17JUN
<i>Cynia tenera</i> Hbn.	ARC	0	1	1	0	0	2	17JUN–29JUN
<i>Bleptina caradrinalis</i> Gn.	NOC	0	2	0	0	0	2	17JUN–17JUN
<i>Bomolocha bijugalis</i> (Wlk.)	NOC	1	1	0	0	0	2	03JUN–06AUG
<i>Zale undularis</i> (Drury)	NOC	0	1	0	0	1	2	13MAY–06JUN
<i>Catocala cerogama</i> Gn.	NOC	1	1	0	0	0	2	18JUN–12AUG
<i>Paectes oculatrix</i> (Gn.)	NOC	0	0	0	2	0	2	07JUN–09AUG
<i>Oruza albocostaliata</i> (Pack.)	NOC	0	1	0	0	1	2	24JUN–01AUG
<i>Acronicta funeralis</i> G. & R.	NOC	0	0	0	2	0	2	21JUN–28JUN
<i>Amphipoea velata</i> (Wlk.)	NOC	0	1	1	0	0	2	06JUL–22JUL
<i>Galgula partita</i> Gn.	NOC	0	1	0	1	0	2	24JUN–19JUL
<i>Polia goodelli</i> (Grt.)	NOC	0	2	0	0	0	2	08JUL–22JUL
<i>Leucania phragmatidicola</i> Gn.	NOC	0	1	0	0	1	2	17JUN–20JUN
<i>Ulolonche culea</i> (Gn.)	NOC	0	0	0	0	2	2	30MAY–30MAY
<i>Xestia normaniana</i> (Grt.)	NOC	1	1	0	0	0	2	12AUG–13AUG
<i>Anaplectoides prasina</i> (D. & S.)	NOC	2	0	0	0	0	2	06AUG–13AUG
<i>Heptagrotis phyllophora</i> (Grt.)	NOC	0	1	0	1	0	2	17JUN–12JUL
<i>Pyrrhia exprimens</i> (Wlk.)	NOC	1	0	1	0	0	2	09JUL–03AUG
<i>Cleora sublunaria</i> (Gn.)	GEO	1	0	0	0	0	1	14MAY–14MAY
<i>Selenia kentaria</i> (G. & R.)	GEO	0	0	0	1	0	1	07JUN–07JUN
<i>Cyclophora packardi</i> (Prout)	GEO	0	0	0	1	0	1	09AUG–09AUG
<i>Dysstroma truncata</i> (Hufn.)	GEO	0	0	1	0	0	1	06JUL–06JUL
<i>Mesoleuca ruficillata</i> (Gn.)	GEO	0	0	1	0	0	1	10AUG–10AUG
<i>Anticlea multifera</i> (Wlk.)	GEO	0	0	0	1	0	1	10MAY–10MAY
<i>Xanthorhoe labradorensis</i> (Pack.)	GEO	0	0	0	1	0	1	10MAY–10MAY
<i>Lacosoma chiridota</i> Grt.	MIM	0	0	0	0	1	1	20JUN–20JUN
<i>Ceratomia amyntor</i> (Geyer)	SPH	0	0	0	0	1	1	20JUN–20JUN
<i>Paonias astylus</i> (Drury)	SPH	0	0	0	1	0	1	19JUL–19JUL
<i>Clostera albosigma</i> Fitch	NOT	0	0	1	0	0	1	08JUN–08JUN
<i>Schizura concinna</i> (J.E. Smith)	NOT	0	0	0	0	1	1	27JUN–27JUN
<i>Zanclognatha cruralis</i> (Gn.)	NOC	0	0	0	1	0	1	24MAY–24MAY
<i>Rivula propinqualis</i> Gn.	NOC	0	1	0	0	0	1	28MAY–28MAY
<i>Metalectra discalis</i> (Grt.)	NOC	0	1	0	0	0	1	10JUN–10JUN
<i>Panopoda carneicosta</i> Gn.	NOC	0	0	0	0	1	1	11JUL–11JUL
<i>Celiptera frustulum</i> Gn.	NOC	0	1	0	0	0	1	05AUG–05AUG

continued

Species	Family*	1990	1991	1992	1993	1994	Total	Inclusive Flight Range
<i>Catocala habilis</i> Grt.	NOC	1	0	0	0	0	1	13AUG–13AUG
<i>Catocala obscura</i> Stkr.	NOC	0	0	0	0	1	1	08AUG–08AUG
<i>Catocala residua</i> Grt.	NOC	1	0	0	0	0	1	13AUG–13AUG
<i>Catocala palaeogama</i> Gn.	NOC	0	0	0	0	1	1	01AUG–01AUG
<i>Catocala relict</i> a Wlk.	NOC	1	0	0	0	0	1	06AUG–06AUG
<i>Catocala crataegi</i> Saund.	NOC	1	0	0	0	0	1	09JUL–09JUL
<i>Abrostola ovalis</i> Gn.	NOC	1	0	0	0	0	1	21MAY–21MAY
<i>Diachrysia aeroides</i> (Grt.)	NOC	1	0	0	0	0	1	16JUL–16JUL
<i>Chrysanympha formosa</i> (Grt.)	NOC	0	0	0	0	1	1	04JUL–04JUL
<i>Syngrapha rectangula</i> (W. Kby.)	NOC	0	0	1	0	0	1	06JUL–06JUL
<i>Panthea furcilla</i> (Pack.)	NOC	0	1	0	0	0	1	08JUL–08JUL
<i>Agriopodes fallax</i> (H.-S.)	NOC	0	1	0	0	0	1	08JUL–08JUL
<i>Agriopodes teratophora</i> (H.-S.)	NOC	0	1	0	0	0	1	17JUN–17JUN
<i>Apamea cariosa</i> (Gn.)	NOC	0	1	0	0	0	1	10JUN–10JUN
<i>Oligia fractilinea</i> (Grt.)	NOC	0	1	0	0	0	1	12AUG–12AUG
<i>Iodopepla u-album</i> (Gn.)	NOC	0	1	0	0	0	1	22JUL–22JUL
<i>Phlogophora periculosa</i> Gn.	NOC	0	1	0	0	0	1	12AUG–12AUG
<i>Ipimorpha pleonectusa</i> Grt.	NOC	1	0	0	0	0	1	13AUG–13AUG
<i>Callopietria cordata</i> (Ljungh)	NOC	1	0	0	0	0	1	18JUN–18JUN
<i>Elaphria grata</i> Hbn.	NOC	0	1	0	0	0	1	10JUN–10JUN
<i>Ogdoconta cinereola</i> (Gn.)	NOC	1	0	0	0	0	1	13AUG–13AUG
<i>Lithophane querquera</i> Grt.	NOC	1	0	0	0	0	1	14MAY–14MAY
<i>Lithophane antennata</i> (Wlk.)	NOC	0	0	0	1	0	1	10MAY–10MAY
<i>Pyreferra hesperidago</i> (Gn.)	NOC	0	0	1	0	0	1	11MAY–11MAY
<i>Eupsilia vinulenta</i> (Grt.)	NOC	0	0	1	0	0	1	11MAY–11MAY
<i>Eupsilia sidus</i> (Gn.)	NOC	0	1	0	0	0	1	01MAY–01MAY
<i>Psaphida resumens</i> Wlk.	NOC	0	1	0	0	0	1	01MAY–01MAY
<i>Copivaleria grotei</i> (Morr.)	NOC	0	1	0	0	0	1	13MAY–13MAY
<i>Leucania ursula</i> (Fbs.)	NOC	1	0	0	0	0	1	30JUL–30JUL
<i>Orthosia revicta</i> (Morr.)	NOC	0	0	0	1	0	1	10MAY–10MAY
<i>Morrisonia evicta</i> (Grt.)	NOC	1	0	0	0	0	1	21MAY–21MAY
<i>Tricholita signata</i> (Wlk.)	NOC	0	0	1	0	0	1	10AUG–10AUG
<i>Peridroma saucia</i> (Hbn.)	NOC	0	1	0	0	0	1	10JUN–10JUN
<i>Xestia bicarnea</i> (Gn.)	NOC	0	1	0	0	0	1	12AUG–12AUG
<i>Euretagrotis attenta</i> (Grt.)	NOC	1	0	0	0	0	1	16JUL–16JUL

*GEO=Geometridae, ARC=Arctiidae, NOT=Notodontidae, LAS=Lasiocampidae, DRE=Drepanidae, NOC=Noctuidae, SAT=Saturniidae, SPH=Sphingidae, LYM=Lymantriidae, EPT=Epiplemidae, THY=Thyatiridae, APA=Apatelodidae, MIM=Mimallonidae

Agricultural and Forestry
Experiment Station
West Virginia University
PO Box 6108
Morgantown, WV 26506-6108

Non-Profit Organization
U.S. Postage
PAID
Morgantown, WV
Permit No. 34

SERIALS DEPARTMENT
EVANSDALE LIBRARY
WEST VIRGINIA UNIVERSITY
PO BOX 6105
MORGANTOWN WV 26506-6105