1-1-1995

Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain section.

Linda Butler

Follow this and additional works at: [https://researchrepository.wvu.edu/
wv_agricultural_and_forestry_experiment_station_bulletins](https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins)

Digital Commons Citation

Butler, Linda, "Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain section." (1995). *West Virginia Agricultural and Forestry Experiment Station Bulletins*. 713.
https://researchrepository.wvu.edu/wv_agricultural_and_forestry_experiment_station_bulletins/604

This Bulletin is brought to you for free and open access by the Davis College of Agriculture, Natural Resources And Design at The Research Repository @ WVU. It has been accepted for inclusion in West Virginia Agricultural and Forestry Experiment Station Bulletins by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

Canopy Arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain Section: II. Macrolepidopterous Larvae on Foliage and Under Burlap Bands

RECEIVED

JAN 24 1996

EVANS DALE LIBRARY
WEST VIRGINIA UNIVERSITY

S127
E1
713
Nov. 1995

Bulletin 713 • November 1995
Agricultural and Forestry Experiment Station
West Virginia University

Authors

Linda Butler is Professor of Entomology and Vicki Kondo and Gregory Chrislip are Entomology Research Assistants, Division of Plant and Soil Sciences, College of Agriculture and Forestry, West Virginia University.

This research was supported by a grant from the USDA Forest Service. We would like to acknowledge the assistance and cooperation of the USDA Forest Service Appalachian Integrated Pest Management Program personnel, our undergraduate field and laboratory assistants and the data management assistance of Dr. Edwin Townsend.

West Virginia University
Agricultural and Forestry Experiment Station
College of Agriculture and Forestry
Rosemary Haggett, Director
Morgantown, WV

Canopy Arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain Section: II. Macrolepidopterous Larvae on Foliage and Under Burlap Bands

Linda Butler, Gregory Chrislip and Vicki Kondo

INTRODUCTION

In recent years, we have conducted a number of studies in West Virginia on species composition of the macrolepidoptera in canopies of hardwood forests. These studies have been for the collection of baseline data only (Butler 1992) and for the evaluation of impacts of gypsy moth suppression efforts (Butler and Kondo 1993; Sample et al. 1995).

In 1989 we began a study to evaluate non-target impact of the insect growth regulator Dimilin (1-(4 chlorophenyl)-3-(2, 6-diflubenzoyl) urea) or diflubenzuron on canopy arthropods. Dimilin has been applied to hundreds of thousands of acres in the United States for suppression of gypsy moth, *Lymantria dispar* (L.) (Lepidoptera: Lymantriidae) (USDA Forest Service 1990). From 1989 through 1994 we sampled canopy arthropods in four closed watersheds on the Fernow Experimental Forest near Parsons in the Allegheny Mountain section of West Virginia. In 1992 Dimilin was applied to two of the watersheds. Elsewhere we present results of impact of Dimilin on canopy arthropods and evaluation of macrolepidopterous moths sampled by blacklight trap. Here we present richness and abundance of macrolepidoptera larvae taken from foliage and collected from burlap bands on mixed birch (*Betula* spp.), black cherry (*Prunus serotina* Ehrh.), mixed maples (*Acer* spp.) and mixed oaks (*Quercus* spp.) during the six-year study period. Used in this context, richness refers to number of species of larvae, abundance refers to total numbers.

MATERIALS AND METHODS

Study Sites

The Fernow Experimental Forest is located on the unglaciated Allegheny Plateau (latitude 39° 05'N, longitude 79° 41'W) at elevations ranging from 533 to 1,112 m. The topography is rugged with slopes of 10 to 60 percent. The experimental forest encompasses 1,902 ha and has been divided into a series of water-

sheds for research purposes. This study was conducted on watersheds 1, 4, 7, and 13.

Prior to establishment of the Fernow Experimental Forest (FEF) in 1934, the land was heavily cut between 1905 and 1910. Additional timbering occurred on several of the four watersheds from the early 1950s through the late 1970s. Currently the common tree species on the better sites include yellow-poplar (*Liriodendron tulipifera* L.), sugar maple (*Acer saccharum* Marsh.), black cherry, white ash (*Fraxinus americana* L.), American basswood (*Tilia americana* L.) and red oak (*Quercus rubra* L.). Dominant tree species on the poorer sites include various species of oak (*Quercus* spp.), hickory (*Carya* spp.), sourwood (*Oxydendrum arboreum* (L.) DC) and sassafras (*Sassafras albidum* (Nutt.) Nees) (Adams et al. 1993).

Importance values (IV) for tree species on watersheds 1, 4, 7, and 13 were determined by Stephenson 1992 (unpublished) in which he evaluated species on five small plots (100 m²) within each watershed, well away from watershed boundaries. All plants with stems greater than 2.5 cm DBH (diameter at breast height) were considered to be trees. Individual tree species were given an importance value by adding the relative basal area plus relative density and dividing by two. Stephenson's importance values are used to characterize tree composition on the four watersheds that comprised this study.

Watershed 1 is 30.1 ha with a NE aspect. It has not been timbered since 1958. Tree species with the highest importance values were sugar maple, basswood, beech (*Fagus grandifolia* Ehrh.), black locust (*Robinia pseudoacacia* L.), cucumber magnolia (*Magnolia acuminata* L.) and red oak. Importance values for the four species groups sampled in this study were 29.2 for sugar maple, 2.6 and 1.5 for yellow birch (*Betula lutea* Michx. f.) and black birch (*Betula lenta* L.), respectively, and 3.7 for red oak. Red maple (*Acer rubrum* L.) and black cherry were sparse on watershed 1.

Watershed 4 is 38.7 ha with an ESE aspect. Most abundant tree species with respect to importance values were beech, sugar maple, moosewood (*Acer pensylvanicum* L.), yellow-poplar, American basswood and red oak. Importance values for trees sampled in this study were 16.6 and 2.4 for sugar maple and red maple, respectively, 1.0 for black birch and 7.4 for red oak. Black cherry was represented on watershed 4 by a few large trees and abundant young trees in the understory, but the young trees were too small to have been counted in Stephenson's importance values. Watershed 4 is a relatively mature forest not having been timbered since 1910.

Watershed 7 is 24.2 ha with an E aspect. It is the youngest of the forest stands, having been last cut and treated with herbicides in 1969. Most abundant species (IV) were black birch, sugar maple, red oak, and yellow-poplar. Importance values of trees sampled in this study were 15.4 for sugar maple, 9.2 for red oak, 33.3 for black birch and 8.4 for black cherry.

Watershed 13 is 14.2 ha with a NNE aspect. Light-selective timbering of this watershed occurred until 1960, but it is a relatively mature stand. Predominant species (IV) were sugar maple, red oak, yellow-poplar, black cherry and black birch. Importance values for species utilized in this study were 30.5 for sugar maple, 1.1 for red maple, 29.3 for red oak, 6.4 for black birch and 6.6 for black cherry.

Sampling Methods

We began sampling macrolepidopterous larvae on all four watersheds in 1989 by means of burlap banding of trees and pole pruning of foliage.

Burlap bands 24 cm wide were stapled at breast height to trees. Ten trees each of black birch and black cherry were banded in late April 1989 on all four watersheds. Ten mixed maples were added on each watershed on July 5, 1989. Ten mixed oaks per watershed were added in early May 1990, giving a total of 160 banded trees for the study. Each year bands were installed in late April and sampling was begun early to mid-May: May 4, 1989; May 14, 1990; May 6, 1991; May 11, 1992; May 10, 1993; and May 9, 1994. Bands were left in place until early to mid-August each year and then removed after the last sampling date: August 15, 1989; August 13, 1990; August 12, 1991; August 10, 1992; August 9, 1993; and August 8, 1994. Each week all bands were checked and all macrolepidoptera larvae removed and returned to the laboratory for identification to species. Burlap bands mimic naturally occurring bark flaps and provide a refuge for sampling caterpillars that move up and down the trunks.

The first foliage samples were taken May 17, 1989 and consisted of two 25-branch tip samples of mixed birch (black and yellow) and two of black cherry on each of the four watersheds, for a total of 16 samples per sampling date. On June 27, 1989 a similar sampling procedure was added for mixed maple (sugar and red) so that 24 samples were being taken for the remainder of the season. Sampling of mixed oak (primarily red oak) was added at the beginning of the 1990 sampling season, giving a total of 32 samples of foliage on each sampling date.

Foliage sampling was begun soon after leaf expansion each spring: May 17, 1989; May 15, 1990; May 7, 1991; May 12, 1992; May 11, 1993; and May 8, 1994 and was continued until mid-August each year. During the six years of the study, foliage samples were taken each week for a total of 15 sampling dates each season. Foliage was collected from low- to mid-canopy with pole pruners that were equipped with large plastic catch bags to collect the foliage that fell.

The 32 foliage samples of mixed birch, black cherry, mixed maple and mixed oak were returned to the laboratory where all macrolepidoptera larvae were removed and identified to species by spot identification, comparison with voucher specimens in the WVU Arthropod Collection, comparison with color slides of macrolepidoptera prepared in our laboratory or by rearing the larva to the adult moth for positive identification.

After larvae were removed, all leaves were removed from twigs in each sample, placed in paper bags, oven dried, and weighed. Samples averaged about 50 grams dry leaf weight.

On May 16, 1992, Dimilin 4L was applied to watersheds 1 and 13 at a rate of 0.5 oz. AI/acre. Elsewhere we will discuss impact of that treatment on macrolepidopterous larvae. Here we present overall richness and abundance of larvae during the six-year study on the Fernow and present information on seasonal range and host associations.

Voucher specimens from the study are deposited in the WVU Arthropod Collection. Species are listed according to the checklist given by Hodges et al. (1983).

RESULTS

A total of 86 species of macrolepidopterous larvae representing 10 families was taken from bands and 111 species from 13 families were collected on foliage from 1989 through 1994 (Table 1). Most larvae collected by both methods were Noctuidae and Geometridae. Under bands, noctuid richness (39 species) was highest with geometrids contributing 24 species. On foliage geometrid richness (43 species) was highest with noctuids representing 36 species. The third richest family was Notodontidae with 12 species on foliage and 6 species under bands. While more families of macrolepidoptera were taken on foliage, sampling by bands allowed evaluation of several species not encountered on foliage, particularly with noctuids and arctiids (Table 1).

The most abundant larva under bands was the gypsy moth *Lymantria dispar* (Table 2). When the study began in 1989 almost no *L. dispar* were found, but their numbers gradually built up and became noticeably abundant by 1994. Not unexpected was the fact that most gypsy moth larvae were found under bands of oaks, their preferred hosts. Of the remaining species of larvae the most abundant were the noctuids *Polia latex* (Gn.), *Orthosia rubescens* (Wlk.), *Abagrotis alternata* (Grt.), *Lithophane hemina* Grt., *L. innominata* (Sm.) and *Zale minerea* (Gn.) and the lasiocampids *Malacosoma disstria* Hbn. and *M. americanum* (F.).

Since burlap bands mimic bark flaps on trees and provide insects with shelter, associating a species of caterpillar with a specific host food tree on the basis of catches under bands is problematic. Many of the species taken under bands are herbivorous on a number of deciduous host trees (Table 2). This is reflected by relatively even numbers of larvae for the host species groups; examples of generalists include *Orthosia rubescens*, *Lithophane hemina* and *Malacosoma disstria*. *Polia latex* is considered a generalist (Wood and Butler 1989) but it was taken in much greater abundance under bands on cherry trees in this study. *Abagrotis alternata* is a general feeder on many tree species (Forbes 1954) but frequently feeds on blueberry (*Vaccinium* spp.) and other low woody plants and

climbs trees to seek shelter. *Malacosoma americanum* strongly prefers black cherry but was found commonly under bands of all tree species in the study.

Most species of larvae recorded from tree bands in this study were in very low numbers. Sixty-one of the 86 species were represented by fewer than 20 individuals over the six-year period and 25 species were recorded as single individuals.

Species of macrolepidopterous larvae taken from foliage of the four tree species groups are listed in Table 3. Five species of geometrid larvae were the most abundant: *Lomographa glomeraria* (Grt.), *Erannis tiliaria* (Harr.), *L. vestaliata* (Gn.), *Melanolophia canadaria* (Gn.) and *Alsophila pometaria* (Harr.). Two noctuids, *Orthosia hibisci* (Gn.) and *Polia latex*, were next most abundant. Of the 17 species represented by more than 100 individuals during the study, 10 were geometrids, 6 were noctuids and one was a lymantriid, the gypsy moth.

Several of the more abundant species are generalists on a number of tree species: *Erannis tiliaria*, *Melanolophia canadaria*, *Alsophila pometaria*, *Orthosia hibisci* and *Polia latex*, for example. *Erannis tiliaria* and *A. pometaria* tend to be outbreak species (Butler 1985), but their numbers were stable and generally low at the FEF.

A number of species showed strong food tree preferences. *Lomographa glomeraria*, *L. vestaliata*, *L. semiclarata* (Wlk.), and *Hydria prunivorata* (Fgn.) are black cherry feeders; *Itame pustularia* (Gn.) and *Symmerista leucitys* Franc. feed on maple and *Catocala ilia* (Cram.) feeds on oak.

Seventy-one of the 111 species on foliage were represented by fewer than 20 individuals; 16 species were represented by a single caterpillar.

The highest richness of larvae on foliage was recorded for mixed oak with 72 species (Table 4). Black cherry accounted for 71 species and had the greatest abundance of larvae at 3,188, primarily because of the large numbers of *Lomographa* spp. Identical species richness (57 species) was noted for black birch and mixed maple. Abundance of larvae was similar among black birch, mixed maple and mixed oak (Table 4). Gypsy moth larvae accounted for 284 of the 1,387 individuals on oak.

Dates of first collection and last collection are given for all species of caterpillars under bands and on foliage (Tables 2, 3). Some larvae occurred throughout most of the May–August period and represent species that are univoltine and abundant, but are not highly synchronized for dates of moth emergence and oviposition. Therefore larvae are hatching over a several week period; *Polia latex* and *Morrisonia confusa* (Hbn.) are examples. Long collection periods are also expected for multivoltine species (species with more than one generation per year), such as the geometrids *Melanolophia canadaria* and *Lomographa vestaliata*. Species that are more restricted to a portion of the season include the spring defoliating geometrids and noctuids such as *Erannis tiliaria*, *Alsophila pometaria* and *Orthosia hibisci*.

DISCUSSION

A total of 130 species of macrolepidopterous larvae were identified during this study (86 from bands and 111 from foliage). Richness and abundance of macrolepidopterous caterpillars are comparable to that reported from other recent studies in West Virginia (Butler 1992; Butler et al. 1995a).

In 1984 and 1985 a baseline study was conducted at Coopers Rock State Forest in which 100 species of macrolepidopterous larvae in 14 families were identified from foliage of the same four tree species groups utilized in the Fernow study. A higher number of species of both noctuids and geometrids primarily accounts for the greater overall richness in the Fernow study.

Ninety-one species of macrolepidopterous caterpillars from 10 families were identified from oak foliage in the *Bacillus thuringiensis kurstaki* (*Btk*) non-target impact study conducted in Pendleton and Grant counties in West Virginia's Eastern Ridge and Valley section in 1990–1992 (Sample et al. 1995; Butler et al. 1995a). The 91 species on oak alone is considerably higher than the 61 species on oak at Coopers Rock and the 72 species on oak at Fernow; however, the sampling intensity of oak in the *Btk* study exceeded that of the Coopers Rock and current study by 2–3 fold. Table 3 indicates the species taken in the current study that were not represented at one or both sites of the two earlier studies. The comparison for Grant and Pendleton counties includes only those oak species for which oak is an expected host.

Coopers Rock and Fernow are both within West Virginia's Allegheny Mountain section and we might expect the caterpillar fauna to be similar. The 38 species taken on foliage at Fernow but not at Coopers Rock include 15 species of geometrids, 18 species of noctuids, a lasiocampid, a notodontid, a lycaenid, and two nymphalids (Table 3). We note that the Coopers Rock study was conducted for two years and the Fernow study for six, but during each year, samples were collected until October at Coopers Rock but only until mid-August at Fernow. Twenty-four of the 38 species not collected at Coopers Rock were represented at Fernow by three or fewer individuals over the six years. One would expect that species that are in very low abundance may have been taken at Coopers Rock with either greater sampling intensity during each of the two years, or by extending the study for more years. All of the 38 species not taken at Coopers Rock are expected to be there with the possible exception of the noctuid *Copipanolis styracis* (Gn.), which has only been collected commonly in the more eastern oak-pine habitats of the state (Butler et al. 1995a).

Species taken at Coopers Rock that were not represented at Fernow include the hesperiid *Erynnis juvenalis* (F.); the geometrids *Semiothisa aemulataria* (Wlk.), *Paleacrita merriccata* Dyar and *Anagoga occiduarua* (Wlk.); the apatelodid *Apatelodes torrefacta* (J.E. Smith); the saturniids *Eacles imperialis* (Drury), *Anisota senatoria* (J.E. Smith), *Automeris io* (F.), *Antheraea polyphemus* (Cram.),

Callosamia promethea (Drury) and *Callosamia angulifera* (Wlk.); the notodontids *Hyperaeschra georgica* (H.-S.) and *Oligocentria lignicolor* (Wlk.); the arctiids *Halysidota harrisii* Walsh and *Lophocampa caryae* Harr.; and the noctuids *Panopoda rufimargo* (Hbn.), *Cerma cerintha* (Tr.), *Acronicta increta* Morr., *A. inclara* Sm., *A. retardata* (Wlk.), *A. impleta* Wlk., *Phosphila turbulenta* Hbn. and *Himella intractata* (Morr.). All of these species were taken at Coopers Rock in very low numbers and/or late in the season, i.e., August–October when samples were not taken at Fernow.

Eighty-six species of macrolepidopterous larvae were taken from bands at Fernow from 1989–1994. In a Dimilin impact study at Coopers Rock in 1990 and 1991, 42 species of larvae representing eight families were taken from bands (Butler and Kondo 1993). The Coopers Rock study employed 80 banded trees from mid-April until mid-September each of the two years in comparison with 160 banded trees evaluated from mid-May until mid-August for six years at Fernow. Some similar results are noted with respect to the composition of the most abundant species. At both sites, the gypsy moth, *Lymantria dispar* was most abundant. *Polia latex*, which was second most abundant at Fernow, was third most abundant at Coopers Rock. Other examples of more abundant species at Fernow are ranked with comparisons of rank for species at Coopers Rock: *Orthosia rubescens* (3 versus 16), *Abagrotis alternata* (4 versus 2), *Lithophane hemina* (5 versus 19), *Malacosoma disstria* (6 versus 14), *Zale minerea* (8 versus 21), and *Malacosoma americanum* (9 versus 10). *Lithophane innominata*, the seventh most abundant species at Fernow, was not taken under bands at Coopers Rock.

Relatively abundant species taken at Coopers Rock that were relatively less abundant at Fernow include *Orthosia hibisci* (number 4 at Coopers Rock, number 15 at FEF), *Halysidota tessellaris* (J.E. Smith) (5 versus 24), *Epiglaea decliva* (Grt.) (6 versus 14), *Orgyia leucostigma* (J.E. Smith) (7 versus 36), *Polia nimbosa* (Gn.) (8 versus 60), *Dasychira basiflava* (Pack.) (9 versus 32) and *Dasychira dorsipennata* (B. & McD.) (11 versus 28). The noctuid *Polia detracta* (Wlk.) was the only species taken under bands at Coopers Rock that was not represented either under bands or on foliage at Fernow.

During 1990–1994 we sampled macrolepidopterous moths on the four watersheds at Fernow by means of blacklight traps (Butler et al. 1995b). The most abundant moths were *Itame pustularia*, *Hydrelia inornata* (Hulst), *Lomographa glomeraria*, *L. vestaliata*, *Halysidota tessellaris*, *Probole amicaria* (H.-S.), *Melanolophia canadaria* and *Iridopsis larvaria* (Gn.). Five of the 12 most abundant larval species on foliage are represented here in this group.

A total of 376 species of macrolepidopterous moths was collected at Fernow (Butler et al. 1995b); in this paper we have listed 130 species of larvae on four host plant groups. Some species of moths were not represented in larval collections because we did not sample the appropriate host. For example, *Euphyia*

unangulata (Haw.) larvae feed on chickweed (Forbes 1948), and *Clemensia albata* Pack. larvae feed on dead leaves (Forbes 1960). A number of species of larvae were collected that were not represented as adults in light traps, for example, *Erannis tiliaria* and *Alsophila pometaria*, whose males fly in November (females are flightless) and *Phigalia titea*, whose males fly in March (females are flightless).

This six year study of macrolepidopterous larvae on foliage and under tree bands at Fernow resulted in the identification of 130 species of larvae on the study plots, indicating a high diversity of species.

SUMMARY

Macrolepidopterous larvae were collected from foliage and bands of black birch, black cherry, mixed maples and mixed oaks from May–August 1989–1994. Collections resulted in 130 species of larvae. No species were sufficiently abundant to be considered in outbreak mode, although gypsy moth larval populations were building in the watersheds. Little defoliation was evident. While foliage pruning and tree banding produced 111 and 86 species of larvae respectively, the sampling methods complement each other for evaluating species richness.

LITERATURE CITED

- Adams, M.B., J.N. Kochenderfer, F. Wood, T.R. Angradi and P. Edwards. 1993. Forty years of hydrometeorological data from the Fernow Experimental Forest, West Virginia. USDA Forest Service. Gen. Tech. Rpt. NE-184. 24 pp.
- Butler, L. 1985. Biology of the half-wing geometer, *Phigalia titea* Cramer (Geometridae), as a member of a looper complex in West Virginia. J. Lepidop. Soc. 39:177–186.
- Butler, L. 1992. The community of macrolepidopterous larvae at Coopers Rock State Forest, West Virginia: a baseline study. Can. Entomol. 124:1149–1156.
- Butler, L. and V. Kondo. 1993. Impact of Dimilin on non-target Lepidoptera: Results of an operational gypsy moth suppression program at Coopers Rock State Forest, West Virginia. WV Agric. and For. Exp. Sta. Bull. 710. 21 pp.
- Butler, L., C. Zivkovich, and B.E. Sample. 1995a. Richness and abundance of arthropods in the oak canopy of West Virginia's Eastern Ridge and Valley Section during a study of impact of *Bacillus thuringiensis* with emphasis on macrolepidoptera larvae. WV Agric. and For. Exp. Sta. Bull. 711.

Butler, L., V. Kondo, and G. Chrislip. 1995b. Canopy arthropods at Fernow Experimental Forest in West Virginia's Allegheny Mountain Section: I. Macrolepidopterous moths collected by blacklight trap during a Dimilin impact study. WV Agric. and For. Exp. Sta. Bull. 712. 17 pp.

Forbes, W.T.M. 1948. Lepidoptera of New York and Neighboring States, Part II, Cornell University Agriculture Experiment Station Memoir 274; 263 pp.

Forbes, W.T.M. 1954. Lepidoptera of New York and Neighboring States, Part III, Cornell University Agriculture Experiment Station Memoir 329; 433 pp.

Forbes, W.T.M. 1960. Lepidoptera of New York and Neighboring States, Part IV, Cornell University Agriculture Experiment Station Memoir 371; 188 pp.

Hodges, R.W., T. Dominic, D.R. Davis, D.C. Ferguson, J.G. Franclemont, E.G. Munroe, and J.A. Powell. 1983. Checklist of the Lepidoptera of America North of Mexico. London. E.W. Classey Ltd. and Washington, D.C. Wedge Entomol. Res. Found. 284 pp.

Sample, B.E., L. Butler, C. Zivkovich, R.C. Whitmore, and R. Reardon. 1995. Effects of *Bacillus thuringiensis* and defoliation by the gypsy moth on native arthropods. Can. Entomol. (Accepted).

Stephenson, S.L. 1992. Department of Biology, Fairmont State College, Fairmont, West Virginia. Personal Communication.

USDA Forest Service. 1990. Gypsy moth news. 24.

Wood, P.S. and L. Butler. 1989. Biology of *Polia latex* (Guenee) (Noctuidae) with descriptions of eggs and larvae. J. Lepidop. Soc. 43:299-304.

TABLE 1.
Distribution of species of macrolepidopterous larvae by family; number of
species of larvae collected from tree foliage and from tree bands
at the Fernow Experimental Forest, WV 1989–1994.

Family	Foliage	Bands
Papilionidae	1	1
Lycaenidae	2	1
Nymphalidae	2	0
Thyatiridae	1	1
Drepanidae	1	0
Geometridae	43	24
Lasiocampidae	2	2
Saturniidae	3	0
Sphingidae	1	2
Notodontidae	12	6
Arctiidae	2	5
Lymantriidae	5	5
Noctuidae	36	39
TOTAL	111	86

TABLE 2
Macrolepidopterous larvae under bands of four host tree groups
at the Fernow Experimental Forest, West Virginia. Species are arranged from
most to least abundant for the combined sampling years of 1989–1994.
First and last collection dates are given.

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Lymantria dispar</i> (L.)	LYM	734	550	878	2267	4429	06MAY	13AUG
<i>Polia latex</i> (Gn.)	NOC	138	402	100	98	738	10JUN	15AUG
<i>Orthosia rubescens</i> (Wlk.)	NOC	107	197	46	135	485	20MAY	04JUL
<i>Abagrotis alternata</i> (Grt.)	NOC	39	40	32	108	219	06MAY	27JUN
<i>Lithophane hemina</i> Grt.	NOC	64	48	66	28	206	20MAY	23JUL
<i>Malacosoma dissτρια</i> Hbn.	LAS	10	60	24	38	132	09MAY	22JUN
<i>Lithophane innominata</i> (Sm.)	NOC	21	24	59	9	113	06JUN	11JUL
<i>Zale minerea</i> (Gn.)	NOC	7	68	20	8	103	11MAY	13AUG
<i>Malacosoma americanum</i> (F.)	LAS	24	46	17	10	97	16MAY	04JUL
<i>Erannis tiliaria</i> (Harr.)	GEO	11	15	17	27	70	13MAY	29JUN
<i>Catocala ultronia</i> (Hbn.)	NOC	9	50	7	2	68	17MAY	12JUL
<i>Melanolophia canadaria</i> (Gn.)	GEO	13	9	11	25	58	06MAY	10AUG
<i>Hyphantria cunea</i> (Drury)	ARC	11	24	6	9	50	24JUN	12AUG
<i>Epiglaea decliva</i> (Grt.)	NOC	8	10	14	11	43	06MAY	07JUN
<i>Orthosia hibisci</i> (Gn.)	NOC	15	9	6	11	41	18MAY	13JUL
<i>Acronicta hasta</i> Gn.	NOC	2	33	2	0	37	28JUN	15AUG

continued

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Alsophila pometaria</i> (Harr.)	GEO	4	9	5	18	36	06MAY	08JUN
<i>Dasychira obliquata</i> (G. & R.)	LYM	4	10	11	10	35	06MAY	05AUG
<i>Itame pustularia</i> (Gn.)	GEO	2	4	11	17	34	09MAY	21JUN
<i>Phigalia titea</i> (Cram.)	GEO	8	6	10	7	31	13MAY	18JUN
<i>Phoberia atomaris</i> Hbn.	NOC	0	6	1	22	29	20MAY	12JUL
<i>Catocala ilia</i> (Cram.)	NOC	1	2	0	24	27	06MAY	09AUG
<i>Hypoprepia fucosa</i> Hbn.	ARC	5	2	6	8	21	09MAY	01AUG
<i>Halysidota tessellaris</i> (J.E. Smith)	ARC	4	2	4	10	20	17JUN	13AUG
<i>Amphipyra pyramidoides</i> Gn.	NOC	5	4	5	6	20	21MAY	15AUG
<i>Catocala amica</i> (Hbn.)	NOC	1	7	5	5	18	03JUN	19JUL
<i>Crocigrapha normani</i> (Grt.)	NOC	4	11	1	0	16	20JUN	19JUL
<i>Dasychira dorsipennata</i> (B. & McD.)	LYM	2	3	4	3	12	28MAY	23JUL
<i>Lomographa vestaliata</i> (Gn.)	GEO	5	3	0	2	10	25MAY	10AUG
<i>Lomographa glomeraria</i> (Grt.)	GEO	0	4	2	4	10	20MAY	27JUN
<i>Lithophane laticinerea</i> Grt.	NOC	5	1	1	2	9	24MAY	21JUN
<i>Dasychira basiflava</i> (Pack.)	LYM	2	3	2	1	8	06MAY	12AUG
<i>Satyrrium calanus</i> (Hbn.)	LYC	0	1	2	4	7	28MAY	17JUN
<i>Papilio glaucus</i> L.	PAP	1	1	1	2	5	10JUN	12JUL
<i>Hydrelia inornata</i> (Hulst)	GEO	1	2	2	0	5	08JUL	15JUL
<i>Orygia leucostigma</i> (J.E. Smith)	LYM	0	0	3	2	5	10JUN	01JUL
<i>Bomolocha baltimoralis</i> (Gn.)	NOC	0	0	4	1	5	30MAY	09AUG
<i>Lithophane grotei</i> Riley	NOC	3	0	1	1	5	30MAY	12JUL
<i>Epimecis hortaria</i> (F.)	GEO	1	2	1	0	4	06MAY	12JUL
<i>Hypagyrtis unipunctata</i> (Haw.)	GEO	1	2	0	1	4	23MAY	13AUG
<i>Zale lunifera</i> (Hbn.)	NOC	0	4	0	0	4	20MAY	16JUL
<i>Achatia distincta</i> Hbn.	NOC	1	2	0	1	4	24JUN	13JUL
<i>Morrisonia confusa</i> (Hbn.)	NOC	2	2	0	0	4	13MAY	15AUG
<i>Lambdina fiscellaria</i> (Gn.)	GEO	0	1	0	2	3	01JUL	15JUL
<i>Paonias myops</i> (J.E. Smith)	SPH	1	1	0	1	3	01JUL	15JUL
<i>Lithophane antennata</i> (Wlk.)	NOC	1	1	1	0	3	23MAY	20JUN
<i>Pseudothyatira</i> <i>cymatophoroides</i> (Gn.)	THY	2	0	0	0	2	08JUL	05AUG
<i>Phigalia strigataria</i> (Minot)	GEO	0	1	0	1	2	20MAY	25MAY
<i>Lomographa semiclarata</i> (Wlk.)	GEO	0	1	0	1	2	25JUL	01AUG
<i>Lambdina pellucidaria</i> (G. & R.)	GEO	0	0	0	2	2	16JUL	22JUL
<i>Tetracis cachexiata</i> Gn.	GEO	0	1	0	1	2	03JUN	25JUL
<i>Eutrapela clemataria</i> (J.E. Smith)	GEO	1	0	0	1	2	28MAY	08JUL

continued

¹LYM = Lymantriidae, NOC = Noctuidae, LAS = Lasiocampidae, GEO = Geometridae, ARC = Arctiidae, LYC = Lycaenidae, PAP = Papilionidae, SPH = Sphingidae, THY = Thyatiridae, NOT = Notodontidae.

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Schizura unicornis</i> (J.E. Smith)	NOT	0	2	0	0	2	18JUN	12AUG
<i>Parallelia bistriaris</i> Hbn.	NOC	0	0	2	0	2	01JUL	30JUL
<i>Acronicta fragilis</i> Gn.	NOC	0	0	1	1	2	15JUL	30JUL
<i>Alypia octomaculata</i> (F.)	NOC	1	1	0	0	2	01JUL	01AUG
<i>Phlogophora periculosa</i> Gn.	NOC	1	1	0	0	2	17MAY	31MAY
<i>Lithophane unimoda</i> (Lint.)	NOC	0	2	0	0	2	31MAY	31MAY
<i>Eupsilia vinulenta</i> (Grt.)	NOC	1	0	0	1	2	24MAY	24MAY
<i>Polia nimbose</i> (Gn.)	NOC	0	1	1	0	2	06MAY	14MAY
<i>Himella intractata</i> (Morr.)	NOC	0	1	0	1	2	20JUN	20JUN
<i>Protoarmia porcelaria</i> (Gn.)	GEO	0	0	0	1	1	26JUL	26JUL
<i>Selenia kentaria</i> (G. & R.)	GEO	1	0	0	0	1	27JUN	27JUN
<i>Probole amicarica</i> (H.-S.)	GEO	0	0	0	1	1	05AUG	05AUG
<i>Eulithis diversilineata</i> (Hbn.)	GEO	0	1	0	0	1	01JUL	01JUL
<i>Hydria prunivorata</i> (Fgn.)	GEO	0	1	0	0	1	25MAY	25MAY
<i>Hydrelia condensata</i> (Wlk.)	GEO	0	0	1	0	1	08AUG	08AUG
<i>Eupithecia herefordaria</i> C. & S.	GEO	0	0	1	0	1	07JUN	07JUN
<i>Cladara atrolitratata</i> (Wlk.)	GEO	0	0	1	0	1	06JUN	06JUN
<i>Darapsa myron</i> (Cram.)	SPH	0	1	0	0	1	04JUL	04JUL
<i>Nadata gibbosa</i> (J.E. Smith)	NOT	0	0	0	1	1	02AUG	02AUG
<i>Symmerista leucitys</i> Franc.	NOT	0	0	1	0	1	08AUG	08AUG
<i>Macrurocampa marthesia</i> (Cram.)	NOT	0	0	0	1	1	15JUL	15JUL
<i>Heterocampa guttivitta</i> (Wlk.)	NOT	0	0	0	1	1	29JUL	29JUL
<i>Lochmaeus manteo</i> Doubleday	NOT	0	0	1	0	1	15JUL	15JUL
<i>Ecpanteria scribonia</i> (Stoll)	ARC	0	0	0	1	1	30MAY	30MAY
<i>Lophocampa caryae</i> Harr.	ARC	0	0	1	0	1	05AUG	05AUG
<i>Idia julia</i> (B. & McD.)	NOC	0	0	1	0	1	09JUL	09JUL
<i>Idia scobialis</i> (Grt.)	NOC	0	0	1	0	1	10JUN	10JUN
<i>Zale lunata</i> (Drury)	NOC	0	1	0	0	1	09JUL	09JUL
<i>Acronicta americana</i> (Harr.)	NOC	0	0	1	0	1	09AUG	09AUG
<i>Acronicta innotata</i> Gn.	NOC	1	0	0	0	1	12AUG	12AUG
<i>Acronicta lobeliae</i> Gn.	NOC	0	0	0	1	1	17JUN	17JUN
<i>Eudryas grata</i> (F.)	NOC	1	0	0	0	1	12AUG	12AUG
<i>Eutolype rolandi</i> Grt.	NOC	1	0	0	0	1	21JUN	21JUN
<i>Psaphida resumens</i> Wlk.	NOC	0	1	0	0	1	01JUN	01JUN

¹LYM = Lymantriidae, NOC = Noctuidae, LAS = Lasiocampidae, GEO = Geometridae, ARC = Arctiidae, LYC = Lycaenidae, PAP = Papilionidae, SPH = Sphingidae, THY = Thyatiridae, NOT = Notodontidae.

TABLE 3. Macrolepidopterous larvae collected from foliage of four host tree groups at the Fernow Experimental Forest, West Virginia.

Species are arranged from most to least abundant for the combined sampling years of 1989-1994. First and last collection dates are given.

Species ^{2,3}	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Lomographa glomeraria</i> (Grt.)	GEO	13	795	4	3	815	10MAY	28JUN
<i>Erannis tiliaria</i> (Harr.)	GEO	275	143	159	214	791	07MAY	19JUN
<i>Lomographa vestaliata</i> (Gn.)	GEO	7	664	7	7	685	30MAY	15AUG
<i>Melanolopia canadaria</i> (Gn.)	GEO	106	323	134	77	640	07MAY	15AUG
<i>Alsophila pometaria</i> (Harr.)	GEO	44	202	62	219	527	07MAY	23JUN
<i>Orthosia hibisci</i> (Gn.)	NOC	217	108	23	107	455	07MAY	28JUN
<i>Polia latex</i> (Gn.)	NOC	77	107	101	51	336	07JUN	15AUG
<i>Itame pustularia</i> (Gn.)	GEO	0	5	287	2	294	07JUN	15AUG
<i>Lymantria dispar</i> (L.)	LYM	28	12	48	201	289	07MAY	29JUN
<i>Hydria prunivorata</i> (Fgn.)	GEO	0	193	0	0	193	18JUN	13AUG
<i>Morrisonia confusa</i> (Hbn.)	NOC	46	77	21	17	161	29MAY	15AUG
<i>Hydrelia inornata</i> (Hulst) ³	GEO	45	21	79	11	156	10MAY	15AUG
<i>Phigalia titea</i> (Cram.)	GEO	30	33	22	68	153	07MAY	13JUN
<i>Campaea perlata</i> (Gn.)	GEO	36	48	44	11	139	07MAY	15AUG
<i>Orthosia rubescens</i> (Wlk.)	NOC	54	25	24	19	122	10MAY	07JUL
<i>Eupsilia sidus</i> (Gn.) ²	NOC	99	1	2	12	114	07MAY	07JUN
<i>Amphipyra pyramidoides</i> Gn.	NOC	10	46	23	24	103	07MAY	15JUN
<i>Lomographa semiclarata</i> (Wlk.)	GEO	0	72	1	1	74	04JUN	20JUL
<i>Hyphantria cunea</i> (Drury)	ARC	21	8	9	35	73	22MAY	13AUG
<i>Eupsilia n. sp.</i> ²	NOC	20	15	7	24	66	10MAY	10JUN
<i>Symmerista leucitys</i> Franc.	NOT	0	0	49	0	49	26JUL	09AUG
<i>Halysidota tessellaris</i> (J.E. Smith)	ARC	7	10	7	20	44	02JUL	15AUG
<i>Tetracis cachexiata</i> Gn.	GEO	11	21	7	4	43	07MAY	15AUG
<i>Lithophane hemina</i> Grt.	NOC	11	17	13	1	42	23MAY	06JUL
<i>Hydrelia condensata</i> (Wlk.) ²	GEO	14	1	22	3	40	13JUL	10AUG
<i>Lochmaeus manteo</i> Doubleday	NOT	2	0	0	37	39	29JUN	14AUG
<i>Peridea basitriens</i> (Wlk.)	NOT	1	0	38	0	39	05JUL	14AUG
<i>Besma quercivoraria</i> (Gn.)	GEO	4	0	0	34	38	11MAY	03AUG
<i>Phigalia strigataria</i> (Minot) ²	GEO	18	5	2	10	35	07MAY	12JUN
<i>Iridopsis larvaria</i> (Gn.)	GEO	1	33	0	0	34	04JUN	03AUG
<i>Besma endropiaria</i> (G. & R.)	GEO	0	3	26	5	34	01JUN	10AUG
<i>Malacosoma distria</i> Hbn. ²	LAS	0	4	2	27	33	12MAY	12JUN
<i>Dryocampa rubicunda</i> (F.)	SAT	0	0	31	0	31	09JUL	16JUL

continued

¹GEO = Geometridae, NOC = Noctuidae, ARC = Arctiidae, LAS = Lasiocampidae, NOT = Notodontidae, LYM = Lymantriidae, SAT = Saturniidae, DRE = Drepanidae, SPH = Sphingidae, THY = Thyatiridae, PAP = Papilionidae, NYM = Nymphalidae, LYC = Lycaenidae.

²Species that were not collected at Coopers Rock, 1984-1985 (Butler 1992).

³Species collected on oak at Fernow that were not collected in Grant and Pendleton counties, 1990-1992 (Butler et al. 1995a). Oak is an expected host for these species.

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Anavitrinella pampinaria</i> (Gn.)	GEO	9	21	0	0	30	14MAY	18JUN
<i>Acronicta fragilis</i> Gn.	NOC	19	7	2	0	28	04JUN	14AUG
<i>Eupithecia miserulata</i> Grt.	GEO	0	19	6	1	26	29MAY	25JUL
<i>Eupithecia herefordaria</i> C. & S. ²	GEO	5	13	6	0	24	07JUN	05JUL
<i>Satyrium calanus</i> (Hbn.)	LYC	0	0	0	23	23	07MAY	25MAY
<i>Pseudothyatira cymatophoroides</i> (Gn.)	THY	20	2	0	1	23	02JUL	10AUG
<i>Hypagyrtis unipunctata</i> (Haw.)	GEO	5	10	7	1	23	15MAY	07AUG
<i>Heterocampa guttivitta</i> (Wlk.)	NOT	10	0	9	2	21	07JUN	25JUL
<i>Probole amicaria</i> (H.-S.)	GEO	0	1	17	1	19	19JUN	13AUG
<i>Zale minerea</i> (Gn.)	NOC	5	9	3	2	19	21MAY	01AUG
<i>Drepana arcuata</i> Wlk.	DRE	17	0	0	0	17	11JUN	07AUG
<i>Nadata gibbosa</i> (J.E. Smith)	NOT	0	0	0	17	17	18JUN	14AUG
<i>Achatia distincta</i> Hbn.	NOC	6	6	0	3	15	29MAY	27JUN
<i>Orgyia leucostigma</i> (J.E. Smith)	LYM	1	8	3	3	15	17MAY	16JUL
<i>Lithophane antennata</i> (Wlk.) ²	NOC	11	3	0	1	15	17MAY	07JUN
<i>Ennomos subsignaria</i> (Hbn.) ²	GEO	0	0	6	8	14	21MAY	26JUN
<i>Acronicta ovata</i> Grt.	NOC	2	0	0	11	13	12JUL	11AUG
<i>Paonias myops</i> (J.E. Smith) ³	SPH	1	7	0	4	12	06JUL	10AUG
<i>Acronicta hasta</i> Gn.	NOC	3	5	2	0	10	06JUL	14AUG
<i>Orthosia alurina</i> (Sm.) ²	NOC	2	3	5	0	10	14MAY	20JUN
<i>Crocigrapha normani</i> (Grt.)	NOC	3	6	1	0	10	24MAY	27JUN
<i>Bomolocha baltimoralis</i> (Gn.)	NOC	3	4	3	0	10	11JUL	02AUG
<i>Parallelia bistriaris</i> Hbn.	NOC	1	0	9	0	10	15MAY	10AUG
<i>Anisota virginiensis</i> (Drury)	SAT	0	0	0	9	9	09JUL	09JUL
<i>Acronicta americana</i> (Harr.) ³	NOC	3	3	2	1	9	09JUL	15AUG
<i>Catocala ultronia</i> (Hbn.) ²	NOC	0	8	0	0	8	25MAY	01JUN
<i>Biston betularia</i> (L.)	GEO	3	4	0	0	7	02JUL	24JUL
<i>Eutrapela clemataria</i> (J.E. Smith)	GEO	0	3	3	1	7	29MAY	23JUN
<i>Malacosoma americanum</i> (F.)	LAS	0	6	0	1	7	24MAY	13JUN
<i>Symmerista albifrons</i> (J.E. Smith) ²	NOT	0	0	0	7	7	26JUL	02AUG
<i>Cleora sublunaria</i> (Gn.) ²	GEO	0	4	2	1	7	26MAY	05JUN
<i>Cladara atroliturlata</i> (Wlk.) ³	GEO	3	1	2	1	7	07JUN	23JUN
<i>Lithophane laticinerea</i> Grt. ²	NOC	0	4	0	3	7	17MAY	22JUN
<i>Papilio glaucus</i> L. ³	PAP	0	6	0	0	6	28JUN	24JUL
<i>Colocasia propinquilinea</i> (Grt.)	NOC	4	0	2	0	6	25JUN	21JUL
<i>Lithophane innominata</i> (Sm.) ^{2,3}	NOC	2	0	2	2	6	07JUN	14JUN
<i>Lithophane querquera</i> Grt. ²	NOC	2	0	0	4	6	14MAY	05JUN
<i>Eupsilia morrisoni</i> (Grt.)	NOC	1	0	0	5	6	07MAY	22MAY

continued

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Protoboarmia porcelaria</i> (Gn.)	GEO	4	1	0	0	5	07MAY	13JUL
<i>Ennomos magnaria</i> Gn.	GEO	0	2	3	0	5	29JUN	13JUL
<i>Furcula borealis</i> (Guer.-Meneville)	NOT	0	5	0	0	5	30JUN	20JUL
<i>Nacophora quernaria</i> (J.E. Smith)	GEO	0	0	0	4	4	02JUL	13JUL
<i>Glena cribrataria</i> (Gn.)	GEO	0	3	0	0	3	30JUL	30JUL
<i>Anacamptodes humaria</i> (Gn.) ²	GEO	0	3	0	0	3	07JUN	20JUN
<i>Paleacrita vernata</i> (Peck) ^{2,3}	GEO	0	0	0	3	3	11MAY	11MAY
<i>Plagodis alcoolaria</i> (Gn.)	GEO	0	0	2	1	3	05JUL	09AUG
<i>Prochoerodes transversata</i> (Drury) ²	GEO	0	2	1	0	3	08JUN	14AUG
<i>Macrurocampa marthesia</i> (Cram.)	NOT	1	0	0	2	3	23JUL	14AUG
<i>Dasychira obliquata</i> (G. & R.)	LYM	0	0	0	3	3	14MAY	23JUL
<i>Zale lunifera</i> (Hbn.)	NOC	0	3	0	0	3	19MAY	19MAY
<i>Acronicta innotata</i> Gn. ²	NOC	3	0	0	0	3	07AUG	14AUG
<i>Lithophane unimoda</i> (Lint.) ²	NOC	0	3	0	0	3	15MAY	25MAY
<i>Epimecis hortaria</i> (F.) ²	GEO	1	0	0	1	2	02JUN	12JUL
<i>Melanolophia signataria</i> (Wlk.) ²	GEO	2	0	0	0	2	08JUN	08JUN
<i>Plagodis serinaria</i> H.-S.	GEO	0	1	0	1	2	09JUL	20JUL
<i>Heterocampa biundata</i> Wlk.	NOT	1	0	1	0	2	27JUN	07AUG
<i>Dasychira dorsipennata</i> (B. & McD.)	LYM	0	0	0	2	2	30MAY	30MAY
<i>Bomolocha deceptalis</i> (Wlk.) ²	NOC	0	2	0	0	2	18JUL	18JUL
<i>Catocala andromedae</i> Gn. ²	NOC	0	0	0	2	2	24MAY	24MAY
<i>Acronicta radcliffei</i> (Harv.) ²	NOC	0	2	0	0	2	19JUN	19JUN
<i>Alypia octomaculata</i> (F.) ²	NOC	0	2	0	0	2	01AUG	01AUG
<i>Lithophane semiusta</i> Grt. ²	NOC	0	0	2	0	2	08JUN	08JUN
<i>Copipanolis styracis</i> (Gn.) ²	NOC	0	0	0	2	2	15MAY	24MAY
<i>Satyrium liparops</i> (Leconte) ^{2,3}	LYC	0	0	0	1	1	24MAY	24MAY
<i>Basilarchia arthemis</i> (Drury) ²	NYM	0	1	0	0	1	12JUL	12JUL
<i>Basilarchia archippus</i> (Cram.) ²	NYM	0	0	0	1	1	30MAY	30MAY
<i>Pero honestaria</i> (Wlk.) ²	GEO	0	0	1	0	1	27JUN	27JUN
<i>Lambdina fiscellaria</i> (Gn.) ²	GEO	0	0	0	1	1	18JUN	18JUN
<i>Lambdina pellucidaria</i> (G. & R.) ^{2,3}	GEO	0	0	0	1	1	14JUL	14JUL
<i>Eugonobapta nivosaria</i> (Gn.) ^{2,3}	GEO	0	0	0	1	1	21MAY	21MAY
<i>Nemoria lixaria</i> (Gn.) ²	GEO	0	1	0	0	1	13JUL	13JUL

continued

¹GEO = Geometridae, NOC = Noctuidae, ARC = Arctiidae, LAS = Lasiocampidae, NOT = Notodontidae, LYM = Lymantriidae, SAT = Saturniidae, DRE = Drepanidae, SPH = Sphingidae, THY = Thyatiridae, PAP = Papilionidae, NYM = Nymphalidae, LYC = Lycaenidae.

²Species that were not collected at Coopers Rock, 1984-1985 (Butler 1992).

³Species collected on oak at Fernow that were not collected in Grant and Pendleton counties, 1990-1992 (Butler et al. 1995a). Oak is an expected host for these species.

Species	Family ¹	No. per host tree					Collection Dates	
		Birch	Cherry	Maple	Oak	Total	First	Last
<i>Actias luna</i> (L.) ³	SAT	0	0	0	1	1	09AUG	09AUG
<i>Peridea angulosa</i> (J.E. Smith)	NOT	0	0	0	1	1	31JUL	31JUL
<i>Schizura unicornis</i> (J.E. Smith)	NOT	0	0	1	0	1	14AUG	14AUG
<i>Oligocentria semirufescens</i> (Wlk.)	NOT	0	1	0	0	1	09AUG	09AUG
<i>Orgyia definita</i> Pack. ²	LYM	0	1	0	0	1	28JUN	28JUN
<i>Acronicta lobeliae</i> Gn. ²	NOC	0	0	0	1	1	28JUN	28JUN
<i>Acronicta haesitata</i> (Grt.) ²	NOC	0	1	0	0	1	17JUL	17JUL
<i>Lithophane grotei</i> Riley	NOC	0	0	0	1	1	24MAY	24MAY

¹GEO = Geometridae, NOC = Noctuidae, ARC = Arctiidae, LAS = Lasiocampidae, NOT = Notodontidae, LYM = Lymantriidae, SAT = Saturniidae, DRE = Drepanidae, SPH = Sphingidae, THY = Thyatiridae, PAP = Papilionidae, NYM = Nymphalidae, LYC = Lycaenidae.

²Species that were not collected at Coopers Rock, 1984-1985 (Butler 1992).

³Species collected on oak at Fernow that were not collected in Grant and Pendleton counties, 1990-1992 (Butler et al. 1995a). Oak is an expected host for these species.

TABLE 4.
Species richness and abundance of macrolepidopterous larvae
on foliage of four host tree groups at Fernow Experimental Forest,
West Virginia, 1989-1994.

Host	Richness	Abundance
black birch	57	1,350
black cherry	71	3,188
mixed maple	57	1,352
mixed oak	72	1,387
TOTAL	111	7,277

Blank Page in Original Bulletin

Agricultural and Forestry
Experiment Station
West Virginia University
PO Box 6108
Morgantown, WV 26506-6108

Non-Profit Organization
U.S. Postage
PAID
Morgantown, WV
Permit No. 34

SERIALS DEPARTMENT
EVANSDALE LIBRARY
WEST VIRGINIA UNIVERSITY
PO BOX 6105
MORGANTOWN WV 26506-6105