

2009

Cryptic Subtelomeric Rearrangements and X Chromosome Mosaicism: A Study of 565 Apparently Normal Individuals with Fluorescent In Situ Hybridization

Jasen L. Wise
West Virginia University

Richard J. Crout
West Virginia University

Daniel W. McNeil
West Virginia University

Robert J. Weyant
University of Pittsburgh

Mary L. Marazita
University of Pittsburgh

See next page for additional authors

Follow this and additional works at: https://researchrepository.wvu.edu/faculty_publications

Digital Commons Citation

Wise, Jasen L.; Crout, Richard J.; McNeil, Daniel W.; Weyant, Robert J.; Marazita, Mary L.; and Wenger, Sharon L., "Cryptic Subtelomeric Rearrangements and X Chromosome Mosaicism: A Study of 565 Apparently Normal Individuals with Fluorescent In Situ Hybridization" (2009). *Faculty & Staff Scholarship*. 2865.

https://researchrepository.wvu.edu/faculty_publications/2865

This Article is brought to you for free and open access by The Research Repository @ WVU. It has been accepted for inclusion in Faculty & Staff Scholarship by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

Authors

Jasen L. Wise, Richard J. Crout, Daniel W. McNeil, Robert J. Weyant, Mary L. Marazita, and Sharon L. Wenger

Cryptic Subtelomeric Rearrangements and X Chromosome Mosaicism: A Study of 565 Apparently Normal Individuals with Fluorescent *In Situ* Hybridization

Jasen L. Wise^{1*}, Richard J. Crout², Daniel W. McNeil², Robert J. Weyant³, Mary L. Marazita⁴, Sharon L. Wenger¹

1 Department of Pathology, West Virginia University, Morgantown, West Virginia, United States of America, **2** School of Dentistry, West Virginia University, Morgantown, West Virginia, United States of America, **3** Department of Dental Public Health and Information Management, University of Pittsburgh School of Dental Medicine, Pittsburgh, Pennsylvania, United States of America, **4** Center for Craniofacial and Dental Genetics, Departments of Oral Biology and Human Genetics, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

Abstract

Five percent of patients with unexplained mental retardation have been attributed to cryptic unbalanced subtelomeric rearrangements. Half of these affected individuals have inherited the rearrangement from a parent who is a carrier for a balanced translocation. However, the frequency of carriers for cryptic balanced translocations is unknown. To determine this frequency, 565 phenotypically normal unrelated individuals were examined for balanced subtelomeric rearrangements using Fluorescent *In Situ* hybridization (FISH) probes for all subtelomere regions. While no balanced subtelomeric rearrangements were identified, three females in this study were determined to be mosaic for the X chromosome. Mosaicism for XXX cell lines were observed in the lymphocyte cultures of 3 in 379 women (0.8%), which is a higher frequency than the 1 in 1000 (0.1%) reported for sex chromosome aneuploidies. Our findings suggest that numerical abnormalities of the X chromosome are more common in females than previously reported. Based on a review of the literature, the incidence of cryptic translocation carriers is estimated to be approximately 1/8,000, more than ten-fold higher than the frequency of visible reciprocal translocations.

Citation: Wise JL, Crout RJ, McNeil DW, Weyant RJ, Marazita ML, et al. (2009) Cryptic Subtelomeric Rearrangements and X Chromosome Mosaicism: A Study of 565 Apparently Normal Individuals with Fluorescent *In Situ* Hybridization. PLoS ONE 4(6): e5855. doi:10.1371/journal.pone.0005855

Editor: Syed A. Aziz, Health Canada, Canada

Received: April 7, 2009; **Accepted:** May 12, 2009; **Published:** June 10, 2009

Copyright: © 2009 Wise et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This project was funded in part by NIH grant #5R01-DE014899. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jasenwise@gmail.com

Introduction

Subtelomeres are the most distal sequences of non-repetitive DNA on the chromosome, and have the highest density of genes in the genome [1]. Any rearrangement or deletion in these gene-rich regions could have severe phenotypic consequences. Numerous studies have shown that 0.5 to 10.7% (variation due to study selection criteria) of patients with unexplained mental retardation (MR) have an unbalanced cryptic subtelomeric rearrangement or deletion [2–11].

Cytogenetically visible balanced translocations have an incidence of 1 in 600 in the general population [12]. Parents, who are carriers of balanced translocations, are at risk for having children with unbalanced gene complements. Fifty percent of the gametes produced by a balanced translocation carrier will have segmental aneuploidy, which can result in a child with an unbalanced rearrangement. The most likely way to identify a balanced carrier is through a child who has been identified with an unbalanced rearrangement.

Several studies have reported that half of all patients with an unbalanced cryptic rearrangement have inherited it from a parent with a cryptic balanced translocation [4,10]. The frequency of

individuals who carry a balanced cryptic translocation is unknown. In this study, 565 unrelated, phenotypically normal individuals were screened with subtelomere FISH probes to determine if balanced cryptic translocation carriers could be identified.

Methods

Subjects for this study were drawn from the Center for Oral Health Research in Appalachia (COHRA) [13], an ongoing cross-sectional oral health etiology study. COHRA ascertains families from two central West Virginia counties and two western Pennsylvania counties and performs a detailed assessment protocol after an informed consent process approved by the Institutional Review Boards (IRB) of the University of Pittsburgh and West Virginia University (WVU). A total of 484 COHRA subjects were included in the current study: 164 male and 320 female. In addition, IRB approval was obtained to use discarded samples from the WVU cytogenetics laboratory. These 81 (22 males and 59 females) samples were selected on the basis of a normal karyotype at least the 550+ band level, which were then deidentified prior to analysis.

Peripheral blood lymphocytes were processed using standard clinical cytogenetic techniques. Cells were dropped onto slides, which were then immersed through the following series of washes in coplin jars: $2\times$ SSC for 10 minutes at 37°C , 1% formaldehyde for 15 minutes at room temperature (RT), $1\times$ PBS for 5 minutes at RT, pepsin solution for 13 minutes at 37°C , $1\times$ PBS for 5 minutes at RT and then air dried. The slides were then placed through a series of ethanol washes of 70%, 85%, and 100% for 1 minute each and allowed to air dry at room temperature.

Working probe solutions were prepared by adding $3\ \mu\text{l}$ of ToTelVysion probe solution (Abbott Molecular Inc, cat# 33-270000) to $30\ \mu\text{l}$ of cDenHyb (InSitus, cat #D002) in a microfuge tube and mixed well. Three μl of each working probe solution was placed in the middle of one of 5 respective circled areas on a slide. A 12 mm circular coverslip was added and all air bubbles driven out. When 5 spots per slide were completed, lab tape was placed across the entire slide and pressed firmly for a tight seal. Slides were placed on a hotplate for 3 minutes at 90°C , then in a light-tight box, and incubated overnight in a 37°C water bath.

The next day, in a minimal light room, the slides were removed from the water bath and de-coverslipped. The slides were then washed in $0.4\times$ SSC/0.3% NP-40 at 73°C for two minutes followed by 30 seconds in $2\times$ SSC/0.1% NP-40 at room temperature. The slides were then completely air dried in the dark. Twenty μl of $1\times$ DAPI counterstain was applied to the slides and coverslipped.

A Leica epi-fluorescent microscope equipped with a DAPI single bandpass, aqua single bandpass, and a red/green dual bandpass filter was used for signal enumeration. Yellow signals were read using the red/green filter. Five metaphase and 5 interphase cells were scored for each of 15 subtelomeric probe sets per subject.

To confirm abnormal subtelomere FISH results regarding X chromosomes, slides were prepared as described previously and hybridized with X/Y centromere probes. Two scorers analyzed 100 cells each for percentage of abnormal cells.

Results

A total of 565 samples were evaluated for cryptic rearrangements using subtelomeric FISH probes. No balanced cryptic rearrangements were observed by FISH, all samples showing normal number and location of signals (Fig. 1).

Among the specimens that were analyzed with subtelomeric FISH probe sets containing Xp/Yp and Xq/Yq, mosaicism for X chromosome aneuploidy was identified in 3 of 379 women (0.8%). The results were confirmed using a separate X/Y centromeric probe set. The FISH results in interphase cells for these individuals identified 89% triple X in a 36 year old, 5% triple X chromosome in a 52 year old, and multiple cell lines including 11% single X, 6% XXX and 2% XXXX in a 54 year old (Fig. 2).

Discussion

Subtelomeric Rearrangements

No balanced cryptic translocations were found among the 565 subjects who were screened by subtelomere FISH. From the small sample size, it was not possible to determine the frequency of balanced subtelomeric translocation carriers. Our sample size was limited due to the number of individuals enrolled in the study who donated blood, loss due to culture failures, and the cost of FISH probes.

Since the frequency for a balanced cryptic rearrangement could not be estimated from our study sample, we estimated the




Figure 1. An example of a FISH hybridized metaphase spread. Four different probe signals are visible: 2p (green signal), 2q (red signal), Xq/Yq (yellow signal) and X (aqua signal). This pattern represents the pattern seen in a normal diploid cell from a female.
doi:10.1371/journal.pone.0005855.g001




Figure 2. Interphase cells hybridized with a FISH probe for the centromere of the X chromosome. A) A cell with one signal for the centromere of the X chromosome or monosomy X. B) Two normal cells showing two signals for the X chromosome, and one cell with monosomy X. C) A cell showing trisomy X. D) A cell with tetrasomy X.
doi:10.1371/journal.pone.0005855.g002

frequency based on reports in the literature. Carrier frequency (X) was estimated based on the equation: $(A)(B)(C)(D)(E) = X$ where A is the 2% of the population with MR [14], B is the 50% of MR patients with unknown etiology [15], C is the estimated 5% incidence of subtelomeric abnormalities in MR patients with unknown etiology [16], D is the 50% percent risk for inheriting the unbalanced rearrangement from a parent with a balanced subtelomeric translocation [4], and E is the 50% chance that a carrier would have a child with an unbalanced rearrangement (due to chromosome segregation in the gametes).

Using this equation, the incidence of cryptic *balanced* subtelomeric rearrangement carriers in the general population is approximately 1 in 8000. Therefore, it is not surprising that the current series of 565 subjects found no one with a cryptic balanced translocation.

The reported incidence of individuals with *unbalanced* subtelomeric rearrangements in the general population has been estimated by Knight and colleagues [4] to be 2.1 in 10,000 (~1 in 4762). Because half of these individuals inherited the rearrangement from a parent [4,10], the frequency of parents who are *balanced* translocation carriers would be half as frequent, or around 1 in 9524. The difference between this calculation and ours is most likely due to the variation in the reported percentages

and criteria for evaluation of MR. Regardless, the incidence of balanced subtelomeric rearrangements is at least 13 times more prevalent than carriers of visible reciprocal translocations.

X Chromosome Mosaicism

Individuals who are mosaic have two or more populations of somatic cells that are genetically different. Aneuploidy of the X chromosome can arise by mitotic nondisjunction or anaphase lag. If this event occurs during early fetal development, higher percentages of mosaicism will occur. If the error occurs at a later time during fetal development or after birth, then lower percentages will be present. Individuals who have low level mosaicism for the X chromosome are less likely to have a clinically relevant phenotype and therefore would go undiagnosed in the general population. There are several factors to consider with low level mosaicism: are the results due to genuine mosaicism, technical artifact [17], or age related?

X chromosome aneuploidy has been attributed to premature centromere division in older women. Several studies have demonstrated that peripheral blood metaphase cells from women generally 50 years of age and older could have an average of 4–5% X chromosome loss and less than 1% gain attributed to mitotic error [18–21]. This finding has been substantiated in interphase

cells using FISH probes, demonstrating that women older than 60 years of age had an average X chromosome loss of 3.4% (as high as 9%) and gain of less than 1% [22–23].

FISH studies have shown that cultured lymphocytes reflect in vivo aneuploidy rates and that there is no significant difference between cultured lymphocyte and uncultured lymphocyte stability [22,24]. All 3 of the individuals identified in this study to have X chromosome mosaicism had gains of X chromosomes. Although two of the three women were over the ages of 50, all three women had a cell line with 5% or greater for an extra X chromosome, suggesting that the aneuploidy is not related to age or tissue culture artifact.

We were unable to find any literature that has estimated the incidence of sex chromosome mosaicism in the general population or the incidence of balanced cryptic subtelomeric rearrangements in the general population. We estimate that the incidence of balanced cryptic translocation to be at least 1 in 8,000. To our

knowledge, the finding of mosaicism in 0.8% of women may be the first reported incidence of low level sex chromosome mosaicism in the general population, much higher than the 0.1% reported in newborn studies for sex chromosome aneuploidy, which was based on analysis of 3–5 cells [25]. Our findings suggest that numerical abnormalities of the X chromosome may be more common in females than previously reported.

Acknowledgments

Many thanks to the COHRA study subjects for their participation.

Author Contributions

Conceived and designed the experiments: JLW SLW. Performed the experiments: JLW. Analyzed the data: JLW SLW. Contributed reagents/materials/analysis tools: RJC DWM RJW MLM SLW. Wrote the paper: JLW RJC DWM RJW MLM SLW.

References

- Helias-Rodzewicz Z, Bocian E, Stankiewicz P, Obersztyń E, Kostyk E, et al. (2002) Subtelomeric rearrangements detected by FISH in three of 33 families with idiopathic mental retardation and minor physical anomalies. *J Med Genet* 39: e53.
- Flint J, Wilkie AO, Buckle VJ, Winter RM, Holland AJ, et al. (1995) The detection of subtelomeric chromosomal rearrangements in idiopathic mental retardation. *Nat Genet* 9: 132–140.
- Knight SJ, Horsley SW, Regan R, Lawrie NM, Maher EJ, et al. (1997) Development and clinical application of an innovative fluorescence in situ hybridization technique which detects submicroscopic rearrangements involving telomeres. *Eur J Hum Genet* 5: 1–8.
- Knight SJ, Regan R, Nicod A, Horsley SW, Kearney L, et al. (1999) Subtle chromosomal rearrangements in children with unexplained mental retardation. *Lancet* 354: 1676–1681.
- Joyce CA, Dennis NR, Cooper S, Browne CE (2001) Subtelomeric rearrangements: results from a study of selected and unselected probands with idiopathic mental retardation and control individuals by using high-resolution G-banding and FISH. *Hum Genet* 109: 440–451.
- Anderlid BM, Schoumans J, Anneren G, Sahlen S, Kyllerman M, et al. (2002) Subtelomeric rearrangements detected in patients with idiopathic mental retardation. *Am J Med Genet* 107: 275–284.
- Van Karnebeek CD, Koevoets C, Sluiter S, Bijlsma EK, Smeets DF, et al. (2002) Prospective screening for subtelomeric rearrangements in children with mental retardation of unknown aetiology: the Amsterdam experience. *J Med Genet* 39: 546–553.
- Bocian E, Helias-Rodzewicz Z, Suchenek K, Obersztyń E, Kutkowska-Kazmierczak A, et al. (2004) Subtelomeric rearrangements: results from FISH studies in 84 families with idiopathic mental retardation. *Med Sci Monit* 10: CR143–151.
- Li R, Zhao ZY (2004) Two subtelomeric chromosomal deletions in forty-six children with idiopathic mental retardation. *Chin Med J (Engl)* 117: 1414–1417.
- Adeyinka A, Adams SA, Lorentz CP, Van Dyke DL, Jalal SM (2005) Subtelomere deletions and translocations are frequently familial. *Am J Med Genet A* 135: 28–35.
- Ravnan JB, Tepperberg JH, Papenhausen P, Lamb AN, Hedrick J, et al. (2006) Subtelomere FISH analysis of 11 688 cases: an evaluation of the frequency and pattern of subtelomere rearrangements in individuals with developmental disabilities. *J Med Genet* 43: 478–489.
- Estop AM, Ciepły KM, Aston CE (1997) The meiotic segregation pattern of a reciprocal translocation t(10;12)(q26.1;p13.3) by fluorescence in situ hybridization sperm analysis. *Eur J Hum Genet* 5: 78–82.
- Polk DE, Weyant RJ, Crout RJ, McNeil DW, Tarter RE, et al. (2008) Study protocol of the Center for Oral Health Research in Appalachia (COHRA) etiology study. *BMC Oral Health* 8: 18.
- Baker E, Hinton L, Callen DF, Altree M, Dobbie A, et al. (2002) Study of 250 children with idiopathic mental retardation reveals nine cryptic and diverse subtelomeric chromosome anomalies. *Am J Med Genet* 107: 285–293.
- Flint J, Wilkie AO (1996) The genetics of mental retardation. *Br Med Bull* 52: 453–464.
- Walter S, Sandig K, Hinkel GK, Mitulla B, Ounap K, et al. (2004) Subtelomere FISH in 50 children with mental retardation and minor anomalies, identified by a checklist, detects 10 rearrangements including a de novo balanced translocation of chromosomes 17p13.3 and 20q13.33. *Am J Med Genet A* 128: 364–373.
- Wenger SL, Golden WL, Dennis SP, Steele MW (1984) Are the occasional aneuploid cells in peripheral blood cultures significant? *Am J Med Genet* 19: 715–719.
- Ford JH, Russell JA (1985) Differences in the error mechanisms affecting sex and autosomal chromosomes in women of different ages within the reproductive age group. *Am J Hum Genet* 37: 973–983.
- Fitzgerald PH (1975) A mechanism of X chromosome aneuploidy in lymphocytes of aging women. *Humangenetik* 28: 153–158.
- Abruzzo MA, Mayer M, Jacobs PA (1985) Aging and aneuploidy: evidence for the preferential involvement of the inactive X chromosome. *Cytogenet Cell Genet* 39: 275–278.
- Russell LM, Strike P, Browne CE, Jacobs PA (2007) X chromosome loss and ageing. *Cytogenet Genome Res* 116: 181–185.
- Guttenbach M, Koschorz B, Bernthaler U, Grimm T, Schmid M (1995) Sex chromosome loss and aging: in situ hybridization studies on human interphase nuclei. *Am J Hum Genet* 57: 1143–1150.
- Mukherjee AB, Alejandro J, Payne S, Thomas S (1996) Age-related aneuploidy analysis of human blood cells in vivo by fluorescence in situ hybridization (FISH). *Mech Ageing Dev* 90: 145–156.
- Catalan J, Surrallés J, Falck GC, Autio K, Norppa H (2000) Segregation of sex chromosomes in human lymphocytes. *Mutagenesis* 15: 251–255.
- Hook E, Hamerton J (1977) The frequency of chromosome abnormalities detected in consecutive newborn studies - differences between studies - results by sex and by severity of phenotypic involvement. Hook E, Porter I, eds. Academic Press, Inc. pp 63–79.