2010

Alterations in T cell function and activation during exposure to the herbicide 3,4 dichloropropionanilide (DCPA) and its metabolites

Tricia L. Lewis
West Virginia University

Follow this and additional works at: <https://researchrepository.wvu.edu/etd>

Recommended Citation

Lewis, Tricia L., "Alterations in T cell function and activation during exposure to the herbicide 3,4 dichloropropionanilide (DCPA) and its metabolites" (2010). *Graduate Theses, Dissertations, and Problem Reports*. 3087.

<https://researchrepository.wvu.edu/etd/3087>

This Dissertation is protected by copyright and/or related rights. It has been brought to you by the The Research Repository @ WVU with permission from the rights-holder(s). You are free to use this Dissertation in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you must obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This Dissertation has been accepted for inclusion in WVU Graduate Theses, Dissertations, and Problem Reports collection by an authorized administrator of The Research Repository @ WVU. For more information, please contact researchrepository@mail.wvu.edu.

**Alterations in T cell function and activation during exposure to the herbicide 3,4
dichloropropionanilide (DCPA) and its metabolites**

Tricia L. Lewis

Dissertation submitted to the
School of Medicine
at West Virginia University
in partial fulfillment of the requirements
for the degree of

Doctor of Philosophy
in
Immunology and Microbial Pathogenesis

John B. Barnett, Ph.D., Chair
Peter M. Gannett, Ph.D.
Michael I. Luster, Ph.D.
Karen H. Martin, Ph.D.
Vazhaikkurichi M. Rajendran, Ph.D.

Department of Microbiology, Immunology and Cell Biology
Morgantown, West Virginia

2010

Keywords: 3,4-dichloropropionanilide, DCPA, propanil, 3,4-dichloroaniline, herbicide,
calcium, T cells, Stim1, metabolites

ABSTRACT

Alterations in T cell function and activation during exposure to the herbicide 3,4 dichloropropionanilide (DCPA) and its metabolites

Tricia L. Lewis

Approximately 5.3 billion pounds of pesticides are applied annually across the United States and 15 of the top 25 most used pesticides are herbicides. Total herbicide use in the United States represents 28% of all worldwide herbicide use. DCPA (3,4 dichloropropionanilide, common name propanil) is a post emergent herbicide that is used extensively for control against several broadleaf plants and grasses. It is the 17th most common herbicide in the United States and 6-9 million pounds are applied annually to 2 million acres of rice fields. In mammals, DCPA is metabolized in the liver and produces 3,4-dichloroaniline (DCA) and propionic acid as its major metabolites. DCA is further biotransformed leading to the production of 2 hydroxylated metabolites; 6-hydroxy-3,4-dichloroaniline (6OH-DCA) and N-hydroxy-3,4-dichloroaniline (NOH-DCA). The immunomodulatory effects of DCPA are well documented but only limited data is available on the effects of its metabolites. Previous studies have shown that DCPA alters transcription factors involved in the expression of IL-2 and decreases mRNA and IL-2 protein in human and mouse T cells. IL-2 is an early cytokine that is secreted by activated T cells and plays an important role in the activation, proliferation and differentiation of several immune cells. Expression of IL-2 relies on activation and influx of calcium through channels in the plasma membrane. This study was conducted to examine the effects of DCPA and its metabolites on T cell activation and function and to propose a mechanism for the observed effects. Human Jurkat T cells, a model cell line for T cell signaling, were exposed to increasing concentrations of DCPA or its metabolites and T cell function was assessed by measuring IL-2 secretion. DCPA and its metabolites all inhibit IL-2 secretion in a concentration-dependent manner, however, NOH-DCA is the most potent inhibitor, followed by DCPA. To better understand the mechanism by which they suppress IL-2, NFAT activity and calcium influx were investigated. Interestingly, DCPA and DCA inhibited IL-2 in a calcium-dependent manner whereas the hydroxylated metabolites inhibited IL-2 in a calcium-independent manner. The calcium-dependent alterations in IL-2, NFAT and calcium influx are influenced by the presence of chlorines, a substitution with fluorines abrogated all effects. Further studies investigating the role of DCPA in calcium release-activated calcium (CRAC) channels revealed that activation of a key protein, Stromal interaction molecule-1 (Stim1), is inhibited by DCPA. Collectively, this data supports the conclusion that DCPA suppresses IL-2 production by inhibiting Stim1. This mechanism describes a novel pathway for immunosuppression.

DEDICATION

I dedicate this dissertation to David Light and our children, Summer and Willem. This work is a testament to their love and patience. It is my hope that I may instill in my children a curiosity and love of learning that will guide and inspire them in all they do. I would also like to dedicate this dissertation in memory of Dr. George Wollner, my uncle, who understood the hard work and sacrifice required to achieve this goal.

ACKNOWLEDGEMENTS

The completion of this work would not have been possible without the encouragement, love and support of so many people. My sincere thanks to my advisor and mentor, John Barnett, whose limitless patience and thoughtful guidance through many obstacles made the completion of this dissertation possible. I also acknowledge my committee members Vazhaikkurichi Rajendran, Peter Gannett, Karen Martin and Michael Luster. I would especially like to thank Karen Martin for all her assistance and suggestions with several of my microscopy experiments and Peter Gannett for his help and insight with all things chemistry. I am also thankful for the valuable comments and suggestions of Rosana Schafer and David Klinke.

I would also like to acknowledge Miranda Hanson, a wonderful and strong friend who kept things in perspective and made me laugh. I am eternally grateful to my fellow grad students Siera Talbott, Laura Kelley and Janna Jackson for helping me maintain my sanity through energizing bike rides and great conversations, which often occurred at the same time. My thanks to Siera for her help with experiments and for a shoulder to laugh and cry on and to Laura for your critical eye and constructive and honest comments. I also want to thank Ron Fecek, Alex Rowe, Irina Ustyugova and Cheryl Walton for their friendship and help throughout my graduate studies.

My family provided endless love and support at times when I needed it most. I would like to thank my sister, Christine, who listened and unconditionally supported me during this long journey and to my Grandparents, Auntie and late Uncle George for their constant support and for encouraging me to reach my goals.

Lastly, I would like to thank my greatest supporter, David Light. Throughout this journey he has been steadfast through all the ups and downs and has sacrificed everything to see me through my graduate studies. No words can convey the deep gratitude and respect I have for his unwavering love and support. He made this achievement possible and I thank him with all my heart.

LIST OF ABBREVIATIONS

2,4-D	2,4-Dichlorophenoxyacetic acid
2-APB	2-Aminoethoxydiphenyl borate
6OH-DCA	6-hydroxy-3,4-dichloroaniline
7-AAD	7-amino-actinomycin
AP-1	Activator protein-1
ASC	Antibody secreting colony
ATP	Adenosine triphosphate
AUC	Area under the curve
BFU	Burst forming unit
Ca	Calcium
CAD	CRAC-activating domain
CD	Cluster of differentiation
CFU	Colony forming unit
con-A	Concavalin-A
CRAC	Calcium release-activated calcium
DAG	Diacylglycerol
DCA	3,4-dichloroaniline
DCPA	3,4-dichloropropionanilide
DDT	Dichlorodiphenyltrichloroethane
DFA	3,4-difluoroaniline
DFPA	3,4-difluoropropionanilide
DMSO	Dimethylsulphoxide
DTH	Delayed type hypersensitivity
ER	Endoplasmic reticulum
Hb	Hemoglobin
HEK293	Human embryonic kidney
hr	Hour
i.p.	Intraperitoneal
I_{CRAC}	Calcium release-activated calcium current
IFN	Interferon
Ig	Immunoglobulin

IL	Interleukin
IP ₃	Inositol 1, 4, 5 triphosphate
lb	Pound
LD ₅₀	Lethal dose 50
LPS	Lipopolysaccharide
MLR	Mixed lymphocyte reaction
Na	Sodium
NADPH	Nicotinamide adenine dinucleotide phosphate
NFAT	Nuclear factor of activated T cells
NF-κB	Nuclear factor kappaB
NK	Natural killer
NOH-DCA	N-hydroxy-3,4-dichloroaniline
PFC	plaque forming colony
PHA	Phytohemagglutinin
PIP ₂	Phosphatidylinositol 4,5-bisphosphate
PKC	Protein kinase C
PLC	Phospholipase C
PMA	Phorbol myristate acetate
PMCA	Plasma membrane Ca-ATPase
ppb	parts per billion
ppm	parts per million
SAM	Sterile alpha motif
SCID	Severe combined immunodeficiency
SERCA	Sarco/Endoplasmic reticulum Ca-ATPase
SHD	Stim1 homerization domain
SOCE	Store-operated calcium entry
Stim	Stromal interaction molecule
TBAB	Tetra-n-butylammonium bromide
TCAB	3,3',4,4' tetrachloroazobenzene
TCR	T cell receptor
Tg	Thapsigargin
TNF	Tumor necrosis factor
TOTP	triorthotolyl phosphate

TABLE OF CONTENTS

ABSTRACT	ii
DEDICATION	iii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS.....	vii
CHAPTER 1 Literature Review	1
DCPA (3,4-dichloropropionanilide)	1
Background	1
Methods and rates of application.....	2
Routes of exposure and risk assessment	2
Phytotoxicity	3
Metabolism	4
Figure 1: Metabolic pathway of DCPA in mammals	6
Ecological effects	7
DCPA metabolites	9
3,4-Dichloroaniline.....	9
N-hydroxy-3,4-dichloroaniline and 6-hydroxy-3,4-dichloroaniline	10
TOXICITY	10
General toxicity	10
Mutagenicity and carcinogenicity.....	11
ACUTE TOXICITY	12
General toxicity	12
Erythrocyte toxicity.....	13
CHRONIC TOXICITY	14
Reproductive Toxicity	15
Nephrotoxicity	15
IMMUNOTOXICITY	16
INNATE AND ADAPTIVE IMMUNITY	17
Innate Immunity	17

Adaptive Immunity	18
T CELL ACTIVATION AND SIGNALING	22
Figure 2 Schematic of T cell signaling	24
T CELLS AND ION CHANNELS	26
CRAC channels	26
CRAC regulation	27
CRAC inhibitors	28
OTHER ION CHANNELS AND PUMPS.....	29
SERCA	29
PMCA	29
RyR	29
Na/Ca exchanger.....	29
Ca uniporter	30
Potassium (K ⁺) channels	30
Non-store-operated cation channels.....	30
REFERENCES	31
CHAPTER 2.....	39
ABSTRACT.....	40
INTRODUCTION.....	41
MATERIALS AND METHODS	45
RESULTS.....	53
Figure 1 DCPA decreases intracellular calcium.....	56
Figure 1B	57
Figure 1C and 1D.....	58
Figure 2A. DCPA does not affect early IP3	60
Figure 2B	61
Figure 2C and 2D	62
Figure 3A. The effect of DCPA and 2-APB	65
Figure 3B and 3C	66
Figure 4 DCPA does not alter the membrane potential	68
Table 1. Effects of DCPA on IL-2.....	70

Figure 5A. DCPA decreases nuclear NFAT.....	72
Figure 5B and 5C	73
Figure 6. DCPA decreases $[Ca^{2+}]_i$ in primary mouse	75
DISCUSSION.....	76
REFERENCES	81
CHAPTER 3.....	85
ABSTRACT	86
INTRODUCTION	87
Figure 1A. DCPA metabolic pathway and structures	90
Figure1B. Stucture of flurorine analogs.	90
MATERIALS AND METHODS	91
RESULTS.....	97
Figure 2A and 2B Cytotoxic effects	98
Figure 2C and 2D Cytotoxic effects	99
Figure 3A and 3B DCPA and DCA inhibit IL-2	102
Figure 3C and 3D NOH-DCA and 6OH-DCA inhibit IL-2	103
Figure 4A DCPA alters NFAT	106
Figure 4B DCA alters NFAT	107
Figure 4C. 6OH-DCA alters NFAT.....	108
Figure 4D. NOH-DCA alters NFAT	109
Figure 5A DCPA alters calcium	111
Figure 5B DCA alters calcium	112
Figure 5C 6OH-DCA does not alter intracellular calcium	113
Figure 5D NOH-DCA does not alter intracellular calcium	114
Figure 6A and 6B Cytotoxic effects of DFPA	116
Figure 6C and 6D DFPA and DFA do not inhibit IL-2	117
Figure 6E and 6F DFPA and DFA DCPA do not alter NFAT	118
Figure 6G and 6H DFPA and DFA do not alter intracellular calcium.....	119
DISCUSSION.....	120
REFERENCES	126
CHAPTER 4.....	130

ABSTRACT.....	131
INTRODUCTION.....	132
Calcium signaling	132
Stromal Interaction Molecule-1 (Stim1)	133
Figure 1. Domain structures in human Stim1	135
Knockdown and mutational studies of Stim1	135
Stim2	136
Orai / CRACM	138
Figure 2. Schematic of Stim1-mediated Orai1 activation of CRAC	140
Orai2/CRACM2 and Orai3/CRACM3	141
MATERIALS AND METHODS	143
RESULTS	145
Figure 3A Formation of Stim1 puncta	146
Figure 3B DCPA inhibits formation of Stim1	147
DISCUSSION	148
REFERENCES	151
CHAPTER 5.....	157
GENERAL DISCUSSION	157
REFERENCES	167

CHAPTER 1

Literature Review

DCPA (3,4-dichloropropionanilide)

Background

DCPA (chemical name 3,4-dichloropropionanilide, common name propanil) is the active ingredient in a commercially available herbicide distributed under the several trade names, including Chem-Rice, Herbax and Propanex. It is a widely used, post-emergent acetanilide contact herbicide and is registered for use against several broadleaf and grassy weeds, such as barnyard grass, crabgrass, pigweed, foxtail and smartweed (USEPA 2006). It has been registered for use as an herbicide in the United States (US) since the 1950s and has recently been required by the United States Environmental Protection Agency (USEPA) to be re-registered due to increased adverse ecological and toxicological effects (USEPA 2006). In the US, DCPA is applied predominantly on rice fields, with lesser use on small grain and wheat crops. Annual use of DCPA is estimated to be 6-9 million pounds per year and is distributed on approximately 2 million acres of crop land (USDA 2006). In the United States, 50-80% of total rice crops are treated with propanil, with major use in California, the Mississippi Delta and the Gulf Coast (USDA 2006). In addition to crop use, DCPA has also recently been registered for turf use on commercial sod farms (USEPA 2006). This new application of DCPA will further increase use and exposure.

Methods and rates of application

Ground boom sprayers and aerial equipment are the most common means of application with hand spraying also occurring in some areas. Commercial DCPA labels indicate that handlers should wear long sleeve shirts, long pants, socks, shoes, waterproof gloves and protective eyewear (USEPA 2006). In addition, in some situations, workers must use maximum protective controls, including closed mixing and loading systems and enclosed cockpits and trucks (USEPA 2006). Application of DCPA to rice fields typically occurs twice, once approximately 15-25 days after the rice is planted and again 10-15 days later. At this time, the rice fields are flooded for 30 days and a third application is possible during the flooding period. After the flooding period, crop fields are drained and the rice is subsequently harvested 140-150 days after planting. The maximum rate of application of propanil, set by the USEPA, on rice fields is 8 lb/acre/season with an average use of 6 lb/acre/season (USEPA 2006). Application of DCPA on turf in commercial sod farms has a maximum rate of 10 lb/acre (USEPA 2006).

Routes of exposure and risk assessment

Workers involved in manufacturing, handling or application of propanil are at greatest risk but there have also been reports of non-occupational exposure, including a child under the age of 6 (USEPA 2006). The routes of exposure include dermal, inhalation, ingestion and contact with eyes. The USEPA's Registration Eligibility Decision (RED) for propanil conducted a risk assessment which included determining a margin of exposure (MOE) for workers who are involved in mixing, loading and application activities (USEPA 2006). For propanil, MOE values of less than 300 are

considered a potential risk concern. Commercial propanil labels indicate that handlers should wear long sleeve shirts, long pants, socks and shoes, waterproof gloves and protective eyewear. Using this minimal and widely used level of protection, the EPA determined that the MOE of all workers (in rice fields and turf) was less than 300, with several handling methods having a MOE less than 100. In addition, in some situations, workers using maximum protective controls (closed mixing and loading systems and enclosed cockpit, cabs or trucks) still had a MOE of less than 300. Post occupational risk assessment, for workers entering treated areas 12 hours after application, was also conducted. Workers entering a field with maximum allowable application of propanil had a MOE of less than 300 while turf workers involved in transplantation, fertilizing or mechanical weeding or harvesting had a MOE less than 100. This assessment indicates that exposure to propanil is a risk concern for all workers, including those with maximal protection.

Phytotoxicity

In rice, as well as broadleaf plants and grasses, DCPA can translocate from the leaves to the growing shoots and then back to other leaves. The translocation and absorption of DCPA occurs at similar rates in both rice and plants targeted by the herbicide (Carey V. F. 1995). The phytotoxic effects of DCPA are influenced by the ability of the plant to produce the enzyme acylamidase (Adachi 1966; Still and Kuzirian 1967). Acylamidase cleaves the amide bond on DCPA and inactivates its herbicidal activity (Yih, McRae et al. 1968). Gaynor *et al* determined that, in the rice plants (*Oryza sativa*), acylamidase is primarily localized to the outer membrane of the mitochondria (Gaynor and Still 1983). Rice leaves produce 60 times more acylamidase

than barnyard grasses and are able to degrade DCPA to avoid its phytotoxic effects. Inhibition of acylamidases can occur during co-application with DCPA and certain organophosphorus or carbamate insecticides rendering the rice plants susceptible to the phytotoxic effects of propanil (Matsunaka 1968; Chang, Smith et al. 1971). Plants susceptible to the herbicidal activity of DCPA express low levels of acylamidase and phytotoxic effects occur primarily through the inhibition of photosynthesis. Specifically, DCPA inhibits the photosynthetic electron chain transport system in chloroplasts (Matsunaka 1968). During photosynthesis two distinct stages are required for the synthesis of sugars. In the light reaction chlorophyll absorbs a photon of light and loses an electron. This electron gets passed through the electron chain transport resulting in the production of NADPH (nicotinamide adenine dinucleotide phosphate). In addition, the light reaction also converts light energy into chemical energy stored in the form of ATP (adenosine triphosphate). Both NADPH and ATP are required for the conversion of CO₂ to sugar that occurs in the dark reaction in the stroma of chloroplasts. Interruption of the electron chain transport prevents the production of sugars required for plant growth. Recently Yun *et al* reported that DCPA also inhibits the enzyme, 4-coumarate:CoA ligase (Yun, Chen et al. 2007). 4-Coumarate is found only in the plant kingdom and upon binding to the cofactor, CoA ligase, plays an important role in the phenylpropanoid pathway, a pathway important in plant growth, mechanical support and cell wall rigidity.

Metabolism

In soil, plants and mammals, metabolism of DCPA occurs by hydrolysis of the amide bond resulting in the production of 3,4-dichloroaniline (DCA) and propionic acid (Still

1968). When McMillan *et al* treated rat hepatic microsomes with radio-labeled DCPA they identified DCA as the major metabolite (McMillan, Freeman *et al* . 1990). Hydrolysis of the amide bond by hepatic acylamidases was supported when formation of DCA was inhibited with acylamidase inhibitors. In addition, they identified 2 minor metabolites produced through the oxidation of DCA; 6-hydroxy-3,4-dichloroaniline (6OH-DCA) and N-hydroxy-3,4-dichloroaniline (NOH-DCA) (McMillan, Leakey *et al* . 1990). Figure 1 outlines the metabolic pathway in mammals and the structures of DCPA and its major metabolites.

The half life of DCPA in soil is reported to be 2-3 days (Chisaka and Kearney 1970; Konstantinou, Zaradis *et al* . 2001) . Recovery of propanil, in sterile-treated soil, is almost 100% after incubation for 25 days (Chisaka and Kearney 1970). Analysis of soil samples from rice fields treated with DCPA reveals that most of the DCPA is metabolized within 15 days and DCA and propionic acid can be readily detectable. Microbial activity in the soil is thought to be responsible for the metabolic breakdown of propanil. Acylamidase activity in bacteria including *Pseudomonas striata* and *Pseudomonas fluorescens* as well as the fungus *Vibrio fisheri* can convert DCPA to DCA (Surovtseva and Funtikova 1978; Zablotowicz, Locke *et al* . 2001) . *Fusarium solani* has also been reported to hydrolyze DCPA to produce DCA and propionic acid (Lanzilotta and Pramer 1970). The propionic acid can serve as a carbon source to *F. solani* to provide energy to the fungus. However, it does not appear to be able to further degrade DCA and an increasing accumulation of DCA can result in inhibited growth of the fungus (Lanzilotta and Pramer 1970). Microbial oxidation of propionic acid releases CO₂ and water within 4 days (Yih, McRae *et al* . 1968; Chisaka and Kearney 1970) .

Figure 1: Metabolic pathway of DCPA in mammals

Soils treated with propanil, at near-field application rates (6lb/acre), reveal that 70-80% of the DCA produced is chemically bound to humic substances (Bartha and Pramer 1967; Bartha 1971). Degradation of the DCA-humic complex was estimated to take 5-10yrs. DCA can also complex with the plant protein, lignin, with no apparent detrimental effects to the plant (Yih, McRae et al. 1968). A small portion (10%) of

unbound DCA can also form complexes with sugars to produce N-(3,4-dichlorophenyl) glucosylamine and other minor sugar conjugates (Still and Kuzirian 1967).

There are also reports that two molecules of DCA, derived microbiologically from propanil, condense and form 3,3',4,4' tetrachloroazobenzene (TCAB) (Bartha 1971; Hill, Rollen et al. 1981). It has been suggested that microbiological action, specifically peroxidase activity, is involved in the condensation of DCA to TCAB in soils (Bartha and Pramer 1967; Bordeleua 1972). TCAB has not been found in plant tissues treated with DCPA (Still, Balba et al. 1981). In a study examining 99 soil samples from the rice-growing states of Arkansas, California, Louisiana, Mississippi, and Texas, TCAB was detected in 6 samples (Carey, Yang et al. 1980). Production of TCAB is thought to occur primarily in loamy soil and is inhibited under dry soil condition (Bartha 1971). Workers involved in the manufacturing and packaging of DCPA report several adverse exposure effects including dizziness, nausea, muscle weakness, fatigue, and chloracne. Only the effect of chloracne was directly related to the presence of TCAB (Kimbrough 1980).

Ecological effects

The effect of DCPA on microbial communities and aquatic animals and plants has been well documented. Contamination of waterways has been reported to occur following release of water from flooded rice fields with reported DCPA levels of 0.1 µg/l (0.1 ppb) in soil water slurry and irrigation water (Papadopoulou-Mourkidou, Karpouzas et al. 2004). A single application of DCPA (6 lb/ac), in a closed flooded rice field, resulted in a DCPA concentration of 200 µg/l (200 ppb) after 24 hrs and decreased to 50 µg/l (50 ppb) for the next 5 days (Papadopoulou-Mourkidou, Karpouzas et al. 2004).

After 10-15 days only metabolites were detectable. The half life of DCPA in water is reported to be approximately 2 days (Dahchour, Bitton et al. 1986).

Several studies have documented the effects of DCPA on aquatic life. Exposure of DCPA to fathead minnows (*Pimephales promelas*) has several deleterious effects with an LD₅₀ value of 3.4mg/l (3.4ppm) after 192 hours of exposure (Call, Brooke et al. 1983). Early life toxicity was observed at 3.8µg/l (3.8ppb) and resulted in significant decreases in egg hatch and increases in dead and deformed hatched minnows. By post-hatch day 54 the mortality rate was 100%. Significant decreases in survival were also seen at 1.2µg/l (1.2ppb), with only 17% surviving past post-hatch day 54. These levels are sufficiently high enough to affect fish populations in the rice paddy water or in the paddy water that is released near spawning or nursery grounds. Adverse effects to DCPA have also been observed in fish and frogs. The frog species, *Xenopus laevis*, has a reported LD₅₀ of 8.64mg/l (8.64ppm) 48 hours after exposure. The 96 hr LC₅₀ for the fish species, Rainbow trout (*Oncorhynchus mykiss*), Bluegill (*Lepomis macrochirus*), and the fingerling channel catfish (*Ictalurus punctatus*) exposed to DCPA ranged from 2.3 - 6mg/l (2.3-6ppm) (McCorkle, Chambers et al. 1977) (Moore, Pierce et al. 1998) (USEPA 2003). Exposure of the metabolite DCA on zebrafish, *Danio rerio*, results in alterations in early life stage development (Nagel, Bresche et al. 1991). Deformations in the spine were observed at 0.25mg/l DCA and an 11 day subchronic test (0.5mg/l) revealed decreases in locomotor activity and mortality. *Daphnia magna* are also very sensitive to DCA as the 48-hr LC₅₀ of 0.14 mg/L (0.14ppm) (Pereira, Antunes et al. 2009).

Microbial communities within rice fields are also altered during exposure to DCPA. Exposure of 20µg/ml (20ppm) DCPA to the cyanobacterium, *Anabaena MH*, resulted in a significant inhibition of growth (Habte and Alexander 1980) with a loss of more than 85% of its chlorophyll with no recovery after 10 days of exposure. Application of 25µg/ml (25ppm) suppressed 98% of algae growth for over 30 days.

DCPA metabolites

3,4-Dichloroaniline

3,4-Dichloroaniline (DCA) is the major metabolite of DCPA as well as other herbicides, including diuron and linuron. In addition, DCA is used as an intermediate in the manufacturing of several pesticides, dyes and pharmaceuticals (ECB 2006). DCA is persistent in soil and water but can be readily photodegraded in the atmosphere. It has a half-life of approximately 1000 days (ECB 2006). DCA is subject to chemical oxidation, reacts with the organic matter in soil and has high volatility. In areas treated with propanil, large amounts of DCA are bound to soil particles making it unavailable for further metabolism or plant uptake (Bartha 1971). However, DCA residues can be detected in most commercial rice (*Oryza sativa*) grain samples at a concentration of 1µg/g of rice (Still 1968; SCTEE 2001). Rats treated with an acute intraperitoneal (i.p.) injection of DCA reach maximum detection of DCA in the liver and plasma within 30-60 minutes with levels decreasing quickly within a few hours (Guilhermino, Soares et al. 1998). Elimination of DCA in rats occurs after 3 days and is primarily excreted in urine and feces. In humans, DCA has been detected in the urine of occupational and non-occupationally exposed subjects (Wittke, Hajimiragha et al. 2001; Turci, Barisano et al.

2006) with levels in the 0.01-6.2µg/l range. DCA-hemoglobin (Hb) adducts have also been detected in workers involved in the application of DCPA to rice fields (Pastorelli 1998). These adducts were still detectable 4 months after the last application of propanil.

N-hydroxy-3,4-dichloroaniline and 6-hydroxy-3,4-dichloroaniline

N-hydroxy-3,4-dichloroaniline (NOH-DCA) and 6-hydroxy-3,4-dichloroaniline (6OH-DCA) are the major mammalian metabolites of DCA. In preparations of rat hepatic microsomes, McMillan *et al* detected 2 oxidative metabolites, NOH-DCA and 6OH-DCA (McMillan, Freeman *et al*. 1990). Further studies in chromium-51 (Cr-51) labeled erythrocytes treated with DCPA indicate that NOH-DCA can undergo rapid redox cycling resulting in the oxidation of oxyHb to metHb. Little is known about the effects of 6OH-DCA but it is also involved in the production of metHb, although the NOH-DCA metabolite is reported to be 10 times more potent (McMillan, Freeman *et al*. 1990). Methemoglobin (metHb) is a form of hemoglobin that does not bind or transport oxygen. Increases in metHb leads to a serious medical disorder known as methemoglobinemia.

TOXICITY

General toxicity

Using metHb levels as a marker for toxicity, the USEPA has assessed a toxicity category for DCPA based on routes of exposure (USEPA 2003). Inhalation and dermal routes, the most common routes of exposure, were placed in Toxicity Category IV. This indicates that caution should be used but little toxicity has been shown. An oral route of

exposure is assigned a Toxicity Category II, with caution indicating slight toxicity. Toxicity Category II, with a hazard warning for moderate toxicity, was indicated as studies demonstrated that DCPA is an eye irritant.

Mutagenicity and carcinogenicity

Conflicting studies are available that assess the possible mutagenic effects of exposure to DCPA and its metabolites. Using the *Salmonella typhimurium* reversion assay, chemical-induced base changes or frameshift mutations in the genome of this organism can be measured (McMillan, Shaddock et al. 1988). Over a range of concentrations, propanil, DCA and NOH-DCA demonstrated no increase in mutations (McMillan, Shaddock et al. 1988). The ability of propanil, DCA and NOH-DCA to induce mutations in Chinese hamster ovary (CHO) cells was also assessed. Although exposure of all 3 chemicals decreased CHO cell viability over a wide range of concentrations, no changes in mutation rates were observed (McMillan, Shaddock et al. 1988). Similarly, no change in DNA damage in rat hepatocytes was observed. However, in more recent studies using the more sensitive *Drosophila* wing spot assay, larvae were treated for 3 days with varying concentrations of DCPA. The wing spot assay can detect a loss of heterozygosity in suitable recessive markers that leads to the formation of mutant clones of cells that are then expressed as spots on the wings. In these assays, DCPA does induce genotoxic effects (Kaya, Creus et al. 2000). A cytogenic assay on DCA indicated that exposure of DCA disturbs spindle formation but with no changes in chromosome formation (Bauchinger, Kulka et al. 1989).

The USEPA has assessed DCPA for carcinogenic potential and has determined that there is “suggestive evidence of carcinogenic potential by all routes of exposure but not sufficient to assess human carcinogenic potential.” {USEPA, 2003 #555}

ACUTE TOXICITY

General toxicity

Early studies on the acute toxicological effects of DCPA established LD₅₀ values for oral exposure in rats and dogs (Ambrose, Larson et al. 1972). An acute oral LD₅₀ of 1384mg/kg for rats and 1217mg/kg for dogs was reported. Oral exposure to DCPA over a 7 day period resulted in toxicity characterized by central nervous system depression within 12 hours of exposure. In studies in Wistar rats, oral doses between 39-170mg/kg of DCA can alter body, spleen, liver and testis weight (Zhang and Lin 2009). Inhalation studies set an LD₅₀ concentration of 1.1mg/kg for a 4hr exposure but LD₅₀ concentrations of more than 5000mg/kg were reported for dermal exposure in rabbits (Stevens 1991). Acute studies in male mice were conducted using a single i.p injection of DCPA at a dose of 400mg/kg (Singleton and Murphy 1973). Signs of toxicity included central nervous depression, loss of righting reflex, and cyanosis. At higher doses, 600 and 800 mg/kg, deaths occurred within 6 to 24 hours after injection. Rats treated with 300mg/kg DCPA and triorthotolyl phosphate (TOTP), an enzyme that inhibits amidases, displayed no signs of cyanosis but the loss of righting reflex, CNS depression or mortality remained (Singleton and Murphy 1973). This data suggests that cleavage of the amide bond is necessary for cyanosis.

Erythrocyte toxicity

Due to the nature of exposure there are limited studies on the effects of DCPA on humans. However there are cases of occupational exposure and DCPA self-poisoning and that help characterize the toxicity of DCPA. DCPA self-poisoning has been reported in Cuba and Japan, but in Sri Lanka, DCPA is the second most lethal herbicide used in deliberate self-poisoning cases (Eddleston, Rajapakshe et al. 2002). Between 2002 and 2007 over 400 patients were identified in self-poisoning cases (Roberts, Heilmair et al. 2009). Symptoms of DCPA poisoning included nausea, vomiting, dizziness, increased heart rate, tissue hypoxia, cyanosis, methemoglobinemia and depression of CNS and respiratory system (Morse, Baker et al. 1979; Kimbrough 1980). Methemoglobinemia is a well characterized effect of DCPA exposure in humans and animal models. Methemoglobin (metHb) is a form of hemoglobin that does not bind or transport oxygen. MetHb levels greater than 20% can result in decreased consciousness, convulsions, shock and, metHb above 70% results in death. Approximately 10% of all patients in Sri Lanka died despite active treatment (Roberts, Heilmair et al. 2009). The reported half life of DCPA, in the human body, was 3.2 hours but the concentration of DCA was detectable and more persistent than that of DCPA (Roberts, Heilmair et al. 2009).

The mechanism for DCPA toxicity on erythrocytes is due to the formation of the hydroxylated metabolites, N OH-DCA and 6OH-DCA (Singleton and Murphy 1973; Guilhermino, Soares et al. 1998). *In vitro* studies in erythrocytes from Sprague-Dawley rat exposed to DCA revealed no increase in metHb levels (McMillan, McRae et al. 1990). However, exposure to N OH-DCA and 6 OH-DCA produced increased levels of

metHb with maximum metHb levels by 90 minutes. Sprague-Dawley rats injected with DCPA (100mg/kg) had detectable levels of DCA and NOH-DCA, in the blood, which reached peak concentration after 20mins and 4.5 hrs, respectively (McMillan, McRae et al. 1990). During methemoglobinemia, oxygen carrying the ferrous ion (Fe^{2+}) of the heme group of the hemoglobin molecule is oxidized by the oxidative metabolites of DCPA to the ferric state (Fe^{3+}). This converts hemoglobin to methemoglobin, a non-oxygen binding form of hemoglobin that binds a water molecule instead of oxygen. Although both NOH-DCA and 6OH-DCA are known to form metHb, the NOH-DCA metabolite is reported to be 10 times more potent (McMillan, Freeman et al. 1990)

In addition to methemoglobinemia, hemolytic anemia has been observed as a result of exposure to propanil. *In vivo* and *in vitro* studies in Sprague-Dawley rat erythrocytes demonstrated that NOH-DCA induced hemolytic anemia, but DCPA and DCA were not direct-acting hemolytic agents (McMillan, Bradshaw et al. 1991). In addition, although 6OH-DCA can induce methemoglobinemia, it does induce hemolytic anemia (Chow and Murphy 1975).

CHRONIC TOXICITY

Chronic toxicity studies were conducted in rats and dogs over a 2 year period and various parameters were measured (Ambrose, Larson et al. 1972). In rats, significant decreases in body weight (from onset), mortality (after 20 months) and Hb levels (after 3 months) were observed at 1600ppm. In addition, higher organ-to-body weight ratios were also reported in the spleen (both sexes), liver (females only) and testes at the 1600ppm dose. Studies in dogs revealed no changes in mortality, hematologic values, or organ-to-weight ratios over a 2 year period. Reproduction

studies in rats revealed no adverse effects in weight, reproductive performance, litter size or litter mortality and no histopathological changes in weaned pups. During a 3 month study of subchronic oral exposures in rats, survival was only affected at the highest concentration (50,000ppm) (Ambrose, Larson et al. 1972). Decreases in Hb levels were observed at all doses above 1000ppm indicating hemolytic anemia. Several increases in organ-to-body weight ratios were also observed. At a dose of 3300ppm increases in kidney and liver ratios in only the females were observed. Increases in spleen ratio were observed in only the females at the 1000ppm dose. At the higher dose (10000ppm) increased testes ratio in the males and increased heart ratios in both the males and females were noted. A similar 4-week study was conducted in dogs with decreased food consumption and weight observed only the two highest doses (10000 and 50000ppm) (Ambrose, Larson et al. 1972).

Reproductive Toxicity

In studies with Wistar rats, oral doses between 39-170 mg/kg of DCA decreased testis weight. In addition, alterations in several testicular enzymes suggests modulatory effects of spermatogenesis, testicular degeneration, and possible deterioration of the germinal epithelium (Zhang, Pan et al. 2009). DCA has also been shown to weakly bind the androgen receptor and is suspected as acting as an endocrine disruptor (Bauer, Meyer et al. 1998; USEPA 2006).

Nephrotoxicity

There are various reports on the toxicity of NOH-DCA and DCA in the kidney. Exposure to DCPA, DCA and NOH-DCA has been reported to result in renal cytotoxicity in Fisher 344 rats (Valentovic, Yahia et al. 1997; Valentovic, Ball et al. 2001). In these

ex vivo experiments, renal cortical slices were exposed to varying concentrations of DCPA and its metabolite and cytotoxicity was determined by measuring lactate dehydrogenase (LDH) levels. Exposure to 500 μ M NOH-DCA resulted in significant cytotoxicity but higher concentrations were required for similar results in renal cortical slices exposed to DCPA (1mM), DCA (2mM) and propionic acid (5mM). Decreases in glutathione levels were observed and cytotoxicity was reversed by addition of glutathione in NOH-DCA exposed samples, indicating an increase in the oxidation of glutathione in order to neutralize the intermediate NOH-DCA.

IMMUNOTOXICITY

The immune system is a complex system comprised of innate and adaptive responses that require surveillance and balance to ensure an optimal response to foreign attacks. In humans, the development of immune cells begins in the bone marrow. Pluripotent stem cells can differentiate down 2 pathways: myeloid or lymphoid. Differentiation down the myeloid pathway results in the production of colony forming units-spleen (CFU-S) progenitor cells that give rise to erythrocytes, platelets, granulocytes and monocytes. In a C57/B6 mouse model of acute exposure to DCPA, there was a dose dependent decrease in CFU-S cells as well erythroid burst forming units (BFU-E), progenitor cells for erythrocytes (Blyler, Landreth et al. 1994) Similar results are also reported in human blood cord progenitor cells but exposure to DCA appears less toxic (Malerba, Castoldi et al. 2002).

Early toxicological studies revealed alterations in both the spleen and the thymus. The spleen-to-body weight ratio was increased in albino rats exposed to oral doses of 3300ppm of DCPA for 3 months and continued in chronic studies conducted over a 2

year period (Ambrose, Larson et al. 1972). In C57/B6 mice an i.p. injection of 200mg/kg of propanil, or a molar-equivalent dose of DCPA (150mg/kg) resulted in increased spleen weight and size (Barnett and Gandy 1989; Barnett, Gandy et al. 1992). Decreases in thymic weight were also observed in DCPA treated mice but not DCA treated mice (Barnett, Gandy et al. 1992; Zhao, Schafer et al. 1995). In addition to thymic atrophy, several thymocyte subpopulations were also decreased. Significant concentration dependent decreases in thymic CD3+CD4+CD8+ and CD3+CD4+CD8- populations were seen at all doses (100-200mg/kg) but no changes were observed in the spleen or mesenteric lymph nodes (Zhao, Schafer et al. 1995).

INNATE AND ADAPTIVE IMMUNITY

Innate and adaptive immune responses are the two main components of an active immune response. Innate immunity is a mechanism of protection that does not require specific recognition of antigens and is important in the early detection of pathogens. Adaptive immunity is an antigen-specific response that includes both humoral and cell-mediated immunity. Alterations in both innate and adaptive responses have been reported after exposure to propanil (reviewed in (Salazar, Ustyugova et al. 2008)).

Innate Immunity

Adverse effects of exposure of DCPA to macrophages have been well documented. *In vivo* studies with peritoneal macrophages from C57/B6 mice exposed to 200mg/kg DCPA resulted in the decreased production and secretion of IL-6 and TNF- α . In both oral and i.p. routes of exposure, DCPA (400mg/kg) significantly decreased

IL-6 and TNF- α production in LPS-stimulated macrophages (Xie, Schafer et al. 1997). *Ex vivo* experiments with peritoneal macrophages resulted in similar decreases (Xie, Schafer et al. 1997). In addition, the ability of LPS-stimulated peritoneal macrophages to phagocytose was also decreased with concomitant decrease in reactive oxygen species (ROS), reactive nitrogen species (RNS) and inducible nitric oxide synthase (iNOS) (Ustyugova, Frost et al. 2007). In the human monocytic cell line, THP-1, TNF- α secretion and phagocytosis was also inhibited (Ustyugova, Frost et al. 2007). The mechanisms for these decreases may be due to a decrease in the p65 subunit of NF- κ B and to alterations in intracellular calcium (Ca) homeostasis (Xie, Schafer et al. 1997; Frost, Neeley et al. 2001). In addition to macrophages, natural killer (NK) cells appear to be sensitive to the effects of DCPA and DCA. Exposure of DCPA and its major metabolite, DCA, results in decreased NK lytic activity. C57/B6 mice treated with 75-200mg/kg DCPA (or the molar equivalent of DCA) resulted in dose-dependent decreases in the ability of NK cells to lyse target cells (Barnett, Gandy et al. 1992).

Adaptive Immunity

B cell response

Recognition of foreign antigens and the production of antigen-specific antibodies are important functions for B cells. Immature B cells are produced in the bone marrow and develop from a progenitor B cell to a Pro-B cell to a pre-B cell and finally reach an immature B cell stage, where it expresses IgM, and migrates to the spleen for differentiation into a mature B cell. Several studies indicate that the humoral immune response is altered by DCPA and its metabolite, DCA. Mice treated with DCPA or DCA,

have increased spleen size and weight but the mechanism for this observation is unknown. Decreased pre-B and IgM populations in the bone marrow of C57/B6 mice exposed DCPA were observed 7 days post exposure but return to normal 24 days post-exposure (de la Rosa, Barnett et al. 2003). Exposure of C57/B6 mice to 400mg/kg of DCPA also reduced the proliferation of LP S-stimulated B cells (Barnett and Gandy 1989). Initial toxicological studies investigating the effects of DCPA on the immune response revealed that i.p. exposure of DCPA or DCA, in C57/B6 mice, leads to decreased in T cell-dependent and T cell-independent antibody responses (Barnett and Gandy 1989; Barnett, Gandy et al. 1992). Interestingly, C57/B6 mice immunized with heat-killed *Streptococcus pneumoniae* and exposed to DCPA have a dose-dependent increase in the number of phosphocholine (PC)-specific IgM, IgG2b and IgG3 antibody secreting B cells (ASC) in the spleen (Salazar, de la Rosa et al. 2005). At 7 days post-exposure, the number of ASC had increased 4-6 fold with no increases in the number of ASC in the bone marrow or serum. The reason for this increase has not been elucidated but the mechanism appears to be ovary-dependent, estrogen and progesterone-independent but requires a functioning steroid synthesis pathway (Salazar, Miller et al. 2006). Male mice produce a smaller increase in PC-specific ASC compared to females when exposed to DCPA but this increase is testes-independent (Salazar, Miller et al. 2006).

T cell response

T cells play an important role in cell-mediated immunity and involve specific recognition of foreign antigens leading to the production of antigen-specific antibodies through CD4+ T cells or the targeted destruction of infected cells (CD8+ T cells). Although there are several types of T cells, studies on the effects of DCPA are limited

to CD4⁺ helper T cells and CD8⁺ cytotoxic T cells. As mentioned above, exposure to DCPA results in a significant decrease in the CD3⁺CD4⁺CD8⁺ and CD3⁺CD4⁺CD8⁻ thymic subpopulations (Zhao, Schafer et al. 1995). Mixed lymphocyte reactions (MLR) and delayed-type hypersensitivity (DTH) reactions were both reduced in C57/B6 mice exposed to high doses of DCPA (400mg/kg) (Barnett and Gandy 1989). In addition, concavalin-A (con-A)-stimulated splenocytes, from mice treated with DCPA, resulted in decreased proliferation as well as decreased IL-2 and IL-6 production (Barnett and Gandy 1989; Zhao, Schafer et al. 1998). Further studies, using murine (EL-4) and human (Jurkat) T cell lines, demonstrated a dose dependent-decrease in IL-2 production, at low concentrations of DCPA (25-100 μ M) when stimulated with PMA or PMA/ionomycin (Io), respectively (Zhao, Schafer et al. 1999; Brundage, Schafer et al. 2004). In the EL-4 cell line, transcription and stability of the IL-2 mRNA was inhibited and in Jurkat cells, decreases in the amount and phosphorylation of c-jun, one of two proteins that make up the transcription factor AP-1, resulted in the decreased DNA binding ability of AP-1.

Cytotoxic CD8⁺ T cells are responsible for identifying infected cells and secreting lytic proteins, such as perforin, granzymes and granulysin, in order to kill infected cells. Investigations of thymic subpopulations demonstrated that CD3⁺CD4⁻CD8⁺ T cells were decreased (Zhao, Schafer et al. 1995). However, studies investigating the effects of DCPA and DCA, demonstrated that neither the parent compound nor its metabolite impaired cytotoxic T cell activity (Barnett, Gandy et al. 1992). More recently, it has been reported that although primary stimulation of CD8⁺ T cells does not impair lytic

function, secondary stimulation of mice CD8⁺ T cells, in the absence of further DCPA exposure, abrogated the lytic function (Sheil, Frankenberry et al. 2006).

T CELL ACTIVATION AND SIGNALING

T cell recognition of a foreign peptide presented on a major histocompatibility complex (MHC) on the surface of an antigen presenting cell (APC), along with co-stimulation of CD28, sets into motion a complex series of signaling events that culminates in the activation of a T cell (reviewed in (Smith-Garvin, Koretzky et al. 2009). The TCR is comprised of a series of $\gamma\epsilon$, $\delta\epsilon$, and $\xi\xi$ dimers that associate with a single $\alpha\beta$ heteromer. These dimers contain specific immunoreceptor tyrosine-based activation motifs (ITAMs) consisting of 2 tyrosines that flank an amino acid sequence that contains key leucine and isoleucines. Ligation of the TCR results in the recruitment of cytosolic Src family protein tyrosine kinases (Lyn and Lck) that phosphorylate tyrosines associated with ITAMs and thereby provide a docking site for Zap-70. Zap-70 is a 70kDa phosphoprotein belonging to the Syk kinase family. Recruitment of Zap-70 leads to a cascade of phosphorylation events important in the activation of downstream pathways. Two key targets of Zap-70 are the adapter proteins, linker of the activation of T cells (LAT) and Src homology-2 (SH2) domain-containing leukocyte phosphoprotein (SLP-76). These 2 proteins help to stabilize the TCR complex and organize effector proteins to allow for activation of multiple pathways. LAT proteins can bind the SH2 domain of PLC- γ and as well as adapter proteins GRB2 (growth factor receptor bound protein-2) and Gads (GRB2-related adapter downstream of Shc). SLP-76 interacts with the SH2 domains and activates adapter proteins Vav1, Nck and Itk (IL-2-induced tyrosine kinase). All these proteins help to increase the stability of the complex and interact to activate PLC γ -dependent pathways, including calcium (Ca^{2+}) and DAG-

mediated pathways, cytoskeletal rearrangements and integrin activation (Smith-Garvin, Koretzky et al. 2009).

Early signaling events, dependent on LAT and SLP-76 activation, result in the recruitment of PLC- γ_1 to the TCR complex. Once activated, PLC- γ_1 hydrolyzes the membrane lipid phosphatidylinositol 4, 5-bisphosphate (PIP₂) resulting in the production of diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP₃). DAG remains in the membrane where it plays a role in the activation of the Ras and PKC θ pathways. Ras-GRP (Ras guanyl nucleotide-releasing protein) contains a DAG-binding domain that recruits it to the membrane where it is phosphorylated by PKC θ and can then convert RAS-GDP to the active Ras-GTP. Ras-GTP activates Raf-1, a Ser/Thr kinase, and phosphorylates and activates MAPK (mitogen-associated protein kinase) pathway. Activation of the MAPK pathway leads to activation of Elk, a transcription factor important for the expression of c-fos. Dimerization of c-fos and c-jun form AP-1, an important transcription factor required for early gene expression. In addition to activation of the MAPK pathway, DAG also recruits PKC θ to the membrane through its DAG-specific lipid-binding domain and activates the NF- κ B pathway (Quest, Ghosh et al. 1997). Activation of NF- κ B occurs when its inhibitory molecule, I κ B, is phosphorylated by the I κ B kinase (IKK) complex (Wan and Lenardo 2010). Phosphorylation of I κ B marks it for ubiquitination and degradation and allows NF- κ B to translocate into the nucleus where it acts as a transcription factor for genes involved in T cell function. Figure 2 outlines a general schematic for T cell activation.

Figure 2 Schematic of T cell signaling

PLC- γ hydrolysis of PIP₂ also results in the production of IP₃. IP₃ is released into the cytosol where it binds to its receptor, IP₃-R on the ER membrane resulting in the release of Ca from the ER stores. Upon depletion of the ER stores, Stim1, a transmembrane protein on the ER, aggregates and initiates clustering of Orai1, a plasma membrane protein, leading to the activation of CRAC channels and the influx of Ca into the cell (described in detail in Chapter 5) (Barr, Bernot et al. 2008). This increase in cytosolic Ca, through the CRAC channels, triggers several signaling events that are critical for T cell function. NFAT is a transcription factor that is strictly regulated by Ca and cooperatively binds to DNA to control the expression of genes important in T cell activation and function (Jain, Loh et al. 1995; Rooney, Sun et al. 1995). In resting T cells, NFAT is phosphorylated at several Ser residues and resides in the cytosol but upon stimulation, is dephosphorylated by calcineurin and can freely translocate into the nucleus. The influx of Ca through CRAC channels following activation of T cells is essential for the activation of NFAT. Binding of Ca, to the 4 ion-binding sites on calmodulin (CaM), leads to the activation of calcineurin (Cn). Cn is a calmodulin-dependent serine/threonine phosphatase with 2 subunits; a catalytic subunit (calcineurin A) and a regulatory subunit (calcineurin B) (Feske 2007). The Ca-CaM complex binds to the regulatory subunit of Cn thereby activating its catalytic activity. Once activated, Cn dephosphorylates NFAT and exposes a nuclear localization sequence (NLS) which leads to binding of importins and translocation of NFAT into the nucleus. In the nucleus, NFAT acts in a coordinate manner with other transcription factors to activate specific genes required for T cell activation, function and differentiation (Gwack, Feske et al. 2007). NFAT-driven gene expression is highly

dependent on sustained Ca influx and calcineurin activity. A decrease in intracellular Ca levels or treatment with the calcineurin inhibitor cyclosporin A results in the immediate export of NFAT from nucleus by NFAT kinases and alterations in gene expression (Gwack, Feske et al. 2007).

T CELLS AND ION CHANNELS

CRAC channels

In T cells, influx of Ca is regulated through Ca release-activated Ca (CRAC) channels. Activation of the IP₃ receptors on the surface of the ER releases Ca from internal ER stores resulting in the aggregation of the ER transmembrane protein, Stim1. Stim1 is a single transmembrane protein that exists as a monomer when ER Ca stores are full (Manji, Parker et al. 2000). During T cell activation, ER stores are depleted and Ca is released from the EF-hand domain, located in the lumen of the ER. Stim1 dimerization occurs and aggregates, or puncta, form in areas within 10-25µm of the plasma membrane initiating clustering of Orai1 proteins on the plasma membrane and activation of the CRAC channel (Liou, Kim et al. 2005; Wu, Buchanan et al. 2006). Orai1 is a 4 transmembrane spanning protein with both the N- and C-terminus located in the cytoplasm (Yeromin, Zhang et al. 2006). Clustering of 4 Orai1 subunits forms the pore forming unit of CRAC channels (Xu, Lu et al. 2006). The complete CRAC channel complex has not been elucidated but expression of Stim1 and Orai1 are sufficient to produce a CRAC current (I_{CRAC}) (Li, Lu et al. 2007; Salido, Sage et al. 2009). CRAC channels were originally described using electrophysiological methods but the recent identification of Stim1 and Orai1 have allowed for a more detailed characterization.

CRAC channels are characterized by activation through ER store depletion, an extreme selectivity for Ca and a low conductance rate, or the rate of ion travel through the channel (Parekh 2006). T cells are small in size, 5-10 μm , and, at rest, the intracellular Ca concentration is around 50 nM (estimated to be about 10,000 free Ca ions) so a very small, Ca selective current, on the scale of picoamps, is capable of evoking a substantial rise in the concentration of cytosolic Ca (Cahalan and Chandy 2009). The selectivity of Ca through CRAC channels is such that monovalent ions are excluded by ion-pore and ion-ion interactions (McNally, Yamashita et al. 2009). Absence of extracellular divalent ions allows the flow of monovalent ions but addition of very low concentrations of Ca blocks monovalent permeation through high affinity Ca binding to the channels. The presence of one Ca ion is suggested to be sufficient to block monovalent flow through the channel (Prakriya 2009).

CRAC regulation

CRAC channels can be regulated by several mechanisms (DeHaven, Smyth et al. 2007; Hogan and Rao 2007). Ca²⁺-dependent potentiation (CDP) enhances CRAC activity through increases in extracellular Ca. The reversal of this, depotentiation, can decrease CRAC activity when extracellular Ca is removed. The mechanism for this is unknown but the degree of potentiation is dependent on the pore occupancy by divalent ions, whereby ions with higher permeability support greater potentiation. CRAC channels are also regulated through fast inactivation where I_{CRAC} decreases in hyperpolarizing environments. Lastly, it has been proposed that high intracellular Ca surrounding CRAC channels inhibits CRAC activity and thereby produce an inhibitory feedback. The nature of inactivation and the binding sites are unknown. Recently it has

been reported that CRAC channels may be regulated by the refilling of the ER stores and the reversal of Stim1 puncta formation. It is hypothesized that as ER Ca stores are refilled, Stim1 dimers disassemble and monomeric Stim1 can bind Ca through its EF-hand domain leading to a disruption of the Stim1/Orai1 assembly (Smyth, Dehaven et al. 2008; Fahrner, Muik et al. 2009).

CRAC inhibitors

CRAC channels can be differentially regulated by 2-APB (2-aminoethyldiphenyl borate). At low concentrations (1-5 μ M) 2-APB can increase I_{CRAC} 2-5 fold but at higher concentrations (>10 μ M) it strongly inhibits I_{CRAC} (Peppiatt, Collins et al. 2003; Peinelt, Lis et al. 2008). At inhibitory concentrations of 2-APB, reversal of Stim1 puncta can be observed in HEK293 cells over-expressing Stim1 (DeHaven, Smyth et al. 2008; Peinelt, Lis et al. 2008). However, when Orai1 is overexpressed no effect is observed indicating that disruption of Stim1-Orai1 is not the mechanism of action (Navarro-Borelly, Somasundaram et al. 2008). Enhancement of CRAC activity at low 2-APB concentrations has been proposed to be mediated through recruitment of CRAC channels that facilitates the association between Stim1 and Orai1 (Navarro-Borelly, Somasundaram et al. 2008; Wang, Deng et al. 2009). Recently analogs of 2-APB have been reported to interrupt Stim1 puncta formation but the specificity of these analogs are unknown (Goto, Suzuki et al.). In T cells, inhibition of CRAC channels decreases Ca influx and inhibits T cell function, which in humans, leads to a SCID-like syndrome characterized by an inability to activate T cells (Feske, Gwack et al. 2006; Thompson, Mignen et al. 2009).

OTHER ION CHANNELS AND PUMPS

SERCA

T cells also express other ion channels that modulate cytosolic Ca. Sarco/Endoplasmic reticulum Ca-ATPase (SERCA) pumps are located on the ER and function to pump Ca out of the cell. Thapsigargin (Tg) is a commonly used inhibitor that irreversibly inhibits the SERCA pump and is used to activate CRAC channels.

PMCA

Plasma membrane Ca-ATPase (PMCA) pumps are located on the plasma membrane of T cells. At rest, PMCA are autoinhibited and when cytosolic Ca increases pump activity is initially slow. Over tens of seconds the Ca-Cam complex binds the C terminus of PMCA and displaces the inhibitory domain. The PMCA pump plays a role in modulating Ca signaling and also to prevent the accumulation of Ca in the cytosol that can lead to apoptosis.

RyR

Ryanodine receptors (RyR) are located on the ER and are activated by changes in Ca, as well as through binding of cADP ribose and NADP. RyR facilitate the movement of Ca from ER stores to the cytosol and are important in later Ca signaling events.

Na/Ca exchanger

Sodium/calcium (Na/Ca) exchangers are located on the plasma membrane and the mitochondria. They function to prevent high Ca levels in the cytosol by sequestering

it in the mitochondria or exporting it out of the cell. Movement of Ca in one direction occurs with the concomitant movement of Na in the other direction.

Ca uniporter

Ca uniporters are located on the mitochondria and are activated by high Ca levels in the cytosol. They are involved in sequestering Ca inside the mitochondria to prevent accumulation of Ca inside the cell.

Potassium (K⁺) channels

T cells express 3 types of K⁺ channels; K_v, K_{Ca}, and TRPM. K_v and TRPM4 and TRPM5 are activated by the depolarization of the plasma membrane (an increase of positive charge inside the cell). K_v channels function to increase the Ca driving potential by exporting K⁺ ions. TRPM channels have been reported to inhibit the Ca driving force but their function in T cells is controversial.

Non-store-operated cation channels

There have been several reports of other cation channels that are not activated by ER Ca depletion. TRPV6(CaT1) channels are constitutively active, non-selective cation channels on the plasma membrane that are inhibited by increases in cytosolic Ca. TRPM2(LTPC7) channels are also plasma membrane bound and are activated by pyrimidine nucleotides, ADPR (adenosine 5'-diphosphoribose) and NAD (nicotinamide adenine dinucleotide) and produce a non-selective cation channel. TRPM7(MIC) channels are also non-selective cation channels that are activated by intracellular Mg (magnesium) levels.

REFERENCES

- Adachi, M., Tonegawa K, Uekima T (1966). "Studies on selective herbicidal activity of 3',4'-dichloropropionanilide herbicide.I. Its penetration into plants and detoxication." Pesticide and Technique **14**: 19-21.
- Ambrose, A. M., P. S. Larson, et al. (1972). "Toxicologic studies on 3',4'-dichloropropionanilide." Toxicol Appl Pharmacol **23**(4): 650-9.
- Barnett, J. B. and J. Gandy (1989). "Effect of acute propanil exposure on the immune response of C57Bl/6 mice." Fundam Appl Toxicol **12**(4): 757-64.
- Barnett, J. B., J. Gandy, et al. (1992). "Comparison of the immunotoxicity of propanil and its metabolite, 3,4-dichloroaniline, in C57Bl/6 mice." Fundam Appl Toxicol **18**(4): 628-31.
- Barr, V. A., K. M. Bernot, et al. (2008). "Dynamic movement of the calcium sensor STIM1 and the calcium channel Orai1 in activated T-cells: puncta and distal caps." Mol Biol Cell **19**(7): 2802-17.
- Bartha, R. (1971). "Altered propanil biodegradation in temporarily air-dried soil." J Agric Food Chem **19**(2): 394-5.
- Bartha, R. and D. Pramer (1967). "Pesticide transformation to aniline and azo compounds in soil." Science **156**(3782): 1617-8.
- Bauchinger, M., U. Kulka, et al. (1989). "Cytogenetic effects of 3,4-dichloroaniline in human lymphocytes and V79 Chinese hamster cells." Mutat Res **226**(3): 197-202.
- Bauer, E. R., H. H. Meyer, et al. (1998). "Application of an androgen receptor assay for the characterisation of the androgenic or antiandrogenic activity of various phenylurea herbicides and their derivatives." Analyst **123**(12): 2485-7.
- Blyler, G., K. S. Landreth, et al. (1994). "Selective myelotoxicity of propanil." Fundam Appl Toxicol **22**(4): 505-10.
- Bordeleua, L., Rosen, JD., Bartha R (1972). "Herbicide-derived chloroazobenzene residues: Pathway of formation." J. Agr. Food Chem **20**(3): 573-577.
- Brundage, K. M., R. Schafer, et al. (2004). "Altered AP-1 (activating protein-1) activity and c-jun activation in T cells exposed to the amide class herbicide 3,4-dichloropropionanilide (DCPA)." Toxicol Sci **79**(1): 98-105.
- Cahalan, M. D. and K. G. Chandy (2009). "The functional network of ion channels in T lymphocytes." Immunol Rev **231**(1): 59-87.
- Call, D. J., L. T. Brooke, et al. (1983). "Toxicity, bioconcentration, and metabolism of the herbicide propanil (3',4'-dichloropropionanilide) in freshwater fish." Arch Environ Contam Toxicol **12**(2): 175-82.

- Carey, A. E., H. S. Yang, et al. (1980). "Residual concentrations of propanil, TCAB, and other pesticides in rice-growing of soils in the United States, 1972." Pestic Monit J **14**(1): 23-5.
- Carey V. F., D. S. O., Hoagland R. E. and Talbert R. E. (1995). "Resistance Mechanism of Propanil-Resistant Barnyardgrass: 1. Absorption, Translocation, and Site of Action Studies." Pesticide Biochemistry and Physiology **52**(3): 182-189.
- Chang, F. Y., L. W. Smith, et al. (1971). "Insecticide inhibition of herbicide metabolism in leaf tissues." J Agric Food Chem **19**(6): 1183-6.
- Chisaka, H. and P. C. Kearney (1970). "Metabolism of propanil in soils." J Agric Food Chem **18**(5): 854-8.
- Chow, A. Y. and S. D. Murphy (1975). "Propanil (3,4-dichloropropionanilide)-induced methemoglobin formation in relation to its metabolism in vitro." Toxicol Appl Pharmacol **33**(1): 14-20.
- Dahchour, A., G. Bitton, et al. (1986). "Degradation of the herbicide propanil in distilled water." Bull Environ Contam Toxicol **36**(4): 556-62.
- de la Rosa, P., J. Barnett, et al. (2003). "Loss of pre-B and IgM(+) B cells in the bone marrow after exposure to a mixture of herbicides." J Toxicol Environ Health A **66**(24): 2299-313.
- DeHaven, W. I., J. T. Smyth, et al. (2008). "Complex actions of 2-aminoethyldiphenyl borate on store-operated calcium entry." J Biol Chem **283**(28): 19265-73.
- DeHaven, W. I., J. T. Smyth, et al. (2007). "Calcium inhibition and calcium potentiation of Orai1, Orai2, and Orai3 calcium release-activated calcium channels." J Biol Chem **282**(24): 17548-56.
- ECB (2006). "DCA Risk Assessment Report." Institute of Health and Consumer Protection (IHCP).
- Eddleston, M., M. Rajapakshe, et al. (2002). "Severe propanil [N-(3,4-dichlorophenyl) propanamide] pesticide self-poisoning." J Toxicol Clin Toxicol **40**(7): 847-54.
- Fahrner, M., M. Muik, et al. (2009). "Mechanistic view on domains mediating STIM1-Orai coupling." Immunol Rev **231**(1): 99-112.
- Feske, S. (2007). "Calcium signalling in lymphocyte activation and disease." Nat Rev Immunol **7**(9): 690-702.
- Feske, S., Y. Gwack, et al. (2006). "A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function." Nature **441**(7090): 179-85.
- Frost, L. L., Y. X. Neeley, et al. (2001). "Propanil inhibits tumor necrosis factor-alpha production by reducing nuclear levels of the transcription factor nuclear factor-kappaB in the macrophage cell line ic-21." Toxicol Appl Pharmacol **172**(3): 186-93.

- Gaynor, J. J. and C. C. Still (1983). "Subcellular Localization of Rice Leaf Aryl Acylamidase Activity." Plant Physiol **72**(1): 80-85.
- Goto, J., A. Z. Suzuki, et al. "Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca^{2+} entry via STIM proteins." Cell Calcium **47**(1): 1-10.
- Guilhermino, L., A. M. Soares, et al. (1998). "Acute effects of 3,4-dichloroaniline on blood of male Wistar rats." Chemosphere **37**(4): 619-32.
- Gwack, Y., S. Feske, et al. (2007). "Signalling to transcription: store-operated Ca^{2+} entry and NFAT activation in lymphocytes." Cell Calcium **42**(2): 145-56.
- Habte, M. and M. Alexander (1980). "Nitrogen Fixation by Photosynthetic Bacteria in Lowland Rice Culture." Appl Environ Microbiol **39**(2): 342-347.
- Hill, R. H., Jr., Z. J. Rollen, et al. (1981). "Tetrachloroazobenzene in 3,4-dichloroaniline and its herbicidal derivatives: propanil, diuron, linuron, and neburon." Arch Environ Health **36**(1): 11-4.
- Hogan, P. G. and A. Rao (2007). "Dissecting ICRAC, a store-operated calcium current." Trends Biochem Sci **32**(5): 235-45.
- Jain, J., C. Loh, et al. (1995). "Transcriptional regulation of the IL-2 gene." Curr Opin Immunol **7**(3): 333-42.
- Kaya, B., A. Creus, et al. (2000). "Use of the Drosophila wing spot test in the genotoxicity testing of different herbicides." Environ Mol Mutagen **36**(1): 40-6.
- Kimbrough, R. D. (1980). "Human health effects of selected pesticides, chloroaniline derivatives." J Environ Sci Health B **15**(6): 977-92.
- Konstantinou, I. K., A. K. Zarkadis, et al. (2001). "Photodegradation of selected herbicides in various natural waters and soils under environmental conditions." J Environ Qual **30**(1): 121-30.
- Lanzilotta, R. P. and D. Pramer (1970). "Herbicide transformation. I. Studies with whole cells of *Fusarium solani*." Appl Microbiol **19**(2): 301-6.
- Lanzilotta, R. P. and D. Pramer (1970). "Herbicide transformation. II. Studies with an acylamidase of *Fusarium solani*." Appl Microbiol **19**(2): 307-13.
- Li, Z., J. Lu, et al. (2007). "Mapping the interacting domains of STIM1 and Orai1 in Ca^{2+} release-activated Ca^{2+} channel activation." J Biol Chem **282**(40): 29448-56.
- Liou, J., M. L. Kim, et al. (2005). "STIM is a Ca^{2+} sensor essential for Ca^{2+} -store-depletion-triggered Ca^{2+} influx." Curr Biol **15**(13): 1235-41.

- Malerba, I., A. F. Castoldi, et al. (2002). "In vitro myelotoxicity of propanil and 3,4-dichloroaniline on murine and human CFU-E/BFU-E progenitors." Toxicol Sci **69**(2): 433-8.
- Manji, S. S., N. J. Parker, et al. (2000). "STIM1: a novel phosphoprotein located at the cell surface." Biochim Biophys Acta **1481**(1): 147-55.
- Matsunaka, S. (1968). "Propanil hydrolysis: inhibition in rice plants by insecticides." Science **160**(834): 1360-1.
- McCorkle, F. M., J. E. Chambers, et al. (1977). "Acute toxicities of selected herbicides to fingerling channel catfish, *Ictalurus punctatus*." Bull Environ Contam Toxicol **18**(3): 267-70.
- McMillan, D. C., T. P. Bradshaw, et al. (1991). "Role of metabolites in propanil-induced hemolytic anemia." Toxicol Appl Pharmacol **110**(1): 70-8.
- McMillan, D. C., J. P. Freeman, et al. (1990). "Metabolism of the arylamide herbicide propanil. I. Microsomal metabolism and in vitro methemoglobinemia." Toxicol Appl Pharmacol **103**(1): 90-101.
- McMillan, D. C., J. E. Leakey, et al. (1990). "Metabolism of the arylamide herbicide propanil. II. Effects of propanil and its derivatives on hepatic microsomal drug-metabolizing enzymes in the rat." Toxicol Appl Pharmacol **103**(1): 102-12.
- McMillan, D. C., T. A. McRae, et al. (1990). "Propanil-induced methemoglobinemia and hemoglobin binding in the rat." Toxicol Appl Pharmacol **105**(3): 503-7.
- McMillan, D. C., J. G. Shaddock, et al. (1988). "Evaluation of propanil and its N-oxidized derivatives for genotoxicity in the *Salmonella typhimurium* reversion, Chinese hamster ovary/hypoxanthine guanine phosphoribosyl transferase, and rat hepatocyte/DNA repair assays." Fundam Appl Toxicol **11**(3): 429-39.
- McNally, B. A., M. Yamashita, et al. (2009). "Structural determinants of ion permeation in CRAC channels." Proc Natl Acad Sci U S A **106**(52): 22516-21.
- Moore, M. T., J. R. Pierce, et al. (1998). "Responses of non-target aquatic organisms to aqueous propanil exposure." Bull Environ Contam Toxicol **61**(2): 169-74.
- Morse, D. L., E. L. Baker, Jr., et al. (1979). "Propanil-chloracne and methomyl toxicity in workers of a pesticide manufacturing plant." Clin Toxicol **15**(1): 13-21.
- Nagel, R., H. Bresch, et al. (1991). "Effect of 3,4-dichloroaniline on the early life stages of the zebrafish (*Brachydanio rerio*): results of a comparative laboratory study." Ecotoxicol Environ Saf **21**(2): 157-64.

- Navarro-Borelly, L., A. Somasundaram, et al. (2008). "STIM1-Orai1 interactions and Orai1 conformational changes revealed by live-cell FRET microscopy." J Physiol **586**(Pt 22): 5383-401.
- Papadopoulou-Mourkidou, E., D. G. Karpouzas, et al. (2004). "The potential of pesticides to contaminate the groundwater resources of the Axios river basin in Macedonia, Northern Greece. Part I. Monitoring study in the north part of the basin." Sci Total Environ **321**(1-3): 127-46.
- Papadopoulou-Mourkidou, E., D. G. Karpouzas, et al. (2004). "The potential of pesticides to contaminate the groundwater resources of the Axios river basin. Part II. Monitoring study in the south part of the basin." Sci Total Environ **321**(1-3): 147-64.
- Parekh, A. B. (2006). "On the activation mechanism of store-operated calcium channels." Pflugers Arch **453**(3): 303-11.
- Pastorelli, R., Catenacci, G., Guanci, M., Fannelli, R., Valoti, E., Minoia, C., Airoidi, L (1998). "3,4 Dichloroaniline-haemoglobin adducts in humans: preliminary data on agricultural workers exposed to propanil." Biomarkers **3**(3): 227-233.
- Peinelt, C., A. Lis, et al. (2008). "2-Aminoethoxydiphenyl borate directly facilitates and indirectly inhibits STIM1-dependent gating of CRAC channels." J Physiol **586**(13): 3061-73.
- Peppiatt, C. M., T. J. Collins, et al. (2003). "2-Aminoethoxydiphenyl borate (2-APB) antagonises inositol 1,4,5-trisphosphate-induced calcium release, inhibits calcium pumps and has a use-dependent and slowly reversible action on store-operated calcium entry channels." Cell Calcium **34**(1): 97-108.
- Pereira, J. L., S. C. Antunes, et al. (2009). "Toxicity evaluation of three pesticides on non-target aquatic and soil organisms: commercial formulation versus active ingredient." Ecotoxicology **18**(4): 455-63.
- Prakriya, M. (2009). "The molecular physiology of CRAC channels." Immunol Rev **231**(1): 88-98.
- Quest, A. F., S. Ghosh, et al. (1997). "DAG second messengers: molecular switches and growth control." Adv Exp Med Biol **400A**: 297-303.
- Roberts, D. M., R. Heilmair, et al. (2009). "Clinical outcomes and kinetics of propanil following acute self-poisoning: a prospective case series." BMC Clin Pharmacol **9**: 3.
- Rooney, J. W., Y. L. Sun, et al. (1995). "Novel NFAT sites that mediate activation of the interleukin-2 promoter in response to T-cell receptor stimulation." Mol Cell Biol **15**(11): 6299-310.

- Salazar, K. D., P. de la Rosa, et al. (2005). "The polysaccharide antibody response after *Streptococcus pneumoniae* vaccination is differentially enhanced or suppressed by 3,4-dichloropropionanilide and 2,4-dichlorophenoxyacetic acid." Toxicol Sci **87**(1): 123-33.
- Salazar, K. D., M. R. Miller, et al. (2006). "Evidence for a novel endocrine disruptor: the pesticide propanil requires the ovaries and steroid synthesis to enhance humoral immunity." Toxicol Sci **93**(1): 62-74.
- Salazar, K. D., I. V. Ustyugova, et al. (2008). "A review of the immunotoxicity of the pesticide 3,4-dichloropropionanilide." J Toxicol Environ Health B Crit Rev **11**(8): 630-45.
- Salido, G. M., S. O. Sage, et al. (2009). "Biochemical and functional properties of the store-operated Ca²⁺ channels." Cell Signal **21**(4): 457-61.
- SCTEE (2001). "Risk Assessment of 3,4-dichloroaniline." Scientific Committee on Toxicity, Ecotoxicity, and the Environment.
- Sheil, J. M., M. A. Frankenberry, et al. (2006). "Propanil exposure induces delayed but sustained abrogation of cell-mediated immunity through direct interference with cytotoxic T-lymphocyte effectors." Environ Health Perspect **114**(7): 1059-64.
- Singleton, S. D. and S. D. Murphy (1973). "Propanil (3,4-dichloropropionanilide)-induced methemoglobin formation in mice in relation to acylamidase activity." Toxicol Appl Pharmacol **25**(1): 20-9.
- Smith-Garvin, J. E., G. A. Koretzky, et al. (2009). "T cell activation." Annu Rev Immunol **27**: 591-619.
- Smyth, J. T., W. I. Dehaven, et al. (2008). "Ca²⁺-store-dependent and -independent reversal of Stim1 localization and function." J Cell Sci **121**(Pt 6): 762-72.
- Stevens, J., Summer, DD. (1991). Handbook of Pesticide Toxicology. New York, Academic Press.
- Still, C. C. and O. Kuzirian (1967). "Enzyme detoxication of 3',4'-dichloropropionanilide in rice and barnyard grass, a factor in herbicide selectivity." Nature **216**(5117): 799-800.
- Still, G. G. (1968). "Metabolic Fate of 3,4-Dichloropropionanilide in Plants: The Metabolism of the Propionic Acid Moiety." Plant Physiol **43**(4): 543-546.
- Still, G. G. (1968). "Metabolism of 3,4-dichloropropionanilide in plants: the metabolic fate of the 3,4-dichloroaniline moiety." Science **159**(818): 992-3.
- Still, G. G., H. M. Balba, et al. (1981). "Studies on the nature and identity of bound chloroaniline residues in plants." J Agric Food Chem **29**(4): 739-46.
- Surovtseva, E. G. and N. S. Funtikova (1978). "[3,4-dichloroaniline cometabolism by representatives of the genus *Pseudomonas*]." Mikrobiologiya **47**(1): 21-5.

- Thompson, J. L., O. Mignen, et al. (2009). "The Orai1 severe combined immune deficiency mutation and calcium release-activated Ca²⁺ channel function in the heterozygous condition." J Biol Chem **284**(11): 6620-6.
- Turci, R., A. Barisano, et al. (2006). "Determination of dichloroanilines in human urine by gas chromatography/mass spectrometry: validation protocol and establishment of Reference Values in a population group living in central Italy." Rapid Commun Mass Spectrom **20**(17): 2621-5.
- USDA (2006). Agricultural Chemical Use Database. N. A. S. Service.
- USEPA (2003). Reregistration Eligibility Decision for Propanil. U. S. E. P. Agency.
- USEPA (2006). Amendment to Reregistration Eligibility Decision for Propanil. U. S. E. P. Agency.
- Ustyugova, I. V., L. L. Frost, et al. (2007). "3,4-dichloropropionaniline suppresses normal macrophage function." Toxicol Sci **97**(2): 364-74.
- Valentovic, M., J. G. Ball, et al. (2001). "3,4-Dichlorophenylhydroxylamine cytotoxicity in renal cortical slices from Fischer 344 rats." Toxicology **162**(3): 149-56.
- Valentovic, M. A., T. Yahia, et al. (1997). "3,4-Dichloroaniline acute toxicity in male Fischer 344 rats." Toxicology **124**(2): 125-34.
- Wan, F. and M. J. Lenardo (2010). "The nuclear signaling of NF-kappaB: current knowledge, new insights, and future perspectives." Cell Res **20**(1): 24-33.
- Wang, Y., X. Deng, et al. (2009). "STIM protein coupling in the activation of Orai channels." Proc Natl Acad Sci U S A **106**(18): 7391-6.
- Wittke, K., H. Hajimiragha, et al. (2001). "Determination of dichloroanilines in human urine by GC-MS, GC-MS-MS, and GC-ECD as markers of low-level pesticide exposure." J Chromatogr B Biomed Sci Appl **755**(1-2): 215-28.
- Wu, M. M., J. Buchanan, et al. (2006). "Ca²⁺ store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane." J Cell Biol **174**(6): 803-13.
- Xie, Y. C., R. Schafer, et al. (1997). "The immunomodulatory effects of the herbicide propanil on murine macrophage interleukin-6 and tumor necrosis factor-alpha production." Toxicol Appl Pharmacol **145**(1): 184-91.
- Xie, Y. C., R. Schafer, et al. (1997). "Inhibitory effect of 3,4-dichloro-propionaniline on cytokine production by macrophages is associated with LPS-mediated signal transduction." J Leukoc Biol **61**(6): 745-52.
- Xu, P., J. Lu, et al. (2006). "Aggregation of STIM1 underneath the plasma membrane induces clustering of Orai1." Biochem Biophys Res Commun **350**(4): 969-76.

- Yeromin, A. V., S. L. Zhang, et al. (2006). "Molecular identification of the CRAC channel by altered ion selectivity in a mutant of Orai." Nature **443**(7108): 226-9.
- Yih, R. Y., D. H. McRae, et al. (1968). "Mechanism of selective action of 3',4'-dichloropropionanilide." Plant Physiol **43**(8): 1291-6.
- Yih, R. Y., D. H. McRae, et al. (1968). "Metabolism of 3',4'-dichloropropionanilide: 3,4-dichloroaniline-lignin complex in rice plants." Science **161**(839): 376-7.
- Yun, M. S., W. Chen, et al. (2007). "Propanil and swep inhibit 4-coumarate:CoA ligase activity in vitro." Pest Manag Sci **63**(8): 815-20.
- Zablotowicz, R. M., M. A. Locke, et al. (2001). "Fluorescent Pseudomonas isolates from Mississippi Delta oxbow lakes: in vitro herbicide biotransformations." Environ Toxicol **16**(1): 9-19.
- Zhang, B. and S. Lin (2009). "Effects of 3,4-dichloroaniline on testicle enzymes as biological markers in rats." Biomed Environ Sci **22**(1): 40-3.
- Zhang, Z. Y., L. J. Pan, et al. (2009). "Functional interactions among STIM1, Orai1 and TRPC1 on the activation of SOCs in HL-7702 cells." Amino Acids.
- Zhao, W., R. Schafer, et al. (1998). "Cytokine production by C57BL/6 mouse spleen cells is selectively reduced by exposure to propanil." J Toxicol Environ Health A **55**(2): 107-20.
- Zhao, W., R. Schafer, et al. (1999). "Propanil affects transcriptional and posttranscriptional regulation of IL-2 expression in activated EL-4 cells." Toxicol Appl Pharmacol **154**(2): 153-9.
- Zhao, W., R. Schafer, et al. (1995). "Changes in primary and secondary lymphoid organ T-cell subpopulations resulting from acute in vivo exposure to propanil." J Toxicol Environ Health **46**(2): 171-81.

CHAPTER 2

3,4-dichloropropionanilide (DCPA) inhibits T cell activation by altering the intracellular calcium concentration following store depletion.¹

Published in Toxicological Sciences 103(1), 97–107 (2008).

Tricia L. Lewis^{*†}, Kathleen M. Brundage^{*†}, Rodney A. Brundage^{‡,2},
and John B. Barnett^{*†1}.

^{*}Department of Microbiology, Immunology and Cell Biology, [†]Center for Immunopathology and Microbial Pathogenesis, West Virginia University School of Medicine, Morgantown, WV 26506. [‡]National Institute for Occupational Health and Safety, Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, Morgantown, WV 26506

Running Title: DCPA alters calcium homeostasis in T cells

¹Address correspondence to:

John B. Barnett, Ph.D.

Department of Microbiology, Immunology and Cell Biology

West Virginia University School of Medicine

PO Box 9177, Morgantown, WV. 26506-9177

Tel. 304-293-4029; Fax 304-293-7823

E-Mail: jbarnett@hsc.wvu.edu

²Present address: Department of Microbiology, Immunology and Cell Biology, West Virginia University School of Medicine, Morgantown, WV 26506.

Key Words: T cells, signal transduction, calcium, 3,4-dichloropropionanilide, propanil.

ABSTRACT

Stimulation of T cells through the T cell receptor (TCR) results in the activation of a series of signaling pathways that leads to the secretion of IL-2 and cell proliferation. Influx of calcium (Ca^{2+}) from the extracellular environment, following internal Ca^{2+} store depletion, provides the elevated and sustained intracellular calcium concentration ($[\text{Ca}^{2+}]_i$) critical for optimal T cell activation. Our laboratory has documented that exposure to the herbicide 3, 4-dichloropropionanilide (DCPA) inhibits intracellular signaling events that have one or more Ca^{2+} dependent steps. Herein we report that DCPA attenuates the normal elevated and sustained $[\text{Ca}^{2+}]_i$ that follows internal store depletion in the human leukemic T cell line, Jurkat cells, and primary BALB/c mouse T cells. DCPA did not alter the depletion of internal Ca^{2+} stores when stimulated by anti-CD3 or thapsigargin demonstrating that early IP_3 -mediated signaling and depletion of Ca^{2+} stores were unaffected. 2-aminoethyldiphenol borate (2-APB) is known to alter the store-operated Ca^{2+} (SOC) influx that follows Ca^{2+} store depletion. Exposure of Jurkat cells to either DCPA or 50 μM 2-APB attenuated the increase in $[\text{Ca}^{2+}]_i$ following thapsigargin or anti-CD3 induced store depletion in a similar manner. At low concentrations, 2-APB enhances SOC influx but this enhancement is abrogated in the presence of DCPA. This alteration in $[\text{Ca}^{2+}]_i$, when exposed to DCPA, significantly reduces nuclear NFAT levels and IL-2 secretion without altering the plasma membrane polarization profile. Taken together, these data indicate that DCPA inhibits T cell activation by altering Ca^{2+} homeostasis following store depletion.

INTRODUCTION

The herbicide DCPA, commonly referred to as propanil, is widely used and applied several times throughout the growing season. It is applied predominantly on rice fields for control against several broadleaf and grassy weeds but, was recently registered for use on turf. Annual use of DCPA is estimated to be 7 million pounds per year and represents use on 50-70% of all rice crops in the United States. The broad application and heavy use of this herbicide underscores the importance of investigating its immunotoxic effects.

The effects of DCPA on mammalian immune cells have been well documented (Xie, Schafer et al. 1997; Zhao, Schafer et al. 1998; Brundage, Schafer et al. 2004; Salazar, Miller et al. 2006; Sheil, Frankenberry et al. 2006; Corsini, Codeca et al. 2007; Ustyugova, Frost et al. 2007). *In vivo* administration of DCPA to mice results in decreased *ex vivo* cytokine production by macrophages (IL-1 β , IL-6 and TNF- α) and T cells (IL-2 and IFN- γ) (Barnett 1992; Zhao, Schafer et al. 1998). Previous studies using LPS-stimulated macrophages demonstrated a decrease in [Ca²⁺]_i after exposure to DCPA (Xie, Schafer et al. 1997). Using the murine EL-4 T cell line and human Jurkat T cell lines, we have demonstrated that exposure to DCPA decreased IL-2 production and IL-2 mRNA levels (Zhao, Schafer et al. 1999; Brundage, Schafer et al. 2004). In addition, Jurkat T cells exposed to DCPA demonstrate decreased DNA binding ability of the transcription factor, activating protein-1 (AP-1), and decreased c-jun protein (Brundage, Schafer et al. 2004). The production of IL-2, an important early cytokine, requires the coordinate activation of transcription factors, AP-1, NF- κ B, and NFAT, which all depend, to varying degrees, on the [Ca²⁺]_i (Garriety, Chen et al. 1994).

Activation of T cells is initiated through recognition of a peptide presented on the surface of an antigen presenting cell, along with co-stimulatory interactions, which triggers a series of events including phosphorylation of the TCR complex and recruitment of kinases and adapter proteins to the plasma membrane (PM). Subsequent activation of phospholipase- $\text{C}\gamma_1$ (PLC- γ_1) results in the hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP₂) and the production of two second messengers, diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP₃) (Feske, Okamura et al. 2003; Panyi, Varga et al. 2004). DAG remains in the PM and activates proteins such as protein kinase C (PKC). IP₃ is released from the PM and interacts with the IP₃-receptor (IP₃-R) on the surface of the endoplasmic reticulum (ER) resulting in the release of Ca^{2+} from the ER. Depletion of ER Ca^{2+} stores activates SOC channels located on the PM allowing for the influx of Ca^{2+} from the extracellular environment. This results in an increased and sustained $[\text{Ca}^{2+}]_i$ and activation, via calmodulin, of calcineurin which leads to the dephosphorylation of NFAT and its translocation into the nucleus. Once in the nucleus NFAT acts as a transcription factor for the production of key cytokines (Feske, Okamura et al. 2003). Ca^{2+} also plays a role in the activation of the transcription factors NF- κB , and AP-1 (Lewis 2001; Quintana, Griesemer et al. 2005). Cooperative binding of these three transcription factors is required for optimal transcription and production of IL-2, an essential early cytokine required for T cell proliferation and differentiation (Garriety, Chen et al. 1994).

The importance of calcium as a second messenger in T cell activation has been well established (Lewis 2001; Quintana, Griesemer et al. 2005; Feske 2007). Two coupled mechanisms exist to provide the necessary $[\text{Ca}^{2+}]_i$ required for optimal T cell activation.

The first mechanism involves an initial increase in $[Ca^{2+}]_i$ through the IP_3 -mediated depletion of ER Ca^{2+} stores. This transient increase is necessary but not sufficient for optimal T cell activation (Feske, Gwack et al. 2006). Emptying of the ER Ca^{2+} stores is coupled to the activation of Ca^{2+} -release activated Ca^{2+} (CRAC) channels on the PM. Upon ER store depletion STIM-1 (stromal interaction molecule-1) redistributes into puncta on the ER and accumulates close to the PM (Zhang, Yu et al. 2005; Wu, Buchanan et al. 2006; Xu, Lu et al. 2006). Orai1 (also known as CRACM1) has recently been described as an essential pore subunit of the CRAC channel (Prakriya, Feske et al. 2006; Vig, Peinelt et al. 2006). During T cell activation, aggregation of STIM-1 on the ER induces clustering of Orai1 on the PM resulting in an influx of Ca^{2+} through the CRAC channel (Xu, Lu et al. 2006). This increase in $[Ca^{2+}]_i$ through the CRAC channels is essential for the activation of transcription factors necessary for cytokines production. Defects in the CRAC channel have been reported in patients with severe combined immunodeficiency (SCID) syndrome, underscoring the importance of Ca^{2+} in T cell activation (Feske, Draeger et al. 2000; Feske, Giltz et al. 2001; Feske, Prakriya et al. 2005).

Due to the importance of Ca^{2+} homeostasis in T cell activation and function, we investigated the effect of DCPA on $[Ca^{2+}]_i$ in T cells, using primary BALB/c mice T cells and the Jurkat human T cell leukemia cell line. The human Jurkat T cell line, has been used as a model for human T cell signaling for over 2 decades (Abraham and Weiss 2004). When investigating the immunotoxic effects of DCPA, Jurkat cells are a valuable tool that allows elucidation of mechanisms involved in exposure to DCPA. Since our previous research has demonstrated alterations in T cell signaling we wanted to define

more clearly the immunosuppressive role of DCPA on T cells. Our experiments indicate that DCPA inhibits T cell activation by attenuating increases in $[Ca^{2+}]_i$ following the depletion of internal Ca^{2+} stores.

MATERIALS AND METHODS

Cell lines

Experiments were performed using the human T cell leukemia cell line, Jurkat clone E6-1, obtained from the ATCC (American Tissue Culture Collection, Manassas, VA). Jurkat cells were maintained in complete RPMI (Mediatech Inc., Herndon, VA) supplemented with 10% heat inactivated fetal bovine serum (v/v) (FBS) (Hyclone Inc. Logan, UT), 100 units/ml penicillin (BioWhittaker), 100 µg/ml streptomycin (BioWhittaker), 20 mM glutamine (BioWhittaker) and 50 µM 2-mercaptoethanol (Sigma, St. Louis, MO). The cultures were kept at 37°C in 5% CO₂.

Mice

Female BALB/c mice (8-10 weeks old) were purchased from Charles River Labs (Wilmington DE). Mice were housed in the vivarium at West Virginia University Health Sciences Center under the care of a full time veterinarian and professional staff. They were given access to food and water *ad libitum* and allowed to acclimate to the facility for at least 1-week prior to use. Experiments were conducted in accordance with all federal and institutional guidelines for animal use and were approved by the West Virginia University Institutional Animal Care and Use Committee.

Isolation of Mouse T cells

Spleens from BALB/c mice were removed aseptically, pooled and made into a single cell suspension. Red blood cells were lysed using Tris-NH₄Cl and resuspended in sorting buffer (PBS, 0.5% bovine serum albumin and 2 mM EDTA) at 2.5×10^5 cells/ml. An enriched (>90% by flow cytometry) population of T cells was isolated through negative selection using the Pan T cell isolation kit (Miltenyi Biotec, Auburn, CA).

Briefly, spleen cells were incubated at 4°C for 10 min with a cocktail of biotin-conjugated monoclonal antibodies specific for CD14, CD16, CD19, CD36, CD56, CD123, and Glycophorin A, as described in the manufacturer's protocol (Miltenyi Biotec). Next, monoclonal anti-biotin antibody-conjugated magnetic microbeads were added to the sample and incubated for 15 min at 4°C. Cells were washed with sorting buffer and resuspended at 1×10^8 cells in 500 μ l of cold buffer. Cells were then loaded onto autoMACS columns (Miltenyi Biotec) and purified T cells were collected through a negative selection process. Purified T cells were stimulated using anti-CD3 (BD Bioscience) and goat anti-Armenian hamster antibody (Jackson ImmunoResearch, West Grove PA) to crosslink the anti-CD3 as previously described (Kubo, Born et al. 1989).

Fluorescence Measurement of $[Ca^{2+}]_i$

Jurkat cells or splenic mouse T cells were loaded with the calcium-indicator dye fluo-3 AM (Invitrogen, Carlsbad, CA) as previously described (Grynkiewicz, Poenie et al. 1985). Briefly, cells were harvested and resuspended to a concentration of 5×10^6 cells/ml and incubated for 30 min (37°C in 5% CO₂) in complete RPMI media (1.5% FBS, v/v) containing 0.1 μ M fluo-3 AM in the presence of 0.02% pluronic F-127 (Invitrogen) and 2.5 mM probenecid (Sigma, St. Louis, MO). Cells were then washed twice in Ca²⁺ and Mg²⁺-free Hanks Balanced Salt Solution (HBSS) (Mediatech, Inc, Herndon, VA) containing 10 mM HEPES, pH 7.4, 2% FBS and 2.5 mM probenecid, resuspended to a concentration of 1×10^6 cells/ml and incubated 30 min at room temperature. Since the addition of 2% FBS is essential for cell viability the media contains a nominal concentration of Ca²⁺ (2.5 μ M). Samples were kept at room

temperature and protected from light until ready for analysis. 2×10^6 cells were placed in a quartz cuvette and the fluorescence was measured using a PTI QM-2000-4 spectrofluorometer (Photon Technology International (PTI), Birmingham, NJ) with constant stirring. The fluorescence of the fluo-3 dye was measured with excitation at 490 nm and emission at 525 nm. Stock solutions of DCPA (ChemServices, West Chester, PA) and 2-APB (CalBioChem, San Diego, CA) were diluted in absolute ethanol (AAPER Alcohol and Chemical Company, Shelbyville, KY). Vehicle samples were treated with an equivalent concentration (0.1%v/v) of ethanol. The fluorescence was measured and digitized at 1 Hz using the software program Felix 1.42b (PTI). In Jurkat cell, approximately 45 s after starting data collection, 2.0 μ l of DCPA, 2-APB or vehicle was added to the cuvette, followed immediately by mouse anti-human CD3 antibody at a final concentration of 5.0 μ g/ml (UCHT1) (BD Biosciences, San Diego, CA). Splenic mouse T cells were stimulated with anti-CD3 (BD Bioscience) and goat anti-Armenian hamster antibody (Jackson ImmunoResearch, West Grove PA) to crosslink the anti-CD3 as previously described (Kubo, Born et al. 1989). When the fluorescence returned to background levels, CaCl_2 (Fluka, Switzerland) was added (final concentration 2.5 mM) to the media to provide an external source of Ca^{2+} . Addition of ionomycin (final concentration 200 μ M) (Sigma, St. Louis, MO) provided evidence that the cells were loaded evenly and that the dye remained in the cytosol. Cell membranes were lysed with 0.1%, v/v Triton X-100 (Fisher Scientific, Hampton, NH) to measure the maximum fluorescence (F_{max}) parameter for calculation of $[\text{Ca}^{2+}]_i$ to monitor compartmentalization of the dye and ensure the amount of dye was not a limiting factor. Ethylene glycol-bis(β -aminoethyl ether)-N,N,N',N'-tetraacetic acid tetrasodium salt (EGTA) (Sigma) was

added to a final concentration of 50 mM (pH 7.5) to lower the free Ca^{2+} to a nominally Ca^{2+} -free level ($F_{\min}$).

Fluorescence values were converted to $[\text{Ca}^{2+}]_i$ using the following equation:

$$\text{free}[\text{Ca}^{2+}]_i = K_D[(F - F_{\min}) / (F_{\max} - F)],$$

where K_D (360 nM) is the dissociation constant of the Fluo-3/ Ca^{2+} complex, F is the measured fluorescence intensity, $F_{\min}$ is the minimum fluorescence at very low $[\text{Ca}^{2+}]_i$ (fluorescence after the addition of 50 mM EGTA) and $F_{\max}$ is the fluorescence measured at high $[\text{Ca}^{2+}]_i$ (fluorescence after the addition of Triton X-100) (Grynkiewicz, Poenie et al. 1985). The background fluorescence obtained from unloaded cells over a 3 minute time period was subtracted from all data points before $[\text{Ca}^{2+}]_i$ was calculated.

Measurement of IL-2 Production

Jurkat cells were cultured in complete RPMI media or RPMI media without Ca^{2+} at 5×10^5 cells/well in 48-well plates (Costar, Corning, NY) coated with mouse anti-human CD3 antibody (10 $\mu\text{g}/\text{ml}$) (BD Sciences). Cells were treated with 25, 50, 100, or 200 μM DCPA, or vehicle control. DCPA solutions were diluted in absolute ethanol and vehicle control samples were given an equivalent amount of absolute ethanol (0.1% v/v). Cells were also simultaneously stimulated with anti-CD28 antibody (2 $\mu\text{g}/\text{ml}$) (BD PharMingen, San Diego, CA). Cells were incubated at 37°C in 5% CO_2 for 48 h after which supernatants were collected and placed at -20°C. IL-2 production was determined using the sandwich ELISA method and following the manufacturer's protocol (BD PharMingen). All cultures and ELISA analyses were performed in triplicate and the experiment was repeated three times.

Nuclear Extracts

Jurkat cells were cultured to 1×10^6 cells/ml in complete RPMI and stimulated with 10 ng/ml PMA (Sigma-Aldrich) and 1 μ g/ml A23187 (Sigma-Aldrich). Cells were treated with 0.1% v/v ethanol (vehicle control), 100 μ M DCPA or received no treatment. After a 1, 2, 4, and 6 h exposure, cells were harvested and centrifuged at 1200 rpm for 8 min at 4°C. Nuclear extracts were prepared as previously described (Schreiber, Matthias et al. 1989). Briefly, cells were resuspended in 1ml PBS, pelleted, resuspended in cold buffer A (10mM Hepes pH 7.9, 10mM KCl, 0.1mM EDTA (disodium ethylenediamine tetraacetate), 0.1mM EGTA, 1mM DTT (dithiothreitol) and 0.5mM PMSF (phenylmethanesulfonyl fluoride)) and incubated on ice for 15 min. After incubation, 25 μ l of a 10% solution of Nonidet NP-40 was added, vortexed and centrifuged for 30 s at 14,000 rpm. Supernatants, containing the cytoplasmic fraction of the cells, were frozen at -70°C. The nuclear pellet was resuspended in cold buffer C (20 mM Hepes pH 7.9, 0.4 M NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM DTT and 1 mM PMSF) and incubated on ice for 15 min with vortexing every 5 min. Samples were centrifuged for 5 min at 14,000 rpm and the supernatants containing the nuclear fraction were stored at -70°C. The protein concentrations of the nuclear extracts were determined using Coomassie plus protein assay reagent kit following the manufacturer's protocol (Pierce, Rockford, IL).

Western Blots

A 30 μ g aliquot of each nuclear extract was boiled for 5 min to denature the proteins and electrophoresed through an 8% Tris polyacrylamide gel with a 4% stacking gel at 25 mAmps for 18 h. Proteins were transferred onto Hybond-P membranes (Amersham Pharmacia, Piscataway, NJ) at 0.1 amps for 20 h. Blots were washed in TBS for 5 min

at room temperature, blocked for 1 h in TBS + 0.1% Tween 20 (TBS/T) plus 5% dry milk at room temperature and then washed three times in TBS/T. Blots were incubated overnight at 4°C with primary antibodies specific for total NFATc2 (Santa Cruz Santa Cruz, CA) or β actin (Santa Cruz) in TBS/T plus 5% BSA. The next day, blots were washed three times in TBS/T, incubated for 1 h at room temperature with anti-Biotin (Cell Signaling Technology, Inc., Danvers, MA) and either a goat anti-mouse IgG-horseradish peroxidase (HRP) (Santa Cruz) or a rabbit anti-goat IgG-HRP (Sigma-Aldrich). Finally, the blots were washed three times in TBS/T and developed using Phototope-HRP detection kit for western blots (Cell Signaling Technology, Inc) and bands were visualized on X-Ray film (BioMax MR, Eastman Kodak Company). Densitometric analysis was performed using Optimus software (Media Cybernetics, Silver Spring, MD) and nuclear NFAT protein levels were normalized to actin protein levels for each sample.

Measurement of membrane potential

The membrane potential of Jurkat cells was measured using the membrane potential-sensitive DiBAC₄(3) dye (Invitrogen). This bis-oxonol dye produces an excitation maximum at approximately 490 nm. As the cell depolarizes increasing amounts of dye enters the cell where it binds to intracellular proteins or membranes and exhibit enhanced fluorescence. Conversely, hyperpolarization is indicated by a decrease in fluorescence. DiBAC₄(3) is excluded from mitochondria because of its overall negative charge, allowing measurement of the plasma membrane potential (Wolff, Fuks et al. 2003). Membrane potential experiments were carried out in a time course similar to the fluo-3 experiments. Briefly, 1×10^6 cells/ml of Jurkat cells were suspended Ca^{2+} and

Mg²⁺-free Hanks Balanced Salt Solution (HBSS) (Mediatech, Inc.) containing 10 mM HEPES, pH 7.4 and 2% FBS containing 20nM DiBAC₄(3). Samples were kept at room temperature and protected from light until ready for analysis. A 1.0 ml (1 × 10⁶ cells) aliquot of loaded cells was placed in a FACS tube and the fluorescence was measured using a FACSCalibur flow cytometer (Becton Dickinson, Franklin Lakes, NJ). Each recorded time point consisted of 10,000 cells. At t=0 a background fluorescence was recorded after which DCPA, or vehicle control was added, followed immediately by anti-CD3 antibody at a final concentration of 5.0 µg/ml (BD Biosciences). Fluorescence measurements were recorded immediately following treatment and anti-CD3 addition (t=1 min) and again at t=3 min and t=5 min. External Ca²⁺, in the form of CaCl₂, (Fluka, Switzerland) was subsequently added (final concentration 2.5 mM) to the cells to provide an external source of Ca²⁺. Fluorescence measurements were collected at 3 time points following addition of external Ca²⁺ (t= 5.5, 6.5, and 9.5 min). Addition of 40 µl KCl (final concentration 100 mM) completely depolarized the cell and fluorescence values were recorded at t=10, 12 and 14 min. Background fluorescence of unloaded cells was subtracted from all data points before the net change in fluorescence was calculated. The fluorescence was represented as the net change in the fluorescence signal as a percent of the initial background fluorescence (F₀) of the T cells at rest as described by $\Delta F/F_0 = (F_t - F_0) * 100 / F_0$ (where F_t is the fluorescence at each time point) (Wolff, Fuks et al. 2003).

Statistical Analysis

All data were analyzed using MS Excel 2003 (Redmond, WA) and Sigma Stat 3.1 (Port Richmond, CA). The area under the curve (AUC) is an indirect measurement of the

increase of $[Ca^{2+}]_i$ over a selected time period. The AUC includes all data points from the time the external Ca^{2+} was added until the $[Ca^{2+}]_i$ reached a plateau and was calculated using Sigma Stat 3.1. The peak $[Ca^{2+}]_i$ was calculated by determining the highest $[Ca^{2+}]_i$ between addition of $CaCl_2$ and addition of ionomycin. Several curves were used for each sample and the mean $\pm$ S.D. is reported. A t-test was used to determine statistical significance for all Ca^{2+} fluorescence and membrane potential experiments. ANOVA was used to determine statistical significance in IL-2 production. An alpha value of <0.05 was considered significant.

RESULTS

Effect of DCPA on $[Ca^{2+}]_i$ in Anti-CD3 Stimulated Jurkat Cells

T cell activation and proliferation requires a cascade of signaling events mediated by two Ca^{2+} sensitive mechanisms. The first involves the IP_3 -mediated depletion of Ca^{2+} stores in the ER and is coupled to the second mechanism, the activation of CRAC channels and the influx of Ca^{2+} . To determine the effect of DCPA on $[Ca^{2+}]_i$ Jurkat cells, loaded with fluo-3 in a nominally Ca^{2+} -free (2.5 μ M) HBSS solution, were treated with 25, 50, 100, 200 μ M DCPA or vehicle (ethanol) control and stimulated with anti-CD3. We determined that DCPA and ethanol, hereafter referred to as the vehicle, did not interfere with the fluorescence of the fluo-3 dye (data not shown). There was no significant difference between cells stimulated with anti-CD3 or cells that were exposed to vehicle and stimulated with anti-CD3 (data not shown). As shown in Figure 1, addition of DCPA or vehicle followed by stimulation with anti-CD3, in a nominally Ca^{2+} -free environment, resulted in an equivalent small, transient increase in fluorescence. This increase in the $[Ca^{2+}]_i$ represents the depletion of IP_3 -sensitive Ca^{2+} stores and its subsequent removal out of the cell by activated Ca^{2+} -ATPases or its sequestration within the cell (Fig. 1A). DCPA and vehicle treated cells exhibited similar changes in $[Ca^{2+}]_i$ indicating that DCPA does not affect the early IP_3 -mediated signaling pathway or the depletion of internal Ca^{2+} stores.

In T cells, optimal activation of several transcription factors requires a sustained and elevated $[Ca^{2+}]_i$ that is maintained by Ca^{2+} influx through CRAC channels following internal Ca^{2+} store depletion. To determine if DCPA alters $[Ca^{2+}]_i$ following store depletion we added Ca^{2+} to the media following anti-CD3 induced store depletion (Fig. 1A). Figure 1B is an expanded view of the change in $[Ca^{2+}]_i$ following store depletion

and addition of external Ca^{2+} , where the fluorescence intensity, seen in Figure 1A, was converted to an $[\text{Ca}^{2+}]_i$. A decrease in the $[\text{Ca}^{2+}]_i$, following store depletion was observed in Jurkat cells exposed to all four DCPA concentrations, when compared to the vehicle control (Fig. 1A and B). Cells were further treated with Triton X-100 to monitor compartmentalization and ensure that the amount of available dye was not a limiting factor. In addition, Triton X-100 and EGTA provided maximum and minimum fluorescence values, respectively, to calculate the $[\text{Ca}^{2+}]_i$. (See Material and Methods). Similar fluorescent levels after addition of ionomycin, Triton X-100 and EGTA were seen independent of treatment (Fig. 1A).

To further quantify the attenuation of the $[\text{Ca}^{2+}]_i$ following store depletion the area under the curve (AUC), peak $[\text{Ca}^{2+}]_i$, and time to peak, were calculated. The AUC was calculated for each sample to determine the relative amount of free Ca^{2+} in the cytosol over a 103 s time period ($t = 337\text{--}440$ s). The AUC and peak $[\text{Ca}^{2+}]_i$ data for cells exposed to increasing concentrations of DCPA and stimulated with anti-CD3 are shown in Figure 1C and D. Exposure of cells to 200 μM DCPA resulted in a significant decrease (79%) in the AUC and the peak $[\text{Ca}^{2+}]_i$ (78%). A 56% decrease in the AUC and a 52% decrease in the peak $[\text{Ca}^{2+}]_i$ was observed in cells treated with 100 μM . Cells treated with 25 and 50 μM DCPA also resulted in decreases in AUC and peak $[\text{Ca}^{2+}]_i$. Although the decreases observed in the presence of 25 and 50 μM were not statistically significant they highlight the trend in the concentration dependent effect of DCPA. In addition, the time to reach peak $[\text{Ca}^{2+}]_i$ following addition of external Ca^{2+} , increased with increasing concentration of DCPA exposure. Vehicle control cells reached a peak $[\text{Ca}^{2+}]_i$ after approximately 65 s. Treatment of cells with 25, 50, 100,

and 200 μ M DCPA resulted in a peak $[Ca^{2+}]_i$ at approximately 75, 81, 93 and 108 s , respectively. Taken together, these results demonstrate that exposure to DCPA decreases $[Ca^{2+}]_i$ following anti-CD3 induced store depletion in a concentration dependent manner.

Figure 1 . DCPA decreases intracellular calcium in a concentration dependent manner.

Jurkat cells were loaded with fluo-3 and at 50s, anti-CD3 was added simultaneously with 25, 50, 100, 200 μM DCPA or vehicle control and changes in $(\text{Ca}^{2+})_i$ were recorded with a spectrofluorometer. At 335 s, the external $[\text{Ca}^{2+}]$ concentration was increased to 2.5 mM with CaCl_2 . Starting at 442 s, ionomycin, Triton X-100 and EGTA were sequentially added to the cell media at the times indicated. **A**, a representative experiment of the complete fluorescence curve in the absence (solid line) and presence of DCPA (dashed lines) at increasing concentrations.

Figure 1B : DCPA decreases intracellular calcium in a concentration dependent manner.

Fluorescence intensity data from Figure 1A were used to calculate $[Ca^{2+}]_i$ for the time frame of interest ($t=320-442$ s). These time points represent the influx of Ca into the cell.

Figure 1C and 1D . DCPA decreases intracellular calcium in a concentration dependent manner. C, Statistical analysis of the area under the curve (AUC) for DCPA and vehicle control from 3 separate experiments. **D,** statistical analysis of the peak $[\text{Ca}^{2+}]_i$ for DCPA and vehicle control from 3 separate experiments. Error bars reflect the $\pm$ S.D. and asterisks (*) indicates statistically significant results, $p < 0.05$. Statistical analysis was performed using ANOVA.

The Effect of DCPA on Thapsigargin Stimulated Jurkat Cells

To determine if exposure to DCPA affected the early IP_3 -induced signaling pathway, the SERCA pump inhibitor thapsigargin was used. Thapsigargin depletes ER Ca^{2+} stores and prevents refilling, thereby activating Ca^{2+} influx through CRAC channels independent of IP_3 production (Bergling, Dolmetsch et al. 1998). The addition of 100 μM DCPA or vehicle control followed by 2 μM thapsigargin resulted in an equivalent small, transient increase in fluorescence, similar to that observed with anti-CD3 stimulation (Fig 2A). However, the normal increase in $[Ca^{2+}]_i$, following internal Ca^{2+} store depletion, is significantly decreased in DCPA-treated cells, compared to the vehicle control cells. (Fig. 2A and B). Cells treated with thapsigargin and exposed to 100 μM DCPA exhibited a significant decrease in the AUC and peak $[Ca^{2+}]_i$ similar to that observed with anti-CD3 stimulation (Figure 2C and D). During a 70s time period ($t=360-430s$) DCPA exposed cells exhibited an approximately 50% decrease in the AUC and an approximate 55% decrease in the peak $[Ca^{2+}]_i$. This data indicates that early IP_3 -induced signaling events and depletion of ER Ca^{2+} stores were not affected by the exposure of Jurkat cells to DCPA. However, the addition of extracellular Ca^{2+} following store depletion resulted in an overall decrease in the available free cytosolic Ca^{2+} .

Figure 2A. DCPA does not affect early IP3-mediated signaling or depletion of ER Ca^{2+} stores.

Jurkat cells were loaded with f fluo-3 in a nominally Ca^{2+} -free solution. At 50 s, thapsigargin was added simultaneously with 100 μM DCPA or vehicle control and changes in $(\text{Ca}^{2+})_i$ were recorded with a spectrofluorometer. At 375 s, the external $[\text{Ca}^{2+}]$ was raised to 2.5 mM with CaCl_2 . Starting at 430 s, ionomycin, Triton X-100 and EGTA were sequentially added to the cell media at the times indicated. **A**, a representative experiment of the complete fluorescence curve of vehicle control (dashed line) and 100 μM DCPA-treated cells (solid line).

Figure 2B. DCPA does not affect early IP_3 -mediated signaling or depletion of ER Ca^{2+} stores. Fluorescence intensity data from Figure 2A were used to calculate $[Ca^{2+}]_i$ for the time frame of interest ($t=360-430$ s). These time points represent the influx of Ca into the cell.

Figure 2C and 2D. DCPA does not affect early IP₃-mediated signaling or depletion of ER Ca²⁺ stores. **C**, statistical analysis of the AUC for DCPA and vehicle control from 3 separate experiments. **D**, statistical analysis of the peak [Ca²⁺]_i for DCPA and vehicle control from 3 separate experiments. Error bars reflect the ± S.D. and asterisks (*) indicates statistically significant results, p<0.05. Statistical analysis was performed using a t-test.

The Effect of DCPA and 2-APB on $[Ca^{2+}]_i$

To further characterize the effect of DCPA on $[Ca^{2+}]_i$, we compared DCPA-induced inhibition to that seen with 2-APB, a known inhibitor and enhancer of Ca^{2+} influx through CRAC channels (Prakriya and Lewis 2001). At low concentrations ($<5 \mu M$) 2-APB enhances Ca^{2+} influx through CRAC channels but at higher concentrations ($>10 \mu M$) it inhibits Ca^{2+} influx (Prakriya and Lewis 2001). Since 2-APB has been shown to alter Ca^{2+} influx following store depletion we compared the effects of 2-APB on $[Ca^{2+}]_i$ to that of DCPA.

Jurkat cells were stimulated with anti-CD3 and the change in $[Ca^{2+}]_i$ was measured in cells treated with vehicle control, $100 \mu M$ DCPA, $2.5 \mu M$ 2-APB, $50 \mu M$ 2-APB or a mixture of $100 \mu M$ DCPA and $2.5 \mu M$ 2-APB (Fig. 3A). As expected there was no change in the release of Ca^{2+} from internal stores regardless of treatment, except for the cells exposed to $50 \mu M$ 2-APB (data not shown). High concentrations of 2-APB have been reported to affect IP_3 -mediated signaling in addition to inhibiting Ca^{2+} influx but IP_3 signaling is not affected at lower enhancing concentrations of 2-APB (Prakriya and Lewis 2001). Figure 3A shows the changes in $[Ca^{2+}]_i$ following store depletion and subsequent addition of external Ca^{2+} . Cells treated with $2.5 \mu M$ 2-APB demonstrated a significant increase in $[Ca^{2+}]_i$ compared to the vehicle control whereas $50 \mu M$ 2-APB abrogates the increase in the $[Ca^{2+}]_i$ (Fig. 3A). When cells were simultaneously treated with $100 \mu M$ DCPA and $2.5 \mu M$ 2-APB the enhanced effect of 2-APB was abrogated.

Cells treated with DCPA or $50 \mu M$ 2-APB revealed significant decreases in the AUC and the peak $[Ca^{2+}]_i$, as compared to the vehicle control (Fig. 3B and C). The AUC and peak $[Ca^{2+}]_i$ were significantly increased (30%) in cells treated with $2.5 \mu M$ 2-APB (Fig. 3B and C). Cells treated simultaneously with $100 \mu M$ DCPA and $2.5 \mu M$ 2-APB

exhibited a 30% decrease in the AUC and peak $[Ca^{2+}]_i$ when compared to the enhanced influx of 2.5 μ M 2-APB (Fig. 3B and C). Together these results provide further evidence that, following store-operated Ca^{2+} depletion, DCPA-exposed cells decrease $[Ca^{2+}]_i$ in a similar manner as 2-APB and DCPA interferes with the ability of 2-APB to enhance Ca^{2+} influx through CRAC channels.

Figure 3A. The effect of DCPA and 2-APB on $[Ca^{2+}]_i$

Jurkat cells were loaded with fluo-3 in a nominally free Ca^{2+} solution. Anti-CD3 was added simultaneously with 100 μ M DCPA, 2.5 μ M 2-APB, 50 μ M 2-APB, a mixture of 100 μ M DCPA and 2.5 μ M or vehicle control (solid line). Fluorescence changes in $(Ca^{2+})_i$ were recorded with a spectrofluorometer and converted to a $[Ca^{2+}]_i$. **A**, a representative experiment of the conversion of the fluorescence intensity to a $[Ca^{2+}]_i$ from $t=300-420$ s for DCPA and vehicle control.

Figure 3B and 3C. The effect of DCPA and 2-APB on $[Ca^{2+}]_i$. **B**, statistical analysis of the AUC for DCPA, 2-APB (2.5 μ M and 50 μ M), 100 μ M DCPA plus 2.5 μ M 2-APB and vehicle control from 3 separate experiments. **C**, statistical analysis of the peak $[Ca^{2+}]_i$ for DCPA, 2-APB (2.5 μ M and 50 μ M), 100 μ M DCPA plus 2.5 μ M 2-APB and vehicle control from 3 separate experiments. Error bars reflect the $\pm$ S.D. and asterisks (*) indicates statistically significant change compared to vehicle control, $p < 0.05$. Statistical analysis was performed using a t-test.

The Effect of DCPA on Membrane Potential

It is well known that changes in the membrane potential of T cells can alter Ca^{2+} influx (Sarkadi, Tor dai et al . 1990) . To determine if the effect on $[\text{Ca}^{2+}]_i$ seen in DCPA-exposed cells was due to alterations in the membrane potential Jurkat cells were loaded with the membrane potential sensitive dye, DiBAC₄(3). As the cell depolarizes increasing amounts of the DiBAC₄(3) dye enters the cell membrane resulting in an increased fluorescence signal whereas a decrease in the fluorescence signal indicates hyperpolarization. Jurkat cells were exposed to 100 μM DCPA or vehicle control and stimulated with anti-CD3 in a nominally Ca^{2+} -free buffer. Changes in the fluorescence signal using flow cytometry are depicted in Figure 4. The background resting T cell fluorescence was recorded at $t = 0$. Fluorescence measurements were taken immediately following the simultaneous addition of the treatment and anti-CD3 ($t = 1$ min), then again at $t = 3$ min and $t = 5$ min. External Ca^{2+} was then added and fluorescence measurements were taken immediately at $t = 5.5$ min, $t = 6.5$ and $t = 9.5$ min. Addition of 50 mM KCl at $t = 10$ min depolarized the cell and demonstrated that the dye responds to this depolarization. Further fluorescence measurements were also taken at $t = 12$ and 14 min. This data demonstrates that the DCPA-induced attenuation of $[\text{Ca}^{2+}]_i$ following store depletion is not due to changes in the membrane potential.

Figure 4 DCPA does not alter the membrane potential of anti-CD3 stimulated Jurkat cells. Jurkat cells were loaded with DiBAC₄(3) in a nominally Ca²⁺ free solution and analyzed via flow cytometry. Open circles are DCPA-treated cells and closed triangles are vehicle control cells. At t = 0 the background resting potential was recorded. Fluorescence measurements were taken immediately following addition of anti-CD3 and treatment (t = 1 min), then again at t = 3 min and t = 5 min. External Ca²⁺ (2.5mM CaCl₂) was then added and fluorescence measurements were recorded at t = 5.5 min, t = 6.5 and t = 9.5 min. 50 mM KCl was added to depolarize the cells and fluorescence measurements were taken at t = 10, 12 and 14 min. Error bars reflect the $\pm$ S.D. from 3 experiments. Statistical analysis was performed using a t-test.

Effect of DCPA on IL-2 Secretion in Jurkat cells

A reduction in the elevated $[Ca^{2+}]_i$ following store depletion has direct downstream effects on Ca^{2+} -dependent pathways. In order to establish a functional consequence of exposure to DCPA we assessed its effect on IL-2 secretion. IL-2 is an essential early cytokine required for T cell proliferation and differentiation. The activation of transcription factors NF- κ B, NFAT and AP-1, depend, to varying degrees, on the $[Ca^{2+}]_i$ and are important in production of IL-2. Jurkat cells were stimulated with anti-CD3 and anti-CD28 and exposed to 25, 50, 100, or 200 μ M DCPA or vehicle control. After 48 h in culture, IL-2 levels in the supernatant were assayed via a sandwich ELISA. The results of a representative experiment are shown in Table 1. In the presence of 2.5 mM extracellular Ca^{2+} , DCPA decreased IL-2 production in a concentration-dependent manner, with no detectable IL-2 production when exposed to 200 μ M DCPA. Exposure of Jurkat cells to 100 μ M DCPA inhibited IL-2 production by 72%. In experiments performed with the same DCPA and vehicle control concentrations but with nominal extracellular Ca^{2+} (≤ 2.5 μ M), no detectable levels of IL-2 were measured (data not shown).

Table 1. Effects of DCPA on IL-2 production

Treatment	IL-2	% control
No treatment	3420±272	97.4
Vehicle control	3510±564	100.0
25 µM DCPA	2720±169	77.5
50 µM DCPA	2262±312	64.4
100 µM DCPA	968±245	27.6
125 µM DCPA	1171±137	33.4
200 µM DCPA	<32 ^a	

Note: IL-2 is measured in µg/ml ± SD

^abelow the level of detection

Effect of DCPA on Nuclear Translocation of NFAT

In order to establish a direct mechanistic consequence of DCPA on the inhibition of Ca^{2+} -dependent signaling events we examined the Ca^{2+} -dependent transcription factor NFAT. A sustained $[\text{Ca}^{2+}]_i$ via Ca^{2+} influx results in the calmodulin-stimulated activation of the protein phosphatase, calcineurin (Lewis 2001; Feske, Okamura et al. 2003; Quintana, Griesemer et al. 2005). Calcineurin dephosphorylates cytoplasmic NFAT allowing its translocation into the nucleus where it acts as a transcription factor for many genes, including IL-2. To assess the effect of DCPA on NFAT we measured the levels of nuclear NFAT at various time points after T cell stimulation. Jurkat cells were treated with vehicle control, 100 μM DCPA or no treatment and stimulated with PMA and A23187. After 1, 2, 4 or 6 h incubations nuclear extracts were made and the level of nuclear NFAT was determined by western blot. Cells exposed to 100 μM DCPA exhibited decreases in nuclear NFAT at all time points (Fig. 5A and 5B). Densitometric analysis of the decrease in nuclear NFAT in DCPA-treated Jurkat cells indicated that 100 μM DCPA decreased nuclear NFAT approximately 30% compared to the vehicle control (Fig. 5C). Since NFAT is a Ca^{2+} -dependent transcription factor, this data demonstrates a direct mechanism linking the decrease in $[\text{Ca}^{2+}]_i$ observed in DCPA-treated cells to decreased IL-2 production.

Figure 5A. DCPA decreases nuclear NFAT levels. Nuclear extracts from Jurkat cells were stimulated and exposed to 100 μ M DCPA at various time points. **A**, a representative blot indicating the bands that represent different forms of NFAT (top blot). The bands in the bottom blot are β -actin levels used as a loading and transfer control. **B**, the ratio of NFAT protein to β -actin protein using densitometry of the bands in part A.

Figure 5B and 5C . DCPA decreases nuclear NFAT levels. **B**, the ratio of NFAT protein to β -actin protein using densitometry of the bands in part A. **C**, the ratio the ratio of NFAT protein to β -actin protein expressed as a percent of the vehicle control.

Effect of DCPA on Primary Mouse T Cells

In order to establish that the DCPA-induced attenuation of $[Ca^{2+}]_i$ was not unique to the Jurkat cell line, we assayed the effect of DCPA on $[Ca^{2+}]_i$ levels in primary mouse T cells. Splenic T cells were isolated from female BALB/c mice by negative selection and magnetic sorting. The T cells were loaded with fluo-3 and the effect of 100 μ M DCPA on $[Ca^{2+}]_i$ was assayed as described previously. DCPA exposure did not affect the anti-CD3 induced release of Ca^{2+} from internal stores (data not shown). However, there was a significant decrease in $[Ca^{2+}]_i$ following store depletion and addition of external Ca^{2+} (Fig. 6). There were not enough data points to assess the change in peak $[Ca^{2+}]_i$ but the AUC (t=517-616 s) was significantly decreased (16%) in the DCPA treated cells (Fig. 6B). Altogether, these results demonstrate that the effect of DCPA on Ca^{2+} influx following store depletion is not limited to the human Jurkat T cells.

Figure 6. DCPA decreases $[Ca^{2+}]_i$ in primary mouse T cells. Purified mouse splenic T cells were loaded with fluo-3. Anti-CD3 was added simultaneously with 100 μ M DCPA or vehicle control and Ca^{2+} fluorescence measured. At 567 s, the external $[Ca^{2+}]$ was raised to 2.5 mM with $CaCl_2$. **A**, a representative experiment showing $[Ca^{2+}]_i$ in the presence of vehicle control (solid line) or 100 μ M DCPA (dashed line). **B**, statistical analysis of the AUC. Error bars reflect the $\pm$ S.D. from 2 experiments and analysis was performed using a student t-test, $p < 0.05$.

DISCUSSION

DCPA is a widely used herbicide that is heavily used on rice crops. Its recent registration for use on turf will further increase not only its use but will increase the likelihood of human exposure to DCPA. Currently, workers involved in manufacturing, handling or applying DCPA are at greatest risk of exposure but there have also been reports of non-occupational exposure, including a child under the age of 6 (EPA 2006). The United States Environmental Protection Agency (EPA) conducted a risk assessment for workers involved in mixing, loading and application of DCPA. The results from this assessment indicate that workers using maximal protection, including closed mixing and loading systems and enclosed cockpit, cabs or trucks, are at risk for exposure to DCPA (EPA 2006). In addition, workers entering treated areas 12 hours after application were also at risk (EPA 2006). These conclusions emphasize the need to investigate the effects of human exposure to DCPA.

The effects of DCPA in the rodent model have been well established. In a mouse model, exposure to DCPA results in thymic atrophy, depressed NK and macrophage functions, altered cytotoxic T cell response, and decreased CD4⁺ T cells (Barnett and Gandy 1989; Zhao, Schafer et al. 1995; Sheil, Frankenberry et al. 2006; Ustyugova, Frost et al. 2007). The effect of DCPA on human immune cells has been limited. Human macrophages exposed to DCPA produce reduced levels of TNF- α , reactive oxygen species and reactive nitrogen species (Ustyugova, Frost et al. 2007). Human T cells exposed to DCPA exhibit decreased IL-2 production and decreases binding of transcription factor AP-1 to DNA (Brundage, Schafer et al. 2004). Recently Corsini *et al* examined agricultural workers exposed to DCPA and reported alterations in leukocyte cytokine production (Corsini, Codeca et al. 2007).

Since the sustained and elevated influx of Ca^{2+} , following store depletion, is necessary for T cell activation and proliferation, we examined the effects of DCPA on $[\text{Ca}^{2+}]_i$. Our experiments demonstrate that DCPA attenuates the increased $[\text{Ca}^{2+}]_i$ following store depletion in a concentration dependent manner. Anti-CD3 or thapsigargin induced store depletion in DCPA-exposed Jurkat cells did not affect IP_3 -mediated release of Ca^{2+} nor the depletion of ER Ca^{2+} stores. However, our experiments demonstrate that DCPA-exposed Jurkat T cells and mouse splenic T cells were able to attenuate the normal elevated $[\text{Ca}^{2+}]_i$ observed following store depletion. As a consequence of this attenuation nuclear NFAT levels and IL-2 production were also decreased.

2-APB is widely used as both an inhibitor and enhancer of Ca^{2+} influx through CRAC channels (Prakriya and Lewis 2001; Prakriya and Lewis 2006). Low concentrations of 2-APB potentiate Ca^{2+} influx through CRAC channels following store depletion but do not affect IP_3 -mediated release of Ca^{2+} from the ER (Prakriya and Lewis 2001). We used 2-APB to further elicit the inhibitory role of DCPA in Ca^{2+} signaling. At low concentrations of 2-APB ($<5.0 \mu\text{M}$) Ca^{2+} influx is enhanced through CRAC channels following store depletion. Jurkat cells exposed to a mixture of $100 \mu\text{M}$ DCPA and $2.5 \mu\text{M}$ 2-APB abrogated the enhanced Ca^{2+} influx seen with $2.5 \mu\text{M}$ 2-APB. This data suggests that T cells exposed to DCPA prevent the increased influx of Ca^{2+} seen at low concentrations of 2-APB by inhibiting events surrounding CRAC channel activation.

Both voltage-gated and Ca^{2+} -activated potassium (K^+) channels are PM channels involved in regulating the polarization state of T cells and modulating the rate of Ca^{2+}

influx through CRAC channels (Lewis 2001; Panyi, Varga et al. 2004; Quintana, Griesemer et al. 2005). Alterations in the K^+ channel can change the polarization state and result in alterations of Ca^{2+} influx (Panyi, Varga et al. 2004). Depolarization of the cell can inhibit Ca^{2+} influx whereas hyperpolarization increases the driving force of Ca^{2+} into the cell. Our experiments indicate that DCPA does not alter the membrane potential of T cells (Fig.4) and therefore DCPA is unlikely to affect these channels.

In T cells, the plasma membrane Ca^{2+} -ATPase (PMCA) pump is considered the primary Ca^{2+} extrusion mechanism. Modulation of PMCA activity can occur in response to an influx of Ca^{2+} through CRAC channels and results in long term stability of the Ca^{2+} signal (Feske, Prakriya et al. 2005). Since we have demonstrated that DCPA-treated cells attenuate the increase in $[Ca^{2+}]_i$ following store depletion it may be possible that DCPA enhances PMCA activity and thereby reduces the apparent amount of cytosolic Ca^{2+} . However, this mechanism appears unlikely since PMCA activity would also result in a decrease $[Ca^{2+}]_i$ during the initial IP_3 -mediated release of Ca^{2+} from internal stores. Figures 1A, 2A and 3A demonstrate no change in $[Ca^{2+}]_i$ during the initial IP_3 -mediated release of Ca^{2+} from internal stores in DCPA-treated cells.

The mitochondria also play an important role in Ca^{2+} homeostasis (Parekh 2003; Quintana, Griesemer et al. 2005). Mitochondria can act as a Ca^{2+} sink and sequester large amounts of Ca^{2+} quickly and release it slowly after Ca^{2+} influx subsides (Parekh 2003). Expression of a uniporter in the mitochondrial inner membrane and a Na^+/Ca^{2+} exchanger allow for Ca^{2+} uptake and release, respectively (Parekh 2003). Alterations in the activity of the Na^+/Ca^{2+} exchanger or altered membrane potential of the

mitochondrial membrane could result in an apparent decrease in Ca^{2+} influx. Although we have not investigated the effects of DCPA on the mitochondria any increased function of Ca^{2+} uptake by the mitochondria would also result in an apparent decrease the initial IP_3 -mediated release of Ca^{2+} from internal stores, which was not observed in DCPA-exposed T cells.

Our data indicates that DCPA alters Ca^{2+} homeostasis in human and murine T cells. Specifically, DCPA is able to attenuate the increase in $[\text{Ca}^{2+}]_i$ following store depletion. The ability of DCPA to abrogate the enhanced Ca^{2+} influx produced by $2.5 \mu\text{M}$ 2-APB through CRAC channels provides some evidence that DCPA is acting on mechanisms involved in the activation or regulation of CRAC channels. This decrease in $[\text{Ca}^{2+}]_i$ has significant downstream consequences on nuclear NFAT levels and secretion of IL-2. Although the exact mechanism by which DCPA exerts its effect is unknown it is clear that the decrease in $[\text{Ca}^{2+}]_i$ results in a functional consequence to human T cells. The data presented here indicates that exposure to DCPA alters $[\text{Ca}^{2+}]_i$ in murine and human Jurkat T cells and may result in immunosuppression resulting in serious consequences on human health.

Funding

This work was supported by the National Institutes of Health (ES11311); National Institute for Environmental Health Sciences (ES010953); Environmental Protection Agency Greater Research Opportunity Fellowship (MA-91684801-0); Core Grant (RR016440).

Acknowledgements

The authors would like to thank Dr. William Atchison for his helpful advice on this project and Dr. Cynthia Cunningham for her help in the flow cytometry facility.

REFERENCES

- Abraham, R. T. and A. Weiss (2004). "Jurkat T cells and development of the T-cell receptor signalling paradigm." Nat Rev Immunol **4**(4): 301-8.
- Barnett, J., Gandy, J., Wilbourn D., Theus, SA. (1992). "Comparison of the immunotoxicity of propanil and its metabolite, 3,4-dichloroaniline, in C57Bl/6 mice." Fund Appl Toxicol **18**: 626-631.
- Barnett, J. B. and J. Gandy (1989). "Effect of acute propanil exposure on the immune response of C57Bl/6 mice." Fundam Appl Toxicol **12**(4): 757-64.
- Bergling, S., R. Dolmetsch, et al. (1998). "A fluorometric method for estimating the calcium content of internal stores." Cell Calcium **23**(4): 251-9.
- Brundage, K. M., R. Schafer, et al. (2004). "Altered AP-1 (activating protein-1) activity and c-jun activation in T cells exposed to the amide class herbicide 3,4-dichloropropionanilide (DCPA)." Toxicol Sci **79**(1): 98-105.
- Corsini, E., I. Codeca, et al. (2007). "Immunomodulatory effects of the herbicide propanil on cytokine production in humans: In vivo and in vitro exposure." Toxicol Appl Pharmacol **222**(2): 202-10.
- EPA (2006). "Amendment to Reregistration Eligibility Decision (RED) for Propanil."
- Feske, S. (2007). "Calcium signalling in lymphocyte activation and disease." Nat Rev Immunol **7**(9): 690-702.
- Feske, S., R. Draeger, et al. (2000). "Impaired NFAT regulation and its role in a severe combined immunodeficiency." Immunobiology **202**(2): 134-50.
- Feske, S., J. Giltzane, et al. (2001). "Gene regulation mediated by calcium signals in T lymphocytes." Nat Immunol **2**(4): 316-24.
- Feske, S., Y. Gwack, et al. (2006). "A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function." Nature **441**(7090): 179-85.
- Feske, S., H. Okamura, et al. (2003). "Ca²⁺/calcineurin signalling in cells of the immune system." Biochem Biophys Res Commun **311**(4): 1117-32.
- Feske, S., M. Prakriya, et al. (2005). "A severe defect in CRAC Ca²⁺ channel activation and altered K⁺ channel gating in T cells from immunodeficient patients." J Exp Med **202**(5): 651-62.

- Garritty, P. A., D. Chen, et al. (1994). "Interleukin-2 transcription is regulated in vivo at the level of coordinated binding of both constitutive and regulated factors." Mol Cell Biol **14**(3): 2159-69.
- Grynkiewicz, G., M. Poenie, et al. (1985). "A new generation of Ca²⁺ indicators with greatly improved fluorescence properties." J Biol Chem **260**(6): 3440-50.
- Kubo, R. T., W. Born, et al. (1989). "Characterization of a monoclonal antibody which detects all murine alpha beta T cell receptors." J Immunol **142**(8): 2736-42.
- Lewis, R. S. (2001). "Calcium signaling mechanisms in T lymphocytes." Annu Rev Immunol **19**: 497-521.
- Panyi, G., Z. Varga, et al. (2004). "Ion channels and lymphocyte activation." Immunol Lett **92**(1-2): 55-66.
- Parekh, A. B. (2003). "Store-operated Ca²⁺ entry: dynamic interplay between endoplasmic reticulum, mitochondria and plasma membrane." J Physiol **547**(Pt 2): 333-48.
- Prakriya, M., S. Feske, et al. (2006). "Orai1 is an essential pore subunit of the CRAC channel." Nature **443**(7108): 230-3.
- Prakriya, M. and R. S. Lewis (2001). "Potentiation and inhibition of Ca(2+) release-activated Ca(2+) channels by 2-aminoethyldiphenyl borate (2-APB) occurs independently of IP(3) receptors." J Physiol **536**(Pt 1): 3-19.
- Prakriya, M. and R. S. Lewis (2006). "Regulation of CRAC channel activity by recruitment of silent channels to a high open-probability gating mode." J Gen Physiol **128**(3): 373-86.
- Quintana, A., D. Griesemer, et al. (2005). "Calcium-dependent activation of T-lymphocytes." Pflugers Arch **450**(1): 1-12.
- Salazar, K. D., M. R. Miller, et al. (2006). "Evidence for a novel endocrine disruptor: the pesticide propanil requires the ovaries and steroid synthesis to enhance humoral immunity." Toxicol Sci **93**(1): 62-74.
- Sarkadi, B., A. Tordai, et al. (1990). "Membrane depolarization selectively inhibits receptor-operated calcium channels in human T (Jurkat) lymphoblasts." Biochim Biophys Acta **1027**(2): 130-40.

- Schreiber, E., P. Matthias, et al. (1989). "Rapid detection of octamer binding proteins with 'mini-extracts', prepared from a small number of cells." Nucleic Acids Res **17**(15): 6419.
- Schwab, C. L., R. Fan, et al. (2005). "Modeling and predicting stress-induced immunosuppression in mice using blood parameters." Toxicol Sci **83**(1): 101-13.
- Sheil, J. M., M. A. Frankenberry, et al. (2006). "Propanil exposure induces delayed but sustained abrogation of cell-mediated immunity through direct interference with cytotoxic T-lymphocyte effectors." Environ Health Perspect **114**(7): 1059-64.
- Ustyugova, I. V., L. L. Frost, et al. (2007). "3,4-dichloropropionaniline suppresses normal macrophage function." Toxicol Sci **97**(2): 364-74.
- Vig, M., C. Peinelt, et al. (2006). "CRACM1 is a plasma membrane protein essential for store-operated Ca²⁺ entry." Science **312**(5777): 1220-3.
- Wolff, C., B. Fuks, et al. (2003). "Comparative study of membrane potential-sensitive fluorescent probes and their use in ion channel screening assays." J Biomol Screen **8**(5): 533-43.
- Wu, M. M., J. Buchanan, et al. (2006). "Ca²⁺ store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane." J Cell Biol **174**(6): 803-13.
- Xie, Y. C., R. Schafer, et al. (1997). "Inhibitory effect of 3,4-dichloro-propionaniline on cytokine production by macrophages is associated with LPS-mediated signal transduction." J Leukoc Biol **61**(6): 745-52.
- Xu, P., J. Lu, et al. (2006). "Aggregation of STIM1 underneath the plasma membrane induces clustering of Orai1." Biochem Biophys Res Commun **350**(4): 969-76.
- Zhang, S. L., Y. Yu, et al. (2005). "STIM1 is a Ca²⁺ sensor that activates CRAC channels and migrates from the Ca²⁺ store to the plasma membrane." Nature **437**(7060): 902-5.
- Zhao, W., R. Schafer, et al. (1998). "Cytokine production by C57BL/6 mouse spleen cells is selectively reduced by exposure to propanil." J Toxicol Environ Health A **55**(2): 107-20.

- Zhao, W., R. Schafer, et al. (1999). "Propanil affects transcriptional and posttranscriptional regulation of IL-2 expression in activated EL-4 cells." Toxicol Appl Pharmacol **154**(2): 153-9.
- Zhao, W., R. Schafer, et al. (1995). "Changes in primary and secondary lymphoid organ T-cell subpopulations resulting from acute in vivo exposure to propanil." J Toxicol Environ Health **46**(2): 171-81.

CHAPTER 3

Differential effects of T cell function by metabolites of the herbicide, propanil.

Tricia L. Lewis^{*} and John B. Barnett^{*1}

^{*}Department of Microbiology, Immunology and Cell Biology, West Virginia University
School of Medicine, Morgantown, WV 26506

Running Title: Differential effects of propanil metabolites.

¹Address correspondence to:

John B. Barnett, Ph.D.

Department of Microbiology, Immunology and Cell Biology

West Virginia University School of Medicine

PO Box 9177, Morgantown, WV 26506-9177

Tel. 304-293-4029; Fax 304-293-7823

E-Mail: jbarnett@hsc.wvu.edu

Key Words: T cells, 3,4-dichloropropionanilide, DCPA, propanil, metabolites, 3,4-dichloroaniline, N-hydroxy-3,4-dichloroaniline, DCA, T cells, fluorine substitution

ABSTRACT

Each year 2 billion pounds of herbicide are used worldwide to control the unwanted growth of plants. In the United States, over 0.5 billion pounds are used, representing 28% of worldwide use. Propanil (chemical name 3,4-dichloropropionanilide, DCPA) is the 17th most commonly used herbicide in the United States and 6-9 million pounds are applied annually to 2 million acres of rice fields. The immunomodulatory effects of DCPA have been well documented but no data is available on the effects of its metabolites. In mammals, hepatic enzymes metabolize DCPA into 3 major metabolites; 3,4-dichloroaniline (DCA), 6-hydroxy-3,4-dichloroaniline (6OH-DCA), and N-hydroxy-3,4-dichloroaniline (NOH-DCA). We report for the first time the immunotoxic effects of propanil metabolites on T cell function. Jurkat T cells were exposed to varying concentrations of DCPA and its metabolites and assayed for viability, IL-2 secretion, NFAT activity and calcium flux. In addition, the fluorine analogs of DCPA and DCA were investigated to determine the relative role of chlorine substituents on T cell immunotoxicity. As previously reported, Jurkat T cells exposed to DCPA decreased IL-2 secretion in a concentration and calcium-dependent manner. Here we report that exposure of Jurkat T cells to DCA also alters IL-2 secretion, NFAT activity and calcium influx, similar to DCPA. Interestingly, exposure to 6OH-DCA and NOH-DCA reduces IL-2 secretion and NFAT activity but has no effect on calcium influx. When both chlorines in DCPA and DCA were substituted with fluorines all effects were abrogated. Our data indicates that metabolites of DCPA have differential effects on T cell function and the presence of chlorines plays a critical role in eliciting these effects.

INTRODUCTION

Approximately 5.3 billion pounds of pesticides are applied annually across the United States and 15 of the top 25 most used pesticides are herbicides (Grube 2004). Propanil (chemical name 3,4-dichloropropionanilide, DCPA) is a post-emergent contact herbicide and is the 17th most used herbicide in the United States (USEPA 2006). Annual use of DCPA is estimated to be 6-9 million pounds per year and is distributed over 2 million acres of crop land, primarily rice fields. Rice plants are able to avoid the herbicidal effects of DCPA because they produce acylamidase, an enzyme that cleaves the amide bond, resulting in the production of 2 metabolites, 3,4-dichloroaniline (DCA) and propionic acid (Still 1968). In mammals, DCPA can also be metabolized to DCA and propionic acid via acylamidases in the liver (McMillan, Leahey et al. 1990). DCA can undergo further biotransformation in the liver to produce the oxidative metabolites, N-hydroxy-3,4-dichloroaniline (NOH-DCA) and 6-hydroxy-3,4-dichloroaniline (6OH-DCA) (Fig.A). Propionic acid is non-toxic and converted to CO₂ and water but limited data is available on the effects of DCA and its oxidative metabolites.

Workers involved in manufacturing, handling or application of DCPA are at greatest risk for exposure but there have also been reports of non-occupational exposure (Pastorelli 1998). Numerous *in vitro* studies have reported the immunomodulatory effects of exposure to DCPA (reviewed in (Salazar, Ustyugova et al. 2008)). Exposure to DCPA alters macrophage phagocytic activity and cytokine production as well as the lytic function of CD8⁺ T cells after secondary stimulation (Sheil, Frankenberry et al. 2006; Ustyugova, Frost et al. 2007; Salazar, Ustyugova et al. 2008). Other immunotoxic effects include decreased natural killer (NK) cell function, and an increase in antibody secreting cells (Barnett and Gandy 1989; Salazar, de la

Rosa et al. 2005). Recent studies from our lab demonstrate that exposure of Jurkat T cells to DCPA results in a concentration dependent decrease in IL-2 production that is mediated by alterations in NFAT translocation and calcium homeostasis (Lewis, Brundage et al. 2008).

Limited data is available on the toxic effects of DCA, NOH-DCA and 6OH-DCA. It has been reported that DCA and NOH-DCA are toxic to the bladder, liver and kidney of rats (Valentovic, Yahia et al. 1997). DCA can also alter the male reproductive system, binds weakly to the androgen receptor and may act as an endocrine disruptor (Bauer, Meyer et al. 1998; Zhang and Lin 2009). Mice exposed to molar equivalent concentrations of DCPA or DCA have decreased natural killer (NK) cell activity (Barnett and Gandy 1989). However, some studies suggest that DCPA may be more toxic than DCA (Zhao, Schafer et al. 1995; Malerba, Castoldi et al. 2002). Corsini et al reported that workers exposed to DCPA had several alterations in immune parameters (Corsini, Codeca et al. 2007). Agricultural workers exposed to DCPA had increased plasma IgG₁ and IL-6 production in whole blood assays and decreased IL-10 and interferon- γ (IFN- γ) when compared to control subjects. *In vitro* assays demonstrated similar results and at molar equivalent concentrations, both DCPA and DCA inhibited IL-10 and IFN- γ production in enriched CD4⁺CD8⁺ samples. In addition, alterations in calcium homeostasis and cytokine production were observed in anti-CD3 and PHA-stimulated peripheral blood mononuclear cells, suggesting that human T cells may be sensitive to the effects of DCPA.

Using Jurkat T cells we investigated the effects of DCA, NOH-DCA, and 6OH-DCA and the role of halogen substitution on T cell function (Fig.1A and 1B). Chlorine-

containing pesticides are a common class of pesticides and 4–5% of all pesticides introduced into the market after 1989 contain a chlorine-carbon bond (Jeschke 2004). The addition of chlorine to many chemical compounds can increase its activity with negative biological consequences. DDT is a well known organochlorine pesticide whose effects are mediated through the presence and positioning of chlorines (Crinnion 2009). Although alterations in calcium-dependent signaling events have been reported in T cells exposed to DCPA, the effects of the metabolites produced in humans are unknown. We report that exposure to DCA alters IL-2 secretion in a calcium-dependent manner but at higher concentrations than that observed with DCPA. In addition, NOH-DCA and 6OH-DCA appear more toxic to T cells than DCPA and DCA and inhibit IL-2 secretion in a calcium-independent manner. Finally, the substitution of fluorines for chlorines results in an abrogation of all investigated effects in Jurkat T cells. This is the first reported study on the effects of DCPA metabolites on T cells and advances our knowledge on the metabolic and structural effects of DCPA.

Figure 1A DCPA metabolic pathway and structures and in mammals

Figure1B Stucture of fluorine analogs

MATERIALS AND METHODS

Cell lines and reagents

Experiments were performed using the human T cell leukemia cell line, Jurkat clone E6-1, obtained from the ATCC (American Tissue Culture Collection, Manassas, VA). Jurkat cells were maintained in complete Roswell Park Memorial Institute (RPMI 1640) (Mediatech Inc., Herndon, VA) supplemented with 10% heat inactivated fetal bovine serum (v/v) (FBS) (Hyclone Inc. Logan, UT), 100 units/ml penicillin (BioWhittaker, Walkersville, MD), 100 µg/ml streptomycin (BioWhittaker), 20 mM glutamine (BioWhittaker) and 50 µM 2-mercaptoethanol (Sigma, St. Louis, MO). The cultures were kept at 37°C in 5% CO₂.

Stock solutions of 3,4-dichloropropionanilide (DCPA) (ChemServices, West Chester, PA), 3,4-dichloroaniline (DCA) (Chem Services), 3,4-difluoropropionanilide (DFPA), and 3,4-difluoroaniline (DFA) (Sigma) were diluted in absolute ethanol (AAPER Alcohol and Chemical Company, Shelbyville, KY). Vehicle samples were treated with an equivalent concentration (0.1% v/v) of ethanol. Stock solutions of N-hydroxy-3,4-dichloroaniline (NOH-DCA) and 6-hydroxy-3,4-dichloroaniline (6OH-DCA) (a generous gift from G. Rankin) were diluted in dimethyl sulphoxide (DMSO) (Sigma) and vehicle samples were treated with an equivalent concentration (0.1% v/v) of DMSO.

Synthesis of DFPA and NOH-DCA

3,4-Difluoropropionanilide (DFPA) was synthesized from 3,4-difluoroaniline (DFA) (Sigma-Aldrich, Milwaukee, WI). Two grams of propanoic acid (Sigma-Aldrich) was added to 1g of DFA and heated to 100°C for 1 h followed by addition of 5 ml of water and continued heating at 100 °C for an additional hour. The precipitate was

cooled to room temperature, filtered through a sintered glass funnel, washed with water and dried *in vacuo* resulting in 1.26 g of DFPA. The crude DFPA was then re-crystallized from a 1:1 water and ethanol solution.

N-hydroxy-3,4-dichloroaniline (NOH-DCA) was synthesized by methods described by Lerman (Lerman, Weinstock-Rosin et al. 2004). In the presence of nitric acid, dichloromethane and tetra-n-butylammonium bromide (TBAB), 3,4-dichlorophenol was converted to 4,5-dichloro-2-nitrophenol, which was analyzed using NMR and mass spectroscopy and was found to be in agreement with others (Lerman, Weinstock-Rosin et al. 2004). Ethanol (100%) was added to 0.32 g 4,5-dichloro-2-nitrophenol and 20 mg platinum dioxide and hydrogenated (30-50 psi) on a Parr shaker for 1 h. The mixture was filtered through celite and concentrated *in vacuo* to yield NOH-DCA.

Production and purity of DFPA, NOH-DCA and the intermediate, 4,5-dichloro-2-nitrophenol were verified using NMR spectra from a Varian Unity-300 NMR spectrometer (Palo Alto, CA) and exact mass data was obtained using a Thermo-Fisher LTQ-FTICR and were in agreement with previously reported analytical data (Lok, Leone et al. 1996; Lerman, Weinstock-Rosin et al. 2004).

Viability assays

Viability assays were performed for DCPA, DCA, NOH-DCA, 6OH-DCA, DFPA, DFA using 7-AAD (7-amino-actinomycin) (BD Pharmingen, San Diego, CA) and following the manufacturer's protocol. Briefly, 1.0×10^6 Jurkat cells were treated with or without varying concentrations of DCPA, DCA, NOH-DCA, 6OH-DCA, DFPA, DFA, and including ethanol and DMSO vehicle controls and incubated at 37°C in 5% CO₂ for 24 hours. Cells were then incubated in PBS with 5 µl (0.25 µg) 7-AAD and incubated on ice

for 20 min in the dark. Cells were then washed and resuspended in 0.4% paraformaldehyde and analyzed by flow cytometry. Emission was detected in the FL-3 channel ($>650\text{nm}$) using a FACSCalibur flow cytometer (Becton Dickinson, Franklin Lakes, NJ).

Calcium Fluorescence Measurements

Jurkat cells were loaded with the calcium-indicator dye fluo-3 AM (Invitrogen, Carlsbad, CA) as previously described (Grynkiewicz, Poenie et al. 1985). Briefly, cells were harvested and re-suspended to a concentration of 5×10^6 cells/ml and incubated for 30 min (37°C in 5% CO_2) in complete RPMI media (1.5% FBS, v/v) containing $0.1 \mu\text{M}$ fluo-3 AM in the presence of 0.02% pluronic F-127 (Invitrogen) and 2.5 mM probenecid (Sigma). Cells were washed twice in Ca^{2+} and Mg^{2+} -free Hanks Balanced Salt Solution (HBSS) (Mediatech Inc.) containing 10 mM N-(2-hydroxyethyl)-piperazine-N'-2-ethanesulfonic acid (HEPES), pH 7.4, 2% FBS and 2.5 mM probenecid, resuspended to a concentration of 1×10^6 cells/ml and incubated 30 min at room temperature. Since the addition of 2% FBS is essential for cell viability, the media contains a nominal concentration of Ca^{2+} ($2.5 \mu\text{M}$). Samples were kept at room temperature and protected from light until ready for analysis. For each sample, 2×10^6 cells were placed in a quartz cuvette and the fluorescence was measured using a PTI QM-2000-4 spectrofluorometer (Photon Technology International (PTI), Birmingham, NJ) with constant stirring. The fluorescence of the fluo-3 dye was measured with excitation at 490 nm and emission at 525 nm. The fluorescence was measured and digitized at 1 Hz using the software program Felix 1.42b (PTI). Data points were collected every second and cells were either treated with vehicle control, DCPA, its

metabolites or analogs and 2 μ M thapsigargin (Sigma) or 2 μ M thapsigargin alone. Reagents were added after baseline data was collected for 45 seconds. Following the return of fluorescence to background levels, 2.5mM CaCl_2 (Sigma) was added to the media. Addition of 200 μ M ionomycin (Sigma) ensured even loading of the cells. Cell membranes were lysed with 0.1% (v/v) Triton X-100 (Fisher Scientific, Hampton, NH) to measure the maximum fluorescence ($F_{\max}$) parameter for calculation of $[\text{Ca}^{2+}]_i$ and to monitor compartmentalization of the dye. To chelate the free Ca^{2+} to a nominally Ca^{2+} -free level ($F_{\min}$), 50mM ethylene glycol-bis(β -aminoethyl ether)-N,N,N',N'-tetraacetic acid tetrasodium salt (EGTA), pH 7.5 (Sigma) was added to the cuvette.

Fluorescence values were converted to $[\text{Ca}^{2+}]_i$ using the following equation:

$$\text{free}[\text{Ca}^{2+}]_i = K_D[(F - F_{\min}) / (F_{\max} - F)],$$

where K_D (360 nM) is the dissociation constant of the Fluo-3/ Ca^{2+} complex, F is the measured fluorescence intensity, $F_{\min}$ is the minimum fluorescence at very low $[\text{Ca}^{2+}]_i$ (fluorescence after the addition of 50 mM EGTA) and $F_{\max}$ is the fluorescence measured at high $[\text{Ca}^{2+}]_i$ (fluorescence after the addition of Triton X-100) (Grynkiewicz, Poenie et al. 1985). The background fluorescence obtained from unloaded cells over a 3 minute time period was subtracted from all data points before $[\text{Ca}^{2+}]_i$ was calculated.

Measurement of IL-2 Production

Jurkat cells were cultured in complete RPMI media at 5×10^5 cells/well in 48-well plates (Costar, Corning, NY) coated with mouse anti-human CD3 antibody (10 μ g/ml) (BD BioSciences, San Diego, CA). Cells were treated with varying concentrations of DCPA, DCA, DFPA, DFA, NOH-DCA, 6OH-DCA, ethanol or DMSO vehicle control. Cells were also simultaneously stimulated with anti-CD28 antibody (2 μ g/ml) (BD BioSciences).

Cells were incubated at 37 °C in 5 % CO₂ for 24 h after which supernatants were collected and placed at -20°C. IL-2 production was determined using the sandwich ELISA method and following the manufacturer's protocol (BD PharMingen). All cultures and ELISA analyses were performed in triplicate and the experiment was repeated at a minimum of three times.

Transfections and Luciferase assay

Jurkat cells were plated in RPMI media with 1.5% FBS at 6×10^5 cells/well in a 6-well plate (Costar). For each well, 750 ng of pNFAT-luc (firefly luciferase plasmid) (Stratagene) and 10ng pRL-TK (Renilla luciferase plasmid) (Promega) were transfected with 1 ul Lipofectamine 2000 (Invitrogen). Cells were incubated for 5 h at 37°C in 5% CO₂ and media was replaced with complete RPMI +10% FBS and incubated overnight at 37° C in 5% CO₂. Following transfection, cells were treated with varying concentrations of DCPA, DCA, DFPA, DFA, NOH-DCA, 6OH-DCA, ethanol or DMSO or left untreated and stimulated with 10ng/ ml PMA and 1ug/ ml A23187. Cells were incubated for 4 h at 37°C in 5% CO₂, centrifuged, lysed and stored at -70°C until ready for analysis. NFAT activity was determined using the Dual Luciferase Assay Kit and following the manufacturer's protocol (Promega). Briefly, 25ul of sample was added to 100ul of Luciferase Assay Reagent II and the firefly luciferase activity measured. Addition of Stop & Glo quenched the firefly luminescence and provides a substrate for the Renilla luciferase activity. Luminescence was detected using a Berthold Lumat LB 9507 (Oak Ridge, TN). NFAT-firefly luciferase transfection efficiency was normalized to the Renilla luciferase activity and the average fold change was reported. All luciferase assays were performed in triplicate and the experiment was repeated at least twice.

Statistical Analysis

All data were analyzed using MS Excel 2007 (Redmond, WA) and Sigma Stat 3.1 (Port Richmond, C A). A NOVA with a Student-Newman Keuls post hoc test was used to determine statistical significance with an alpha value of <0.05 considered significant.

RESULTS

Hydroxy metabolites of DCPA are more cytotoxic than its parent.

In mammals, metabolism of DCPA results in the hydrolysis of the amide side chain and production of the 3,4-DCA with further biotransformation resulting in the formation of 6OH-DCA and NOH-DCA. To assess the cytotoxicity of DCPA and its metabolites on Jurkat T cells, viability and proliferation assays were conducted over a range of concentrations. For cells exposed to DCPA, DCA, DFPA, and DFA, ethanol (0.1%v/v) was used as a vehicle control and DMSO was used as a vehicle control for 6OH-DCA and NOH-DCA. Jurkat T cells were treated or left untreated, loaded with 7-AAD and viability was assessed after 24 hours (Fig.2). The viability of cells treated with increasing concentrations of DCPA up to 100 μ M did not decrease viability (Fig.2A). Treatment with 200 μ M DCPA decreases viability by approximately 14.6% (Fig.2A) and subsequent studies were conducted using a maximum concentration of 100 μ M DCPA. DCA appeared to be less toxic and concentrations up to 200 μ M were not cytotoxic (Fig.2B). However, Jurkat T cells were more sensitive to the hydroxylated DCA metabolites and cytotoxicity was observed at 100 μ M for both 6OH-DCA and NOH-DCA (Fig.2C and 2D). This concentration was excluded for both 6OH-DCA and NOH-DCA in all subsequent assays. Proliferation assays were also performed and proliferation of Jurkat T cells was inhibited only at those concentrations that were also cytotoxic (data not shown).

Figure 2A and 2B Cytotoxic effects of DCPA and DCA

Jurkat T cells were treated with or without varying concentrations of A) DCPA, B) DCA, or their vehicle control, ethanol for 24 h. Cells were stained with 7-AAD to determine viability. Results using flow cytometry are representative of 3 experiments each performed in triplicate. Error bars reflect $\pm$ SD.

Figure 2C and 2D Cytotoxic effects of 6OH-DCA and NOH-DCA

Jurkat T cells were treated with or without varying concentrations of, C) 6OH-DCA, D) NOH-DCA, or their vehicle control DMSO for 24 h. Cells were stained with 7-AAD to determine viability using flow cytometry. Results are representative of 3 experiments each performed in triplicate. Error bars reflect $\pm$ SD.

DCPA and its metabolites decrease IL-2 secretion.

We have previously reported that exposure to DCPA decreases IL-2 production in Jurkat T cells in a calcium-dependent manner (Lewis 2008). IL-2 is an important early cytokine produced by T cells and is critical for the activation and proliferation of several immune cells, including NK cells, T cells and B cells. To determine if exposure to metabolites of DCPA alter IL-2 production, Jurkat T cells were treated over range of concentrations with or without vehicle controls (DMSO for the hydroxylated metabolites and ethanol for all others) and stimulated with anti-CD3 and anti-CD28. After 24 h in culture, IL-2 secretion levels in the supernatant were assessed using an ELISA. DCPA decreased IL-2 secretion in a concentration-dependent manner with significant decreases observed at 25 μ M, 50 μ M, and 100 μ M DCPA and represented a 20%, 48% and 74% decrease in IL-2 production, respectively (Fig. 3A). This data is in agreement with previously reported decreases of IL-2 in response to DCPA treatment.

Metabolism of DCPA is reported to occur in the liver through the action of acylamidases which cleave the amide side chain resulting in the production of DCA (McMillan). Several studies have reported detectable levels of DCA in blood and urine in DCPA-exposed individual. When Jurkat cells are exposed to increasing concentrations of DCA (25-200 μ M), concentration-dependent decreases in IL-2 secretion are also observed (Fig. 3B). At a concentration of 50 μ M decreases are observed but statistically significant decreases are only observed at 100 μ M and 200 μ M DCA (Fig. 3B). Exposure to 50 μ M, 100 μ M and 200 μ M DCA resulted in a 10%, 45% and 78% decrease in IL-2, respectively. These results suggest that, in T cells, DCPA is more potent inhibitor of IL-2 than its metabolite, DCA.

In mammals, hydroxylation of DCA to 6OH-DCA and NOH-DCA occurs in the liver and both oxidative metabolites are capable of converting oxy-hemoglobin (Hb) to met-Hb. In Jurkat T cells, NOH-DCA and 6OH-DCA are cytotoxic at 100 μ M so to assess their effects on IL-2 secretion lower concentrations were evaluated. Both 6OH-DCA and NOH-DCA decreased IL-2 secretion however NOH-DCA is a more potent inhibitor of IL-2 secretion (Fig. 3C and 3D). 6OH-DCA decreased IL-2 production by 30% at the highest concentration (50 μ M) whereas 25 μ M and 50 μ M NOH-DCA decreased IL-2 secretion by 51% and 90%, respectively. No change in IL-2 secretion was observed in cells treated with 5 μ M NOH-DCA. This data indicates that Jurkat T cells are highly sensitive to exposure of NOH-DCA.

Figure 3A and 3B DCPA and DCA inhibit IL-2 secretion

Jurkat T cells were treated with or without varying concentrations of A) DCPA, B) DCA, and stimulated with anti-CD3 and anti-CD28 for 24h. Supernatants were analyzed by ELISA and results are representative of three separate experiments. Error bars reflect $\pm$ SD and asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

Figure 3C and 3D NOH-DCA and 6OH-DCA inhibit IL-2 secretion

Jurkat T cells were treated with or without varying concentrations of C) 6OH-DCA, D) NOH-DCA and stimulated with anti-CD3 and anti-CD28 for 24 h. Supernatants were analyzed by ELISA and results are representative of three separate experiments. Error bars reflect $\pm$ SD and asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

D CPA and its metabolites alter NFAT activity

In T cells, optimal transcription of the IL-2 gene requires coordinated binding of three transcription factors; NFAT (nuclear factor of activated T cells), NF- κ B (nuclear factor kappaB) and AP-1 (activator protein-1). We have previously reported that DCPA-exposed Jurkat cells have decreased AP-1 DNA binding ability, decreased c-jun phosphorylation and decreased nuclear NFAT levels. To further understand the effects of DCPA and its metabolites on IL-2 secretion Jurkat T cells were co-transfected with a NFAT Luciferase plasmid (pNFAT-luc) and a Renilla Luciferase plasmid (pRL-TK), to control for transfection efficiencies. In resting T cells, NFAT is localized to the cytoplasm in a phosphorylated state and upon stimulation becomes de-phosphorylated by the calcium-dependent phosphatase, calcineurin. De-phosphorylation of NFAT exposes a nuclear localization sequence facilitating its translocation into the nucleus and binding to the promoter region of the targeted gene. In this system, stimulation of Jurkat cells leads to the de-phosphorylation of endogenous NFAT that binds to the NFAT DNA binding sites on the pNFAT-luc plasmid which controls expression of the firefly luciferase. In cells exposed to 25-100 μ M DCPA there is a significant decrease in NFAT activity (Fig 4A), with a 37 %, 63% and 87% decrease in NFAT activity in cells treated with 25 μ M, 50 μ M and 100 μ M DCPA, respectively. Similarly, cells treated with DCA also have decreased NFAT activity at those concentrations that resulted in decreased IL-2 secretion (Fig 3B). Cells exposed to 100 μ M and 200 μ M DCA statistically decreased NFAT activity levels by 66 and 18%, respectively (Fig 4B). 6OH-DCA-treated cells had a 30% decrease in NFAT activity at 50 μ M but no change at 25 μ M (Fig.4C). NOH-DCA decreased NFAT activity by 33% and 47% at 25 μ M and

50 μ M, respectively (Fig.4D). Although 50 μ M NOH-DCA inhibited IL-2 secretion by 90% NFAT activity is only decreased by 47%, whereas 100 μ M DCPA decreased IL-2 by 74% with an 87 % decrease in NFAT activity. This data suggests that NOH-DCA may inhibit IL-2 through an alternate mechanism than DCPA. In all experiments, cells treated with cyclosporin, an inhibitor of NFAT that prevents dephosphorylation, produced similar levels of NFAT activity as those in unstimulated cells.

Figure 4A DCPA alters NFAT activity

Jurkat T cells were co-transfected with a firefly luciferase-NFAT reporter plasmid and a Renilla luciferase plasmid to control for transfection efficiency. Cells were treated with or without varying concentrations of DCPA or vehicle control and stimulated with PMA and A23187 for 5 h. Unstimulated and cyclosporin treated cells were included as controls. Cell lysates were analyzed for luciferase activity and normalized to Renilla activity. Results are representative of at least 2 experiments each performed in triplicate. Asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

Figure 4B DCA alters NFAT activity

Jurkat T cells were co-transfected with a firefly luciferase-NFAT reporter plasmid and a Renilla luciferase plasmid to control for transfection efficiency. Cells were treated with or without varying concentrations of DCA or vehicle control and stimulated with PMA and A23187 for 5 h. Unstimulated and cyclosporin treated cells were included as controls. Cell lysates were analyzed for luciferase activity and normalized to Renilla activity. Results are representative of at least 2 experiments each performed in triplicate. Asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

Figure 4C. 6OH-DCA alters NFAT activity.

Jurkat T cells were co-transfected with a firefly luciferase-NFAT reporter plasmid and a Renilla luciferase plasmid to control for transfection efficiency. Cells were treated with or without varying concentrations 6OH-DCA or vehicle control and stimulated with PMA and A23187 for 5 h. Unstimulated and cyclosporin treated cells were included as controls. Cell lysates were analyzed for luciferase activity and normalized to Renilla activity. Results are representative of at least 2 experiments each performed in triplicate. Asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

Figure 4D. NOH-DCA alters NFAT activity

Jurkat T cells were co-transfected with a firefly luciferase-NFAT reporter plasmid and a Renilla luciferase plasmid to control for transfection efficiency. Cells were treated with or without varying concentrations of NOH-DCA or vehicle control and stimulated with PMA and A 23187 for 5 h. Unstimulated and cyclosporin treated cells were included as controls. Cell lysates were analyzed for luciferase activity and normalized to Renilla activity. Results are representative of at least 2 experiments each performed in triplicate. Asterisks (*) indicate statistically significant results, $p < 0.05$ using ANOVA with a Student-Newman Keuls post hoc test.

DCPA and its metabolites have differential effects on Ca homeostasis.

It has been previously reported that exposure of Jurkat T cells to DCPA results in decreased intracellular calcium influx following ER Ca store depletion. To examine the effect of DCPA metabolites on calcium homeostasis, Jurkat T cells were loaded with a calcium-sensitive dye and changes in intracellular calcium were monitored over time. At the start of these experiments, the extracellular media is essential Ca-free and depletion of ER Ca stores with thapsigargin, an inhibitor of the SERCA (Sarco/endoplasmic reticulum Ca ATPase) pump, results in a small, transient increase in intracellular Ca (Fig.5). Addition of 2mM Ca to the media results in a large and sustained increase in intracellular Ca that reflects store-operated calcium influx (Fig 5). Jurkat T cells exposed to DCPA decrease calcium influx following store depletion in a concentration dependent manner (Fig 5A). At concentrations of 100 μ M and 200 μ M DCA, decreases in intracellular calcium are also observed (Fig 5B). It should be noted that DCPA and DCA-treated cells do not alter the transient increase in intracellular calcium observed when ER calcium stores are depleted. Interestingly, cells treated with 6OH-DCA and NOH-DCA did not inhibit ER calcium store depletion or calcium influx (Fig 5C and 5D). This data supports the conclusion that hydroxylated metabolites alter IL-2 secretion through a different mechanism than that of DCPA and DCA.

Figure 5A DCPA alters calcium influx

Jurkat T cells were loaded with the calcium-sensitive dye, fluo-3, and treated with or without varying concentrations of DCPA or vehicle control. Immediately following addition of treatment, 2 μ M thapsigargin was added to deplete Ca stores. When fluorescence returned to baseline 2 mM CaCl_2 was added and the effect on Ca influx was recorded. Results are representative of at least 3 experiments.

Figure 5B DCA alters calcium influx

Jurkat T cells were loaded with the calcium-sensitive dye, fluo-3, and treated with or without varying concentrations of DCA or vehicle control. Immediately following addition of treatment, 2 μ M thapsigargin was added to deplete Ca stores. When fluorescence returned to baseline 2 mM CaCl_2 was added and the effect on Ca influx was recorded. Results are representative of at least 3 experiments.

C

Figure 5C 6OH-DCA does not alter intracellular calcium concentrations

Jurkat T cells were loaded with the calcium-sensitive dye, fluo-3, and treated with or without varying concentrations of 6OH-DCA or vehicle control. Immediately following addition of treatment, 2 μM thapsigargin was added to deplete Ca stores. When fluorescence returned to baseline 2 mM CaCl₂ was added and the effect on Ca influx was recorded. Results are representative of at least 3 experiments.

Figure 5D NOH-DCA does not alter intracellular calcium concentrations

Jurkat T cells were loaded with the calcium-sensitive dye, fluo-3, and treated with or without varying concentrations of NOH-DCA or vehicle control. Immediately following addition of treatment, 2 μ M thapsigargin was added to deplete Ca stores. When fluorescence returned to baseline 2 mM CaCl₂ was added and the effect on Ca influx was recorded. Results are representative of at least 3 experiments.

Chlorine substituents play a role in the immunotoxic effects of DCPA

To assess the role that chlorines, in the 3 and 4 positions on DCPA, may have on the immunotoxic effects in T cells, we synthesized the fluorine analogs of DCPA, DFPA (see materials and methods) and used DFA, a commercially available fluorine analog of DCA. Concentrations up to 400 μ M of both DFPA and DFA were not cytotoxic to Jurkat T cells (Fig 6A and 6B). Proliferation assays were also conducted and no changes in proliferation after 24 hours were observed at concentration up to 400 μ M (data not shown). To determine the effect of DFPA and DFA on IL-2 secretion, Jurkat T cells were stimulated with anti-CD3 and anti-CD28 and IL-2 was measured as described above. No changes in IL-2 secretion were detected when Jurkat cells were exposed to DFPA or DFA with increasing concentrations up to 400 μ M (Fig 6C and 6D). These results indicate the presence of chlorine at the 3 and 4 positions play an important role in the inhibiting IL-2 secretion in DCPA and DCA exposed T cells. To confirm the apparent inert effects of fluorine substitution, NFAT activity and Ca influx were also examined as described above. Consistent with the IL-2 secretion data, Jurkat T cells exposed to DFPA and DFA did not alter NFAT activity or Ca homeostasis (Fig 6E-6H). This data provides clear evidence that the immunotoxic effects of DCPA and its metabolites, on T cells, can be attributed, in large part, to the presence and location of chlorines at the 3 and 4 positions.

Figure 6A and 6B Cytotoxic effects of DFPA and DFA

Jurkat T cells were treated with or without varying concentrations of A) DFPA or B) DFA and vehicle control for 24h. Cells were stained with 7-AAD to determine viability and analyzed with flow cytometry. Results are representative of 3 experiments each performed in triplicate. Error bars reflect $\pm$ SD.

Figure 6C and 6D DFPA and DFA do not inhibit IL-2 secretion

Jurkat T cells were treated with or without varying concentrations of A) DFPA or B) DFA, and vehicle control and stimulated with anti-CD3 and anti-CD28 for 24 h. Supernatants were collected and analyzed by ELISA. Results are representative of three separate experiments. Error bars reflect $\pm$ SD.

Figure 6E and 6F DFPA and DFA DCPA do not alter NFAT activity

Jurkat T cells were co-transfected with a firefly luciferase-NFAT reporter plasmid and a Renilla luciferase plasmid to control for transfection efficiency. Cells were treated with or without varying concentrations of E) DFPA or F) DFA and vehicle control and stimulated with PMA and A23187 for 5 h. Unstimulated and cyclosporin treated cells were included as controls. Cell lysates were analyzed for luciferase activity and normalized to Renilla activity. Results are representative of at least 2 experiments each performed in triplicate.

Figure 6G and 6H DFPA and DFA do not alter intracellular calcium concentrations

Jurkat T cells were loaded with the calcium-sensitive dye, fluo-3, and treated with or without varying concentrations of G) DFPA, H) DFA and vehicle control. 2 μ M thapsigargin was added to deplete Ca stores. When fluorescence returned to baseline 2 mM CaCl_2 was added and the effect on Ca influx was recorded. Results are representative of at least 3 experiments.

DISCUSSION

In the United States, agricultural use of herbicides represents 58% of total pesticide use and approximately 2 billion pounds are applied to crop fields every year (Grube 2004). DCPA is a commonly used herbicide in the United States and is used on 50-80% of all rice crops for control against broadleaf and grassy plants. The toxic effects of DCPA on the immune system have been well documented (reviewed in (Salazar, Ustyugova et al. 2008). In mouse models, *in vivo* and *in vitro* exposure to DCPA decreases the phagocytic ability of macrophages and its ability to produce IL-6 and TNF- α when stimulated with LPS (Xie, Schafer et al. 1997; Xie, Schafer et al. 1997; Frost, Neeley et al. 2001; Ustyugova, Frost et al. 2007). Primary stimulation of CD8⁺ T cells exposed to DCPA is not affected but the secondary response, in the absence of further exposure to DCPA, abrogates lytic activity (Barnett and Gandy 1989; Seil, Frankenberry et al. 2006). In human Jurkat T cells, DCPA decreases IL-2 protein and mRNA levels as well as the binding ability of the transcription factor AP-1 (activator protein-1) and the protein levels and phosphorylation of c-jun (Zhao, Schafer et al. 1999; Brundage, Schafer et al. 2004). The decreases in IL-2 secretion observed in DCPA-treated T cells are mediated by alterations in NFAT translocation and calcium homeostasis (Lewis, Brundage et al. 2008). Although the immunotoxic effects of DCPA are well documented little is known about the effects of its metabolites.

In mammals, DCPA is metabolized in the liver by acylamidases resulting in the production of DCA which can be hydroxylated to form 6OH-DCA and NOH-DCA (Fig.1). Exposure to DCA can occur in several ways, as breakdown products of DCPA and other herbicides such as linuron and diuron and also through the manufacturing of several pesticides, dyes and pharmaceuticals (SCTEE 2001). In humans, DCPA has

an estimated half life of 3.2 hours and is quickly metabolized into DCA which can be detected in the urine and blood of occupationally and non-occupationally exposed humans (Wittke, Hajimiragha et al. 2001; Turci, Barisano et al. 2006; Roberts, Heilmair et al. 2009).

Limited data is available on the immunotoxic effects of DCA but Barnett *et al* reported that molar equivalent concentrations of DCA and DCPA decreased natural killer (NK) cell activity and increased spleen weight (Barnett, Gandy et al. 1992). NOH-DCA and 6OH-DCA are reactive metabolites that have been studied for their effects on erythrocytes (Singleton and Murphy 1973). Both NOH-DCA and 6OH-DCA can bind hemoglobin leading to methemoglobinemia, a serious medical condition that results in the formation of methemoglobin adducts that do not bind or transport oxygen. NOH-DCA has also been reported as a nephrotoxicant and can induce hemolytic anemia (McMillan, Bradshaw et al. 1991; Valentovic, Ball et al. 2001). No data exists on the immunotoxic effects of NOH-DCA and 6OH-DCA.

Studies on exposure and the effects of DCPA in human subjects are limited. Richards *et al* measured air levels of DCPA in farms adjacent to rice fields and determined that individuals living near rice fields are at risk for exposure to DCPA (Richards, McClure et al. 2001). Accompanying this study, McClure *et al* reported that individuals living in areas sampled by Richards et al did not appear to be at higher risk for altered immune functions (McClure, Helm et al. 2001). Although individuals surrounding rice fields may not be at risk, there is evidence to suggest that workers in the agricultural and manufacturing sectors are at risk. Corsini *et al* examined agricultural workers exposed before, during and after exposure to DCPA (Corsini,

Codeca et al. 2007). DCPA was readily detected in the urine of workers at concentrations up to 332ng/ml. Whole blood assays of the workers revealed that cells stimulated with PHA had decreased IL-10 and IFN- γ production. In addition, Pastorelli et al. report that DCPA-Hb adducts are detectable in agricultural workers exposed to DCPA for as long as 4 months following the last application of the herbicide (Pastorelli 1998). This suggests that metabolites of DCPA can remain in the body for an extended period of time.

We report here for the first time the immunotoxic effects of the DCPA metabolites, DCA, NOH-DCA and 6OH-DCA on human Jurkat T cell function. Jurkat T cells have been used for over 20 years as a model for human T cell signaling (Abraham 2004). Previous research has demonstrated that exposure of DCPA to human Jurkat T cells inhibits IL-2 secretion in a concentration and Ca-dependent manner. Similar to DCPA, DCA inhibits IL-2 secretion, NFAT activity and calcium homeostasis, although at higher concentrations than that of DCPA (Fig 3B, 4B, 5B). This is in agreement with others who have reported that immune parameters, including T cell dependent antibody production, myelotoxic effects, and IL-6 response in mouse T cells, were altered by DCPA but required higher concentrations of DCA to produce similar effects (Barnett, Gandy et al. 1992; Malerba, Castoldi et al. 2002)(Zhao unpubl data).

NOH-DCA appears more cytotoxic to T cells than both DCPA or DCA and inhibits IL-2 secretion more potently, with a 90% reduction in IL-2 at 50 μ M NOH-DCA whereas at 50 μ M DCPA IL-2 is decreased by only 50% (Fig 2A, 2C, 3A, 3C). However, a 2.5 fold decrease in NFAT activity in cells exposed to 50 μ M DCPA resulted in a 50% decrease in IL-2 but 50 μ M NOH-DCA decreased IL-2 by 90% with only a 2 fold change

in NFAT activity (Fig. 4A and 4C). The effect of these metabolites on calcium homeostasis suggests that the mechanism by which NOH-DCA inhibits IL-2 secretion is different from that of DCPA (Fig 5A and 5C). NOH-DCA and 6OH-DCA appeared to inhibit IL-2 in a Ca-independent manner, whereas the effects of DCPA and DCA on T cell function are elicited in a Ca-dependent manner. There is no data available on the immunotoxic effects of 6OH-DCA, however, McMillan *et al* reported that although 6OH-DCA produced metHb adducts NOH-DCA was 10 times more potent (McMillan, Leakey *et al.* 1990). Our data supports a diminished effect of 6OH-DCA on T cell function. 6OH-DCA and NOH-DCA are equally cytotoxic but 6OH-DCA only decreased IL-2 secretion at 50µM compared to 5µM NOH-DCA (Fig 2C and 2D). Calcium homeostasis is also unaffected by treatment with 6OH-DCA. The mechanism for the decreased in IL-2 and NFAT activity in 6OH-DCA and NOH-DCA exposed cells is unknown.

Approximately 45% of all herbicides contain a carbon-chlorine bond and 1/6th of all organochlorines requires special safety precautions for use in the workplace (Naumann 2000). In many cases the biological activity of the compound is conferred by the presence of the chlorines (Naumann 2000). The best known example is DDT, as the removal of two specific chlorines renders this pesticide inactive (Crinnion 2009). Other toxic organochlorine pesticides include atrazine and 2,4-D. In order to determine the role that chlorine substituents have in DCPA-exposed T cells, we substituted fluorines for both of the chlorines found in DCPA and DCA. In the last 20 years, the number of fluorinated chemicals has increased significantly and now 28% of all halogenated agrochemicals are fluorinated (Jeschke 2004). The position and number of fluorines in agrochemicals determines the activity of many pesticides (Jeschke 2004).

Unlike DCPA and DCA, DFPA and DFA are not cytotoxic (up to 400 μ M) and do not alter IL-2 secretion, NFAT activity or calcium homeostasis (Fig. 6). This indicates that chlorine substitution plays an important role in exerting immunotoxic effects in T cells. It has been reported that 3,4-DFA is toxic to the liver but at concentrations higher than that for 3,4-DCA, suggesting a negative biological role for chlorine at the 3 and 4 positions (Hong, Anestis et al. 2000). Several possibilities exist for the differential effects observed with chlorine and fluorine. First, fluorines are highly electronegative and act only as hydrogen acceptors whereas chlorine and other halogens act as both hydrogen acceptors and donors. Second, trifluoro-substitution can increase lipophilicity but mono- or difluoro-substitution has been reported to decrease it (Purser, Moore et al. 2008). It has recently been reported that, in Jurkat T cells, DCPA becomes localized to the cytosol, indicating that it can pass through the membrane (Hanson, Peer et al. 2010). Increases in electronegativity and lipophilicity could prevent binding to proteins targeted by DCPA. Lastly, fluorines are similar in size to hydrogen and may not produce the conformation required to elicit the effects of DCPA. These specific properties of fluorine may be responsible for the apparent reversal of effects observed in DFPA and DFA treated cells.

In conclusion, metabolites of DCPA display differential effects on Jurkat T cells. Although DCA elicits similar Ca-dependent effects on IL-2 secretion, the parent compound, DCPA, is more toxic. However, DCPA is quickly metabolized to DCA and persists in the body possibly allowing DCA to accumulate to concentrations greater than that of DCPA. Hydroxylation of DCA can result in the production of NOH-DCA and 6OH-DCA which, in Jurkat T cells, can be cytotoxic and can alter IL-2 secretion in a Ca-

independent manner at low concentration (5 μ M NOH-DCA). Further studies are required to determine the mechanism for the decreases in IL-2 observed in NOH-DCA-exposed T cells and to more fully understand the mechanism by which DCPA elicits its effects.

REFERENCES

- Barnett, J. B. and J. Gandy (1989). "Effect of acute propanil exposure on the immune response of C57Bl/6 mice." Fundam Appl Toxicol **12**(4): 757-64.
- Barnett, J. B., J. Gandy, et al. (1992). "Comparison of the immunotoxicity of propanil and its metabolite, 3,4-dichloroaniline, in C57Bl/6 mice." Fundam Appl Toxicol **18**(4): 628-31.
- Bauer, E. R., H. H. Meyer, et al. (1998). "Application of an androgen receptor assay for the characterisation of the androgenic or antiandrogenic activity of various phenylurea herbicides and their derivatives." Analyst **123**(12): 2485-7.
- Brundage, K. M., R. Schafer, et al. (2004). "Altered AP-1 (activating protein-1) activity and c-jun activation in T cells exposed to the amide class herbicide 3,4-dichloropropionanilide (DCPA)." Toxicol Sci **79**(1): 98-105.
- Corsini, E., I. Codeca, et al. (2007). "Immunomodulatory effects of the herbicide propanil on cytokine production in humans: In vivo and in vitro exposure." Toxicol Appl Pharmacol **222**(2): 202-10.
- Crinnion, W. J. (2009). "Chlorinated pesticides: threats to health and importance of detection." Altern Med Rev **14**(4): 347-59.
- Frost, L. L., Y. X. Neeley, et al. (2001). "Propanil inhibits tumor necrosis factor-alpha production by reducing nuclear levels of the transcription factor nuclear factor-kappaB in the macrophage cell line ic-21." Toxicol Appl Pharmacol **172**(3): 186-93.
- Grube, A. (2004). Pesticides Industry Sales and Usage. P. Office of Prevention, and Toxic Substances, U.S. Environmental Protection Agency.
- Grynkiewicz, G., M. Poenie, et al. (1985). "A new generation of Ca²⁺ indicators with greatly improved fluorescence properties." J Biol Chem **260**(6): 3440-50.
- Hanson, M. L., C. J. Peer, et al. (2010). "Subcellular localization of the amide class herbicide 3,4-dichloropropionanilide (DCPA) in T cells and hepatocytes." J Toxicol Environ Health A **73**(1): 1-4.
- Hong, S. K., D. K. Anestis, et al. (2000). "Haloaniline-induced in vitro nephrotoxicity: effects of 4-haloanilines and 3,5-dihaloanilines." Toxicol Lett **114**(1-3): 125-33.

- Jeschke, P. (2004). "The unique role of fluorine in the design of active ingredients for modern crop protection." Chembiochem **5**(5): 571-89.
- Lerman, L., M. Weinstock-Rosin, et al. (2004). "An Improved Synthesis of Hydroxyindoles." Synthesis **18**: 3043-3046.
- Lewis, T. L., K. M. Brundage, et al. (2008). "3,4-Dichloropropionanilide (DCPA) inhibits T-cell activation by altering the intracellular calcium concentration following store depletion." Toxicol Sci **103**(1): 97-107.
- Lok, R., R. Leone, et al. (1996). "Facile Rearrangements of Alkynylamino Heterocycles with Noble Metal Cations." The journal of organic chemistry **61**(10): 3289-3297.
- Malerba, I., A. F. Castoldi, et al. (2002). "In vitro myelotoxicity of propanil and 3,4-dichloroaniline on murine and human CFU-E/BFU-E progenitors." Toxicol Sci **69**(2): 433-8.
- McClure, G. Y., R. M. Helm, et al. (2001). "Evaluation of immune parameters in propanil-exposed farm families." Arch Environ Contam Toxicol **41**(1): 104-11.
- McMillan, D. C., T. P. Bradshaw, et al. (1991). "Role of metabolites in propanil-induced hemolytic anemia." Toxicol Appl Pharmacol **110**(1): 70-8.
- McMillan, D. C., J. E. Leahey, et al. (1990). "Metabolism of the arylamide herbicide propanil. II. Effects of propanil and its derivatives on hepatic microsomal drug-metabolizing enzymes in the rat." Toxicol Appl Pharmacol **103**(1): 102-12.
- Naumann, K. (2000). "Influence of chlorine substituents on biological activity of chemicals: a review." Pest Manag Sci **56**(1): 3-21.
- Pastorelli, R., Catenacci, G., Guanci, M., Fannelli, R., Valoti, E., Minoia, C., Airoidi, L (1998). "3,4 Dichloroaniline-haemoglobin adducts in humans: preliminary data on agricultural workers exposed to propanil." Biomarkers **3**(3): 227-233.
- Purser, S., P. R. Moore, et al. (2008). "Fluorine in medicinal chemistry." Chem Soc Rev **37**(2): 320-30.
- Richards, S. M., G. Y. McClure, et al. (2001). "Propanil (3,4-dichloropropionanilide) particulate concentrations within and near the residences of families living adjacent to aerially sprayed rice fields." Arch Environ Contam Toxicol **41**(1): 112-6.

- Roberts, D. M., R. Heilmair, et al. (2009). "Clinical outcomes and kinetics of propanil following acute self-poisoning: a prospective case series." BMC Clin Pharmacol **9**: 3.
- Salazar, K. D., P. de la Rosa, et al. (2005). "The polysaccharide antibody response after *Streptococcus pneumoniae* vaccination is differentially enhanced or suppressed by 3,4-dichloropropionanilide and 2,4-dichlorophenoxyacetic acid." Toxicol Sci **87**(1): 123-33.
- Salazar, K. D., I. V. Ustyugova, et al. (2008). "A review of the immunotoxicity of the pesticide 3,4-dichloropropionanilide." J Toxicol Environ Health B Crit Rev **11**(8): 630-45.
- SCTEE (2001). "Risk Assessment of 3,4-dichloroaniline." Scientific Committee on Toxicity, Ecotoxicity, and the Environment.
- Sheil, J. M., M. A. Frankenberry, et al. (2006). "Propanil exposure induces delayed but sustained abrogation of cell-mediated immunity through direct interference with cytotoxic T-lymphocyte effectors." Environ Health Perspect **114**(7): 1059-64.
- Singleton, S. D. and S. D. Murphy (1973). "Propanil (3,4-dichloropropionanilide)-induced methemoglobin formation in mice in relation to acylamidase activity." Toxicol Appl Pharmacol **25**(1): 20-9.
- Still, G. G. (1968). "Metabolic Fate of 3,4-Dichloropropionanilide in Plants: The Metabolism of the Propionic Acid Moiety." Plant Physiol **43**(4): 543-546.
- Turci, R., A. Barisano, et al. (2006). "Determination of dichloroanilines in human urine by gas chromatography/mass spectrometry: validation protocol and establishment of Reference Values in a population group living in central Italy." Rapid Commun Mass Spectrom **20**(17): 2621-5.
- USEPA (2006). Amendment to Reregistration Eligibility Decision for Propanil. U. S. E. P. Agency.
- Ustyugova, I. V., L. L. Frost, et al. (2007). "3,4-dichloropropionaniline suppresses normal macrophage function." Toxicol Sci **97**(2): 364-74.
- Valentovic, M., J. G. Ball, et al. (2001). "3,4-Dichlorophenylhydroxylamine cytotoxicity in renal cortical slices from Fischer 344 rats." Toxicology **162**(3): 149-56.

- Valentovic, M. A., T. Yahia, et al. (1997). "3,4-Dichloroaniline acute toxicity in male Fischer 344 rats." Toxicology **124**(2): 125-34.
- Wittke, K., H. Hajimiragha, et al. (2001). "Determination of dichloroanilines in human urine by GC-MS, GC-MS-MS, and GC-ECD as markers of low-level pesticide exposure." J Chromatogr B Biomed Sci Appl **755**(1-2): 215-28.
- Xie, Y. C., R. Schafer, et al. (1997). "The immunomodulatory effects of the herbicide propanil on murine macrophage interleukin-6 and tumor necrosis factor-alpha production." Toxicol Appl Pharmacol **145**(1): 184-91.
- Xie, Y. C., R. Schafer, et al. (1997). "Inhibitory effect of 3,4-dichloro-propionaniline on cytokine production by macrophages is associated with LPS-mediated signal transduction." J Leukoc Biol **61**(6): 745-52.
- Zhang, B. and S. Lin (2009). "Effects of 3,4-dichloroaniline on testicle enzymes as biological markers in rats." Biomed Environ Sci **22**(1): 40-3.
- Zhao, W., R. Schafer, et al. (1999). "Propanil affects transcriptional and posttranscriptional regulation of IL-2 expression in activated EL-4 cells." Toxicol Appl Pharmacol **154**(2): 153-9.
- Zhao, W., R. Schafer, et al. (1995). "Changes in primary and secondary lymphoid organ T-cell subpopulations resulting from acute in vivo exposure to propanil." J Toxicol Environ Health **46**(2): 171-81.

CHAPTER 4

The inhibitory effects of 3,4-dichloropropionanilide (DCPA) on Stromal Interaction Molecule-1 (Stim1) puncta formation.

Tricia L. Lewis* and John B. Barnett*

* Department of Microbiology, Immunology and Cell Biology, West Virginia University
School of Medicine, Morgantown, WV 26506

Running title: DCPA inhibits Stim1 puncta formation

Address correspondence to:

John B. Barnett, Ph.D.

Department of Microbiology, Immunology and Cell Biology

West Virginia University School of Medicine

P.O. Box 9177

Morgantown, West Virginia 26506-9177

Tel. 304-293-4029; Fax 304-293-7823

E-Mail: jbarnett@hsc.wvu.edu

Keywords: pr opanil, 3, 4-dichloropronanilide, DCPA, Stromal I nteracting M olecule-1,
Stim-1

ABSTRACT

Calcium (Ca) is a ubiquitous second messenger that is required in almost all cells and at almost all stages of life. This widespread use of Ca in intracellular signaling pathways requires tight regulation. Calcium release-activated calcium (CRAC) channels are a type of store-operated (SOC) channel that have been identified and described in detail, especially in immune cells. Activation of CRAC channels results from the depletion of Ca located in endoplasmic reticulum (ER) stores. Depletion of the ER stores signals the ER protein, Stim1 to aggregate into punctate structures near the plasma membrane where it clusters with the plasma membrane protein, Orai1. Formation of the Stim1/Orai1 complex activates CRAC channels allowing the selective entry of Ca into the cell. Alterations in the activation and assembly of Stim1 and Orai1 have serious consequences on human health. DCPA is a post-emergent herbicide used in rice fields and numerous studies suggest that it has negative effects on the immune system. We have previously reported that DCPA inhibits Ca influx in Jurkat T cells with adverse effects on IL-2 production and NFAT activity. Here we examine the effects of DCPA on Stim1 puncta formation. Upon ER Ca store depletion, HEK293 cells expressing YFP-Stim1 formed puncta-rich regions near the plasma membrane of the cell. DCPA-treated cells formed few punctate structures and failed to form the puncta-rich regions when ER stores were depleted. Here we propose a mechanism and identify DCPA as an inhibitor of Stim1 puncta formation.

INTRODUCTION

Calcium signaling

Calcium (Ca) is a ubiquitous second messenger that is important in all aspects of life, from conception to death. Many signaling systems are used to regulate the flow of Ca into and out of a cell. Store-operated Ca entry (SOCE) is used by virtually all animal cells to increase intracellular Ca levels in response to a stimulus (Feske 2009). As the name implies, entry of Ca into the cell is regulated by the levels of free Ca stored in the lumen of the ER. SOCE is initiated by stimulation of receptors on the surface of a cell which leads to the production of IP₃. IP₃ binds to the IP₃ receptor on the surface of the ER resulting in the release of Ca from the ER stores. Stim1, an ER protein with an EF-hand domain extending into the lumen of the ER, acts as a Ca sensor to detect changes in Ca. Multimerization of Stim1 initiates interactions with Orai1 proteins, located in the plasma membrane, and results in the activation of CRAC channels (Fig 2). The formation of Stim1-Orai1 clusters is essential for CRAC activation and translocation of Stim1 to areas within 10-25nm of the membrane occurs within 6-10 seconds prior to CRAC activation (Wu, Buchanan et al. 2006). Opening of CRAC channels enables the refilling of ER Ca stores and functions to produce the sustained and elevated Ca signaling required for activation of various signaling pathways. In T cells, influx of Ca through CRAC channels and the subsequent activation of the Ca-dependent transcription factor NFAT are critical for the expression of IL-2, an important early cytokine responsible for activation, proliferation and differentiation of several immune cells (Vig and Kinet 2009).

Stromal Interaction Molecule-1 (Stim1)

Stim1 is a single transmembrane protein that has been identified as an essential protein in SOC signaling in many cell types. Originally described in 1996 this protein, then designated SIM, was identified in a screening for proteins that bind B cell precursors (Oritani and Kincade 1996). The gene for this protein was cloned from the human chromosome region 11p15.5 and is evolutionarily conserved from *Drosophila* to mammals. Further studies revealed that Stim1 had tumor suppressor abilities in some cancer cell lines (Sabbioni, Barbanti-Brodano et al. 1997; Sabbioni, Veronese et al. 1999). The link between SOCE and Stim1 came in 2005 when Roos et al discovered that, in siRNA against Stim, in *Drosophila*, decreased Ca influx (Roos, DiGregorio et al. 2005). Simultaneously, Liou et al screened 2304 human genes using siRNA for each gene and looked for those altering Ca influx in HeLa cells (Liou, Kim et al. 2005). They identified 2 genes, Stim1 and Stim2. Stim1 and Stim2 are ubiquitously expressed in almost all mammalian tissue and, to date, the role of Stim1 in SOCE has been identified in numerous cell types including T cells, B cells, macrophages, mast cells, platelets, skeletal muscle, smooth muscle, vascular endothelium and neuronal cells (Soboloff, Spassova et al. 2006; Frischauf, Schindl et al. 2008).

Stim1 is characterized as a 685 amino acid, 90k Da, Type I transmembrane domain protein (Williams, Manji et al. 2001) (Fig 1). The N terminus resides in the lumen of the ER and the C-terminus in the cytosol. The N terminus is a highly conserved region that contains a single sterile alpha motif (SAM) domain that is modified by N-linked glycosylation at 2 sites (Asn131 and Asn171). The SAM domain is thought to mediate several hetero- and homotypic interactions. Deletion of the SAM

domain results in the failure of Stim1 to form puncta and CRAC channels are not activated (Baba, Hayashi et al. 2006). In addition, the presence of an EF-hand domain, in the N terminus acts as a Ca sensor for ER Ca stores. A single mutation (Asp76Ala) in the EF-hand domain of Stim1 results in prelocalized puncta formation in the absence of ER Ca store depletion along with constitutive activation of the CRAC current (I_{CRAC}) (Liou, Kim et al. 2005; Zhang, Yu et al. 2005).

The C-terminus of Stim1 is located in the cytosol and contains 2 highly conserved alpha helical coiled-coil regions, an erzin-radixin-moesin (ERM) domain and Ser and Lys-rich regions (Stathopoulos, Li et al. 2006). Within the ERM domain, a Stim/Orai-activating region (SOAR) and a Stim1 homomerization domain (SHD) have recently been identified (Muik, Fahrner et al. 2009; Yuan, Zeng et al. 2009). Deletion of the ERM domain disrupts Stim1 puncta formation and CRAC activation. Further mutational studies have identified a CRAC-activating domain (CAD), also called Orai-activating small fragment (OASF), within the SOAR domain (Muik, Fahrner et al. 2009; Park, Hoover et al. 2009). Expression of CAD alone is sufficient to activate Orai1 and deletion of CAD results in diffuse Stim1 expression when ER Ca stores are depleted (Park, Hoover et al. 2009). Stim1 also contains a CRAC modulatory domain (CMD), C terminal to the SHD, which is reported to regulate the degree of coupling to Orai1 as well as CRAC activation (Derler, Fahrner et al. 2009). The extreme C terminus of Stim1 contains a polybasic region which can bind calmodulin, in a calcium-dependent manner, and is critical for translocation to the plasma membrane (Bauer, O'Connell et al. 2008).

Figure 1. Domain structures in human Stim1 and Stim2

It is estimated that 20-30% of Stim1 is expressed on the plasma membrane but limited and conflicting data is available to determine its function (Manji, Parker et al. 2000). Some studies have identified a role for plasma membrane bound Stim1 in platelet aggregation and vascular smooth muscle cell migration (Jardin, Lopez et al. 2009) (Li, Sukumar et al. 2008). In addition, a Stim1 antibody, applied externally, blocks I_{CRAC} in Jurkat T cells and SOCE in HEK293 cells, revealing that Stim1 on the plasma membrane may be functionally important (Soboloff, Spassova et al. 2006; Spassova, Soboloff et al. 2006). However, Liou et al reported did not detect Stim1 on the plasma membrane of HeLa cells, indicating the role of Stim1 on the plasma membrane may be cell type specific (Liou, Fivaz et al. 2007).

Knockdown and mutational studies of Stim1

Numerous knockdown and mutational studies have contributed to the understanding of Stim1 in SOCE. Knockdown of Stim1 in several cell types, as well as in *Drosophila*, result in inhibition of I_{CRAC} (Ong, Liu et al. 2007; Lyfenko and Dirksen

2008; Picard, McCarl et al. 2009). Overexpression of Stim1 and Orai1, in HEK293 and Jurkat T cells, greatly increased I_{CRAC} with a 10-60 fold increase in Ca influx but with no change in basal resting Ca levels (Soboloff, Spassova et al. 2006; Soboloff, Spassova et al. 2006). However, overexpression of Orai1 alone does not alter I_{CRAC} , but overexpression of Stim1 increased I_{CRAC} 2 fold, indicating an important role for Stim1 in CRAC activation (Vig, Peinelt et al. 2006).

Stim1^{-/-} mice die *in utero* or shortly after birth but studies in mice with conditionally targeted deletion of *Stim1* have been reported (Oh-Hora, Yamashita et al. 2008). Conditional deletion of Stim1 in T cells and fibroblasts showed almost no Ca²⁺ influx in response to depletion of ER Ca²⁺ stores. In addition, TCR-stimulated production of IL-2, IFN- γ , and IL-4 production is significantly decreased in T cells from Stim1^{-/-} mice (Oh-Hora, Yamashita et al. 2008).

Recently, three individuals from a single family were identified with a homozygous nonsense mutation in Stim1. These individuals have no mutations in Orai but have decreased SOCE and profound immunodeficiencies. *Ex vivo* studies demonstrated that SOCE could be partially rescued by introducing functional Stim1 but only partially rescued with Stim2 (Picard, McCarl et al. 2009).

Stim2

Stim2 was originally identified by Williams et al during a screening for Stim1-like sequences (Williams, Manji et al. 2001). Differential levels of phosphorylation account for two molecular mass isoforms (105kDa and 115kDa). Stim2 is ubiquitously expressed and although Stim1 and Stim2 have high sequence homology, there are several important differences that may account for differences in function (Fig 1)

(Williams, Manji et al. 2001). The N-termini of both proteins are almost identical, with the noted exception that Stim1 has 2 N-linked glycosylation sites in the SAM domain and Stim2 only has one site. The greatest differences lie within the cytosolic C-terminal domain. As mentioned previously, Stim1 is expressed in the ER and the plasma membrane, however, an ER retention sequence in the C-terminus of Stim2, but not on Stim1, confines its expression to the ER (Wang, Deng et al. 2009). In addition, the C-terminus of Stim2 contains phosphorylated Ser/His residues and a large Pro-rich domain.

In vitro and *in vivo* experiments have demonstrated that Stim2 can oligomerize with Stim1, indicating a possible functional interaction between Stim1 and Stim2 (Stathopoulos, Zheng et al. 2009; Wang, Deng et al. 2009). Overexpression of Stim2 does not result in puncta formation when ER stores are depleted but puncta formation does occur when both Stim1 and Stim2 are overexpressed (Soboloff, Spassova et al. 2006). In HEK 293 and Jurkat T cells, overexpression of Stim2 alone decreased Ca influx and I_{CRAC} but knockdown of Stim2, using siRNA, has little effect on Ca influx or I_{CRAC} (Dziadek and Johnstone 2007). Interestingly, overexpression of both Stim2 and Orai1 slightly increase baseline I_{CRAC} when ER stores were filled and addition of 50 μ M 2-APB, normally inhibitory, increased I_{CRAC} when Stim2 and Orai1 are overexpressed. (Parvez, Beck et al. 2008) This data suggests that Stim2 acts as a regulator of SOC signaling. Further studies revealed that expression of the mutant Stim1(D76A) forms puncta when ER stores are full and produced constitutive Ca entry but when co-expressed with Stim2, puncta formation and constitutive Ca entry could be reversed (Brandman, Liou et al. 2007). This suggests that Stim2 can interfere with Stim1-

mediated SOC at a point downstream of puncta formation. It has been proposed that the effects of Stim2 are dependent on its ratio with Stim1 and that after store depletion, Stim1 and Stim2 aggregate and organize into puncta (Soboloff, Spassova et al. 2006). If the Stim1-Stim2 ratio is high, functional coupling to Orai1 occurs and SOCE proceeds but if the ratio is low, Stim2 interferes with coupling and inhibits SOCE.

There are also some studies that suggest that Stim2 is involved in stabilizing basal cytosolic and ER Ca levels (Brandman, Liou et al. 2007). Knockdown of Stim2 in HEK293 cells reduced basal cytosolic and ER Ca levels whereas overexpression increased basal Ca levels. Similar studies have reported that Stim2 responds to changes in ER Ca near resting levels, whereas Stim1 is activated only when ER Ca stores are depleted (Frischauf, Schindl et al. 2008).

Orai / CRACM

In 2006, Feske et al identified a protein in patients with a hereditary form of severe combined immunodeficiency (SCID) syndrome whose T cells failed to activate CRAC channels (Feske 2009). They named this protein Orai1, after the Greek mythological Orai, keepers of heaven's gate. At the same time Vig et al identified the same protein through a siRNA screening of the Drosophila genome for proteins that altered Ca influx and designated the protein CRACM1 (Vig, Peinelt et al. 2006). Orai1 or CRACM1 is distributed diffusely in the plasma membrane at rest but aggregates and interacts with Stim1 in overlapping clusters after store depletion to activate CRAC channels (Fig 2) (Vig, Beck et al. 2006; Navarro-Borelly, Somasundaram et al. 2008). Förster resonance energy transfer (FRET) analysis confirmed that Stim1 and Orai1 co-immunoprecipitate and overlapping puncta occur in parallel with increases in FRET

(Navarro-Borelly, Somasundaram et al. 2008). The coupling of Stim1/Orai1 to activate CRAC channels is dynamic but fully reversible when ER stores are replenished (Mignen, Thompson et al. 2008).

The Orai family consists of 3 members, Orai1, Orai2 and Orai3. A functional CRAC channel requires the formation of an Orai tetramer. The exact contribution from each family member has not been fully elucidated but appears to be dependent on cell type (Mignen, Thompson et al. 2008). Orai1/CRACM1 is the dominant form expressed in most mammalian cells. It is a 33kDa, 301 amino acid protein with 4 transmembrane (TM) spanning domains with both N and C termini located in the cytosol. Orai1 has been shown to form the essential pore-forming unit of the CRAC channel (Prakriya, Feske et al. 2006; Feske 2010). The N terminus of Orai1 contains a Pro/Arg-rich region (not found in Orai2 or Orai3) with a conserved polybasic (Arg/Lys) motif before TM1. The Pro/Arg-rich region is thought to play a role in Orai1 assembly. The C-terminus region contains a coiled-coil domain and protein interaction domains. Deletion of the complete N terminus in Orai1 abolishes Ca²⁺ influx but co-clustering interaction with Stim1 is retained, indicating that the C-terminus contains regions important in the interaction with Stim1 (Li, Lu et al. 2007; Muik, Frischauf et al. 2008).

Feske et al Pflugers Arch 2010

Figure 2. Schematic of Stim1-mediated Orai1 activation of CRAC channels

The transmembrane (TM) regions are important in forming the CRAC channel pore. Based on mutational analysis, TM1 and TM3 and the extracellular loop between TM1 and TM2 form the CRAC channel pore (Vig, Beck et al. 2006; Yeromin, Zhang et al. 2006). A Glu106Asp mutation in TM1 and a Glu190Gln mutation in TM3 both decrease the Ca selectivity of the pore through a proposed mechanism that increases the size of the pore (Yeromin, Zhang et al. 2006; Yamashita, Navarro-Borely et al.

2007). As p to A la mutations i n am ino acids 110, 112, and 114 i n t he TM 1-TM2 extracellular loop also alters ion selectivity (Yeromin, Zhang et al. 2006).

Interaction with Stim1 is thought to be mediated through an amphipathic domain on the C terminus of Orai1 and also through two hydrophobic Lys residues. Isolation of a 107 amino acid region, within the coiled-coil domain of the C-terminus of Stim1, can co-immunoprecipitate with Orai1 and activate CRAC channels independent of the ER Ca stores (Wang, Deng et al. 2009).

In HEK 293 and Jurkat T cells, overexpression of both Stim1 and Orai1 produce an extremely large I_{CRAC} but overexpression of Orai1 alone produces no increase in I_{CRAC} (Mercer, Dehaven et al. 2006; Soboloff, Spassova et al. 2006; Vig, Beck et al. 2006). Knockdown of Orai1 abolishes Ca influx and CRAC activity (Vig, Beck et al. 2006). Expression of Orai1 and Stim2 increased cell proliferation of HEK 293 cells and endothelial cells, with no evident role for Stim1 (Potier, Gonzalez et al. 2009; El Boustany, Katsogiannou et al. 2010). Silencing Orai1 or Stim2 abolished proliferation but silencing Stim1 had no effect.

Orai2/CRACM2 and Orai3/CRACM3

Orai1 is the dominant form of Orai expressed in most cells lines, however, Orai2 and Orai3 can also produce I_{CRAC} upon co-expression with Stim1 (Lis, Peinelt et al. 2007). In HEK 293 cells, overexpression of both Stim1 and Orai2 increased Ca influx and I_{CRAC} but to a lesser degree than in cells overexpressing both Stim1 and Orai1 (Soboloff, Spassova et al. 2006). In addition, overexpression of both Stim1 and Orai2 has no effect in T cells from SCID patients (Feske 2010). When Stim1 and Orai3 are overexpressed, Ca influx did not increase (Lis, Peinelt et al. 2007). Chimeric constructs

of Orai2 with the Orai1 C-terminus increases I_{CRAC} , and deletion of part of N-term of Orai1 results in a diminished Ca influx, comparable to that of wild type Orai2 and Orai3 (Li, Lu et al. 2007; Yuan, Zeng et al. 2009). In addition, four Orai1 proteins are thought to homodimerize to form an active $CRAC$ channel but, based on co-immunoprecipitation experiments, heteromultimerization can also occur between Orai2 and Orai3 (Penna, Demuro et al. 2008; Maruyama, Ogura et al. 2009). These studies suggest multiple functions for Orai proteins which may be cell-type specific.

MATERIALS AND METHODS

Cell lines

Experiments were performed using human embryonic kidney-293 (HEK-293) cells (a generous gift from J. Soboloff). HEK 293 cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Mediatech, Herndon, VA) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS) (Hyclone, Logan, UT), 100 units/ml penicillin (BioWhittaker, Walkersville, MD), 100 μ g/ml streptomycin (BioWhittaker), 2.0 mM glutamine (BioWhittaker). The cultures were kept at 37°C in 5% CO₂.

Reagents

Stock solutions of DCPA (ChemServices, West Chester, PA) were diluted in absolute ethanol (AAPER Alcohol and Chemical Company, Shelbyville, KY). Vehicle samples were treated with an equivalent concentration (0.1%v/v) of ethanol.

Transfections

The day prior to transfection, 0.7×10^5 HEK 293 cells were added to Delta T culture dishes (Bioprotech, Butler, PA) in DMEM supplemented with 10% FBS, without antibiotics and incubated overnight at 37°C in 5% CO₂. The day of transfection, HEK 293 cells were transfected using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) as recommended by the manufacturer. Briefly, for each plate 1 μ l of Lipofectamine 2000 was placed in 50 μ l of OptiMem (Invitrogen) and incubated at room temperature for 5 minutes. The transfection solution was added to a tube with 50 μ l OptiMem and 0.5 μ g YFP-Stim1 plasmid (a generous gift from J. Soboloff) and incubated at room temperature for 20 minutes. The DNA-transfection complex was added to the HEK 293 cells and incubated at 37°C in 5% CO₂ for 5 hours, after which the media was replaced

with fresh, complete DMEM and incubate for 24 hours at 37°C in 5% CO₂. On the day of the experiment the cells were placed in buffer containing 140 mM NaCl, 5 mM KCl, 1 mM MgCl₂, 10 mM glucose, 15 mM Hepes, 0.1% BSA and 2 mM CaCl₂. Cells were kept at 37°C throughout the experiment and reagents added to the plate as indicated. YFP-Stim1 images were acquired using a 60x/1.49 Apo TIRF objective and 1x1 binning on a Nikon Eclipse TE2000-E inverted fluorescent microscope equipped with the Perfect Focus system and a Photometrics CoolSNAP HQ CCD camera. A z-stack of fluorescent images was collected at 1 µm intervals every 20 seconds for 10 minutes. Cells were maintained at 37°C with a Biotechs Delta T dish heater. Images were acquired and analyzed using the NIS-Elements software package. Images shown are from a single focal plane.

RESULTS

In T cells exposure to DCPA leads to decreased IL-2 production, NFAT activity and calcium influx. Since Ca influx in T cells is regulated by CRAC channels we sought to determine the effects of DCPA on Stim1 puncta formation. Characterization of the role of Stim1 in CRAC channel activation has been thoroughly investigated in both Jurkat and HEK 293 cells (Parekh 2006; Feske 2007; Prakriya 2009). In both cell types, Stim1 puncta formation, its interaction with Orai1 and activation of CRAC channels all occur in a similar manner. HEK 293 cells are adherent, easy to transfect cells that have a relatively large cytoplasmic space that allows easy identification of puncta formation. These cells were used to determine the effects of DCPA on Stim1 puncta formation. HEK 293 cells were transfected with a YFP-Stim1 plasmid and ER Ca stores were depleted with 2 μ M thapsigargin approximately 70-80 s after data collection began (Fig 3A). Cells treated with DCPA received 100 μ M DCPA just prior to the addition of 2 μ M thapsigargin (Tg). Images were collected over a 16 minute period. Prior to treatment, all cells displayed a diffuse distribution of YFP-Stim1. Upon depletion of ER Ca stores with Tg in control cells, puncta formation was observed after approximately 7.5 minutes and continued to form over time (white arrows indicate areas of puncta formation). Puncta-rich regions are apparent around the membranes of the cells. Upon ER store depletion, cells treated with DCPA appeared to form some puncta but fail to form the puncta-rich regions observed in control cells (Fig.3B). These results suggest that DCPA can block Ca influx through CRAC channels by inhibiting Stim1 puncta formation.

Figure 3A Formation of Stim1 puncta-rich regions in control cells

HEK 293 cells were transfected with YFP-Stim1 and store-operated Ca^{2+} influx was initiated by depletion of ER Ca stores with $2\mu M$ Tg. Tg was added to control cells at $t=70$ sec. The cells presented here represent an average response of 30-40 cells.

Figure 3B DCPA inhibits formation of Stim1 puncta-rich regions

HEK 293 cells were transfected with YFP-Stim1 and store-operated Ca influx was initiated by depletion of ER Ca stores with $2\mu\text{M}$ Tg. $100\mu\text{M}$ DCPA was added prior to Tg at $t=67$ sec. The cells presented here represent an average response of 30-40 cells.

DISCUSSION

Regulation of intracellular calcium levels in cells is essential for almost all aspects of cell function, and alterations in calcium homeostasis can have profound effects on cell function and viability. Many cells, including HEK 293 and Jurkat T cells, use SOC channels to regulate Ca influx (Peinelt, Vig et al. 2006; Vig and Kinet 2009). SOCE is a form of Ca influx that is dependent on the depletion of its ER Ca stores in order to activate Ca channels on the plasma membrane. CRAC channels are the most widely known and described SOC channel and were identified over 20 years ago (Vig and Kinet 2007; Prakriya 2009; Vig and Kinet 2009). Until recently, CRAC channels could only be characterized by electrophysiological methods. The mechanism by which CRAC channels function has become clearer with the identification of 2 essential proteins, Stim1 and Orai1. Stim1 is a single transmembrane protein that predominantly resides in the ER, with some studies indicating a role for Stim1 in the plasma membrane (Baba and Kurosaki 2009). It contains an EF-hand domain that acts as a Ca sensor in the lumen of the ER. In unstimulated cells, both Stim1 and Orai1 proteins are diffusely distributed in the ER and plasma membrane, respectively (Liou, Kim et al. 2005). Depletion of ER Ca stores releases Ca from the Stim1 EF-hand and induces a conformational change that facilitates aggregation of Stim1 and results in the formation of overlapping clusters with Orai1, in areas close to the plasma membrane (Luik, Wang et al. 2008). This Stim1/Orai1 complex activates CRAC channels and allows Ca to flow into the cell (Fahrner, Muik et al. 2009). Once inside the cell, Ca plays an important role in activating multiple signaling pathways, including the NFAT (Gwack, Feske et al. 2007).

Alterations in Stim1 can have profound effects on cell function and, in T cells, can result in severe immunodeficiencies (Feske 2009). Recently three individuals, from a single family, were identified with a homozygous nonsense mutation in Stim1 but with no mutations in Orai1 (Picard, McCarl et al. 2009). These patients have profound immunodeficiencies and limited T cell SOCE. In addition, several other *in vitro* and *in vivo* studies support the important role for Stim1 in SOCE and calcium signaling (Feske, Prakriya et al. 2005; Feske 2010). Knockdown of Stim1 in several cell types, including Hek 293 and Jurkat T cells, resulted in inhibition of I_{CRAC} whereas overexpression of Stim1 increased I_{CRAC} 2 fold (Liou, Kim et al. 2005). *In vivo* experiments, with a conditional knockdown of Stim1 in mice, revealed almost no Ca^{2+} influx in response to depletion of Ca stores (Oh-Hora, Yamashita et al. 2008). TCR-stimulated CD4⁺ T cells from these mice also produce significantly less IL-2, IFN- γ , and IL-4.

Inhibitors of SOCE have been described but most have proven non-specific for CRAC channels (DeHaven, Smyth et al. 2008; Nam, Shin et al. 2009; Salido, Sage et al. 2009). A widely used modulator of CRAC channels is 2-APB (Bootman, Collins et al. 2002). It is a bimodal modulator that activates CRAC at low concentrations (<5 μ M) and inhibits CRAC at higher concentrations (>10 μ M) (DeHaven, Smyth et al. 2008). We have previously reported that DCPA inhibits Ca^{2+} influx in Jurkat T cells and attenuates the enhanced Ca^{2+} influx observed with low concentrations of 2-APB (Lewis, Brundage et al. 2008). Here we more clearly define the mechanism by which DCPA alters Ca^{2+} influx. HEK 293 cells have a diffuse expression of YFP-Stim1 when cells are at rest and aggregate into punctate structures when ER Ca^{2+} stores are depleted. DCPA decreased the puncta-rich regions observed in control cells (Fig 3B). Some puncta do appear to

form in DCPA-treated cells but the dense puncta-rich regions observed in control cells were never observed. We conclude that DCPA inhibits Stim1 puncta formation in HEK 293 cells and extend the results here to conclude that DCPA also inhibits Stim1 puncta formation in Jurkat T cells. Both cells lines are known to express CRAC channels and overexpression and mutational studies in both cells lines respond in a similar manner. Our data, in Jurkat T cells, supports a mechanism whereby inhibition of Stim1 results in decreased Ca influx, NFAT activity and IL-2 production. In addition, CRAC channels are the only Ca channels expressed on T cells that respond to ER store depletion. A recent study examining the localization of DCPA in Jurkat T cells concluded that, after stimulation, DCPA is localized to the cytosol and not found in the membrane or subcellular fractions (Hanson, Peer et al. 2010). This data supports a role for a mechanism in which DCPA interact with Stim1 in the cytosol to inhibit puncta formation. An analog of 2-APB was recently identified as a potential inhibitor of Stim1 but the specificity of its actions are unknown (Goto, Suzuki et al. 2010). No other known inhibitors of Stim1 have been reported. DCPA may serve as a novel inhibitor of Stim1 with the potential for use as a tool to better understand Stim1/Orai1 interactions and CRAC channels.

REFERENCES

- Baba, Y., K. Hayashi, et al. (2006). "Coupling of STIM1 to store-operated Ca^{2+} entry through its constitutive and inducible movement in the endoplasmic reticulum." Proc Natl Acad Sci U S A **103**(45): 16704-9.
- Baba, Y. and T. Kurosaki (2009). "Physiological function and molecular basis of STIM1-mediated calcium entry in immune cells." Immunol Rev **231**(1): 174-88.
- Bauer, M. C., D. O'Connell, et al. (2008). "Calmodulin binding to the polybasic C-termini of STIM proteins involved in store-operated calcium entry." Biochemistry **47**(23): 6089-91.
- Bootman, M. D., T. J. Collins, et al. (2002). "2-aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca^{2+} entry but an inconsistent inhibitor of InsP3-induced Ca^{2+} release." Faseb J **16**(10): 1145-50.
- Brandman, O., J. Liou, et al. (2007). "STIM2 is a feedback regulator that stabilizes basal cytosolic and endoplasmic reticulum Ca^{2+} levels." Cell **131**(7): 1327-39.
- DeHaven, W. I., J. T. Smyth, et al. (2008). "Complex actions of 2-aminoethoxydiphenyl borate on store-operated calcium entry." J Biol Chem **283**(28): 19265-73.
- Derler, I., M. Fahrner, et al. (2009). "A Ca^{2+} release-activated Ca^{2+} (CRAC) modulatory domain (CMD) within STIM1 mediates fast Ca^{2+} -dependent inactivation of ORAI1 channels." J Biol Chem **284**(37): 24933-8.
- Dziadek, M. A. and L. S. Johnstone (2007). "Biochemical properties and cellular localisation of STIM proteins." Cell Calcium **42**(2): 123-32.
- El Boustany, C., M. Katsogiannou, et al. (2010). "Differential roles of STIM1, STIM2 and Orai1 in the control of cell proliferation and SOCE amplitude in HEK293 cells." Cell Calcium.
- Fahrner, M., M. Muik, et al. (2009). "Mechanistic view on domains mediating STIM1-Orai coupling." Immunol Rev **231**(1): 99-112.
- Feske, S. (2007). "Calcium signalling in lymphocyte activation and disease." Nat Rev Immunol **7**(9): 690-702.
- Feske, S. (2009). "ORAI1 and STIM1 deficiency in human and mice: roles of store-operated Ca^{2+} entry in the immune system and beyond." Immunol Rev **231**(1): 189-209.

- Feske, S. (2010). "CRAC channelopathies." Pflugers Arch.
- Feske, S., M. Prakriya, et al. (2005). "A severe defect in CRAC Ca²⁺ channel activation and altered K⁺ channel gating in T cells from immunodeficient patients." J Exp Med **202**(5): 651-62.
- Frischauf, I., R. Schindl, et al. (2008). "The STIM/Orai coupling machinery." Channels (Austin) **2**(4): 261-8.
- Goto, J., A. Z. Suzuki, et al. (2010). "Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca(2+) entry via STIM proteins." Cell Calcium **47**(1): 1-10.
- Gwack, Y., S. Feske, et al. (2007). "Signalling to transcription: store-operated Ca²⁺ entry and NFAT activation in lymphocytes." Cell Calcium **42**(2): 145-56.
- Hanson, M. L., C. J. Peer, et al. (2010). "Subcellular localization of the amide class herbicide 3,4-dichloropropionanilide (DCPA) in T cells and hepatocytes." J Toxicol Environ Health A **73**(1): 1-4.
- Jardin, I., J. J. Lopez, et al. (2009). "Store-operated Ca²⁺ entry is sensitive to the extracellular Ca²⁺ concentration through plasma membrane STIM1." Biochim Biophys Acta **1793**(10): 1614-22.
- Lewis, T. L., K. M. Brundage, et al. (2008). "3,4-Dichloropropionanilide (DCPA) inhibits T-cell activation by altering the intracellular calcium concentration following store depletion." Toxicol Sci **103**(1): 97-107.
- Li, J., P. Sukumar, et al. (2008). "Interactions, functions, and independence of plasma membrane STIM1 and TRPC1 in vascular smooth muscle cells." Circ Res **103**(8): e97-104.
- Li, Z., J. Lu, et al. (2007). "Mapping the interacting domains of STIM1 and Orai1 in Ca²⁺ release-activated Ca²⁺ channel activation." J Biol Chem **282**(40): 29448-56.
- Liou, J., M. Fivaz, et al. (2007). "Live-cell imaging reveals sequential oligomerization and local plasma membrane targeting of stromal interaction molecule 1 after Ca²⁺ store depletion." Proc Natl Acad Sci U S A **104**(22): 9301-6.
- Liou, J., M. L. Kim, et al. (2005). "STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx." Curr Biol **15**(13): 1235-41.

- Lis, A., C. Peinelt, et al. (2007). "CRACM1, CRACM2, and CRACM3 are store-operated Ca²⁺ channels with distinct functional properties." Curr Biol **17**(9): 794-800.
- Luik, R. M., B. Wang, et al. (2008). "Oligomerization of STIM1 couples ER calcium depletion to CRAC channel activation." Nature **454**(7203): 538-42.
- Lyfenko, A. D. and R. T. Dirksen (2008). "Differential dependence of store-operated and excitation-coupled Ca²⁺ entry in skeletal muscle on STIM1 and Orai1." J Physiol **586**(Pt 20): 4815-24.
- Manji, S. S., N. J. Parker, et al. (2000). "STIM1: a novel phosphoprotein located at the cell surface." Biochim Biophys Acta **1481**(1): 147-55.
- Maruyama, Y., T. Ogura, et al. (2009). "Tetrameric Orai1 is a teardrop-shaped molecule with a long, tapered cytoplasmic domain." J Biol Chem **284**(20): 13676-85.
- Mercer, J. C., W. I. Dehaven, et al. (2006). "Large store-operated calcium selective currents due to co-expression of Orai1 or Orai2 with the intracellular calcium sensor, Stim1." J Biol Chem **281**(34): 24979-90.
- Mignen, O., J. L. Thompson, et al. (2008). "Orai1 subunit stoichiometry of the mammalian CRAC channel pore." J Physiol **586**(2): 419-25.
- Muik, M., M. Fahrner, et al. (2009). "A Cytosolic Homomerization and a Modulatory Domain within STIM1 C Terminus Determine Coupling to ORAI1 Channels." J Biol Chem **284**(13): 8421-6.
- Muik, M., I. Frischauf, et al. (2008). "Dynamic coupling of the putative coiled-coil domain of ORAI1 with STIM1 mediates ORAI1 channel activation." J Biol Chem **283**(12): 8014-22.
- Nam, J. H., D. H. Shin, et al. (2009). "Inhibition of store-operated Ca²⁺ entry channels and K⁺ channels by caffeic acid phenethyl ester in T lymphocytes." Eur J Pharmacol **612**(1-3): 153-60.
- Navarro-Borelly, L., A. Somasundaram, et al. (2008). "STIM1-Orai1 interactions and Orai1 conformational changes revealed by live-cell FRET microscopy." J Physiol **586**(Pt 22): 5383-401.
- Oh-Hora, M., M. Yamashita, et al. (2008). "Dual functions for the endoplasmic reticulum calcium sensors STIM1 and STIM2 in T cell activation and tolerance." Nat Immunol **9**(4): 432-43.

- Ong, H. L., X. Liu, et al. (2007). "Relocalization of STIM1 for activation of store-operated Ca(2+) entry is determined by the depletion of subplasma membrane endoplasmic reticulum Ca(2+) store." J Biol Chem **282**(16): 12176-85.
- Oritani, K. and P. W. Kincade (1996). "Identification of stromal cell products that interact with pre-B cells." J Cell Biol **134**(3): 771-82.
- Parekh, A. B. (2006). "On the activation mechanism of store-operated calcium channels." Pflugers Arch **453**(3): 303-11.
- Park, C. Y., P. J. Hoover, et al. (2009). "STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1." Cell **136**(5): 876-90.
- Parvez, S., A. Beck, et al. (2008). "STIM2 protein mediates distinct store-dependent and store-independent modes of CRAC channel activation." Faseb J **22**(3): 752-61.
- Peinelt, C., M. Vig, et al. (2006). "Amplification of CRAC current by STIM1 and CRACM1 (Orai1)." Nat Cell Biol **8**(7): 771-3.
- Penna, A., A. Demuro, et al. (2008). "The CRAC channel consists of a tetramer formed by Stim-induced dimerization of Orai dimers." Nature **456**(7218): 116-20.
- Picard, C., C. A. McCarl, et al. (2009). "STIM1 mutation associated with a syndrome of immunodeficiency and autoimmunity." N Engl J Med **360**(19): 1971-80.
- Potier, M., J. C. Gonzalez, et al. (2009). "Evidence for STIM1- and Orai1-dependent store-operated calcium influx through ICRAC in vascular smooth muscle cells: role in proliferation and migration." Faseb J **23**(8): 2425-37.
- Prakriya, M. (2009). "The molecular physiology of CRAC channels." Immunol Rev **231**(1): 88-98.
- Prakriya, M., S. Feske, et al. (2006). "Orai1 is an essential pore subunit of the CRAC channel." Nature **443**(7108): 230-3.
- Roos, J., P. J. DiGregorio, et al. (2005). "STIM1, an essential and conserved component of store-operated Ca²⁺ channel function." J Cell Biol **169**(3): 435-45.
- Sabbioni, S., G. Barbanti-Brodano, et al. (1997). "GOK: a gene at 11p15 involved in rhabdomyosarcoma and rhabdoid tumor development." Cancer Res **57**(20): 4493-7.

- Sabbioni, S., A. Veronese, et al. (1999). "Exon structure and promoter identification of STIM1 (alias GOK), a human gene causing growth arrest of the human tumor cell lines G401 and RD." Cytogenet Cell Genet **86**(3-4): 214-8.
- Salido, G. M., S. O. Sage, et al. (2009). "Biochemical and functional properties of the store-operated Ca²⁺ channels." Cell Signal **21**(4): 457-61.
- Soboloff, J., M. A. Spassova, et al. (2006). "Calcium signals mediated by STIM and Orai proteins--a new paradigm in inter-organelle communication." Biochim Biophys Acta **1763**(11): 1161-8.
- Soboloff, J., M. A. Spassova, et al. (2006). "STIM2 is an inhibitor of STIM1-mediated store-operated Ca²⁺ Entry." Curr Biol **16**(14): 1465-70.
- Soboloff, J., M. A. Spassova, et al. (2006). "Orai1 and STIM reconstitute store-operated calcium channel function." J Biol Chem **281**(30): 20661-5.
- Spassova, M. A., J. Soboloff, et al. (2006). "STIM1 has a plasma membrane role in the activation of store-operated Ca(2+) channels." Proc Natl Acad Sci U S A **103**(11): 4040-5.
- Stathopulos, P. B., G. Y. Li, et al. (2006). "Stored Ca²⁺ depletion-induced oligomerization of stromal interaction molecule 1 (STIM1) via the EF-SAM region: An initiation mechanism for capacitive Ca²⁺ entry." J Biol Chem **281**(47): 35855-62.
- Stathopulos, P. B., L. Zheng, et al. (2009). "Stromal interaction molecule (STIM) 1 and STIM2 calcium sensing regions exhibit distinct unfolding and oligomerization kinetics." J Biol Chem **284**(2): 728-32.
- Vig, M., A. Beck, et al. (2006). "CRACM1 multimers form the ion-selective pore of the CRAC channel." Curr Biol **16**(20): 2073-9.
- Vig, M. and J. P. Kinet (2007). "The long and arduous road to CRAC." Cell Calcium **42**(2): 157-62.
- Vig, M. and J. P. Kinet (2009). "Calcium signaling in immune cells." Nat Immunol **10**(1): 21-7.
- Vig, M., C. Peinelt, et al. (2006). "CRACM1 is a plasma membrane protein essential for store-operated Ca²⁺ entry." Science **312**(5777): 1220-3.

- Wang, Y., X. Deng, et al. (2009). "STIM protein coupling in the activation of Orai channels." Proc Natl Acad Sci U S A **106**(18): 7391-6.
- Williams, R. T., S. S. Manji, et al. (2001). "Identification and characterization of the STIM (stromal interaction molecule) gene family: coding for a novel class of transmembrane proteins." Biochem J **357**(Pt 3): 673-85.
- Wu, M. M., J. Buchanan, et al. (2006). "Ca²⁺ store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane." J Cell Biol **174**(6): 803-13.
- Yamashita, M., L. Navarro-Borelly, et al. (2007). "Orai1 mutations alter ion permeation and Ca²⁺-dependent fast inactivation of CRAC channels: evidence for coupling of permeation and gating." J Gen Physiol **130**(5): 525-40.
- Yeromin, A. V., S. L. Zhang, et al. (2006). "Molecular identification of the CRAC channel by altered ion selectivity in a mutant of Orai." Nature **443**(7108): 226-9.
- Yuan, J. P., W. Zeng, et al. (2009). "SOAR and the polybasic STIM1 domains gate and regulate Orai channels." Nat Cell Biol **11**(3): 337-43.
- Zhang, S. L., Y. Yu, et al. (2005). "STIM1 is a Ca²⁺ sensor that activates CRAC channels and migrates from the Ca²⁺ store to the plasma membrane." Nature **437**(7060): 902-5.

CHAPTER 5

GENERAL DISCUSSION

The overall goal of this dissertation was to determine the effects that DCPA and its metabolites have on T cell activation and function and to propose a possible mechanism to explain the effects. Numerous studies have been conducted on the immunotoxic effects of DCPA (reviewed in (Salazar, Ustyugova et al. 2008)). Earlier studies in mouse models and human cell lines, laid the foundation for a more detailed investigation into the effects of DCPA on T cells. In addition, it is known that DCPA is metabolized quickly in the liver of mammals, but little immunotoxic data is available on the effects of its metabolites (McMillan, Freeman et al. 1990). The first goal was to determine the effects of DCPA and its metabolites on human T cell function and activation by examining immune parameters and signaling events. Once this was determined, a possible mechanism for these effects was then investigated. The results of these experiments shed light on the importance of investigating metabolic products when evaluating the immunotoxicity of chemicals and reveal that the effects of DCPA on T cells may provide a novel mechanism for immunosuppression.

In T cells, IL-2 is one of the first cytokines to be expressed and its transcription is dependent on the cooperative DNA binding of AP-1, NFAT and NF- κ B (Jain, Loh et al. 1995; Serfling, Berberich-Siebelt et al. 2007). NFAT is a transcription factor that is strictly calcium-dependent. In T cells, activation of NFAT occurs through a series of complex signaling events that begin with the recognition of a foreign peptide, and co-stimulation of the CD28 receptor. ITAMs, located on the intracellular portion of the TCR, act as a staging area for the phosphorylation and recruitment of several kinases

and adapter proteins that form the immunological synapse. One important consequence of this is the recruitment of PLC- γ which hydrolyzes the membrane lipid PIP₂ resulting in the production of IP₃ and DAG. DAG remains in the membrane where it plays a role in the activation of Ras and PKC θ pathways. IP₃ is released into the cytosol and binds to the IP₃-R on the ER allowing the release of Ca from the internal ER stores. Depletion of the ER stores releases Ca from the EF-hand domain on Stim1 allowing aggregation of Stim1 and interaction with Orai1 near the plasma membrane (Luik, Wang et al. 2008). Clustering of Stim1 and Orai1 activate CRAC channels and allows the selective entry of Ca into the cell. This increase in intracellular Ca enables it to bind to sites on calmodulin (CaM). The Ca-CaM complex binds to the regulatory subunit of calcineurin, thereby activating its phosphatase activity. Dephosphorylation of NFAT, by calcineurin, exposes a nuclear localization sequence which leads to translocation of NFAT into the nucleus. NFAT-driven gene expression is highly dependent on a sustained and elevated Ca influx (Rao 2009). Patients with T cells that have altered CRAC function also display changes in NFAT activity and IL-2 production that results in a SCID-like syndrome (Feske, Gwack et al. 2006). In addition, IL-2 is important in T cell homeostasis and tolerance and deficiencies can impair T cell activation, proliferation and lead to age-dependent tolerance and anergy (Smith-Garvin, Koretzky et al. 2009).

Immunotoxic studies indicate that exposure of DCPA to C57/B6 mice resulted in thymic atrophy, decreased thymic weight and alterations in several thymocyte populations (Barnett and Gandy 1989; Blyler, Landreth et al. 1994). Mice treated with DCPA also have enlarged spleens and decreased IL-2 and IL-6 production in

splenocytes stimulated with con-A, a T cell mitogen (Zhao, Schafer et al. 1998). Similar decreases in IL-2 production, as well as mRNA levels, were reported in mouse (EL-4) and human T (Jurkat) cell lines (Zhao, Schafer et al. 1999; Brundage, Schafer et al. 2004). Additional studies confirmed that human Jurkat T cells exposed to DCPA altered c-jun protein levels as well as phosphorylation levels (Brundage, Schafer et al. 2004). AP-1 is a transcription factor that is important in T cells for the expression of IL-2 and is comprised of c-jun and c-fos. Reductions in c-jun resulted in the decreased DNA binding ability of AP-1 with downstream decreases in IL-2. Transcription of IL-2 is dependent on the cooperative binding of 3 transcription factors; AP-1, NFAT and NF- κ B (Jain, Loh et al. 1995). Alterations in the timing and binding of these transcription factors can have adverse effects on T cell function (Jain, Loh et al. 1995; Hogan, Chen et al. 2003).

Although immunological studies indicate that DCPA can alter T cell function, there is sparse data on the immunotoxic effects of its metabolites. In humans, DCPA is metabolized by hepatic acylamidases to produce DCA which can undergo further biotransformation to produce NOH-DCA and 6OH-DCA (McMillan, Leakey et al. 1990). Roberts et al examined DCPA self-poisoning patients and reported that the estimated half-life of DCPA in the body is 3.2 hours and that DCA was detectable and more persistent than DCPA (Roberts, Heilmair et al. 2009). No data is available on the half life of DCA in humans, but in rats, elimination of DCA is reported to occur after 3 days. In humans, DCA is detectable in both occupationally and non-occupationally exposed individual and DCA-Hb adducts are detectable in agricultural workers four months after the last application of DCPA (Pastorelli 1998; Wittke, Hajimiragha et al. 2001; Turci,

Barisano et al. 2006). This indicates that exposure to DCPA may result in an accumulation of DCA in the body long after DCPA had been metabolized. Toxicity studies on DCA are limited but there have been reports that DCA can alter reproductive, endocrine, and liver functions (Valentovic, Yahia et al. 1997; Bauer, Meyer et al. 1998; Zhang and Lin 2009). Limited immunotoxicity studies indicate that DCA may be as toxic as DCPA, as molar equivalent doses of DCPA and DCA increased spleen weight and size and decreased NK cell function (Barnett, Gandy et al. 1992). In contrast, myelotoxic effects are observed in mice exposed to DCPA, but not when exposed to DCA (Malerba, Castoldi et al. 2002). *In vitro* studies, using a murine T cell line, resulted in decreased IL-6 production when exposed to 50µM DCPA but IL-6 was only inhibited by DCA at higher concentration (300µM) (Zhang unpublished data).

NOH-DCA and 6OH-DCA are metabolites that have been documented in their ability to form met-Hb adducts (McMillan, McRae et al. 1990). Both of these oxidative metabolites can convert oxy-Hb to met-Hb, a form of Hb that does not bind or transport oxygen. Formation of met-Hb can lead to a serious medical condition known as methemoglobinemia. NOH-DCA is also a nephrotoxicant and can induce hemolytic anemia (McMillan, Bradshaw et al. 1991; Valentovic, Ball et al. 2001). 6OH-DCA is also nephrotoxic but does not induce hemolytic anemia. In addition, no immunotoxicity studies on either NOH-DCA or 6OH-DCA have been conducted.

The immunotoxic effects of DCPA and its metabolites on T cells are reported here for the first time. As a test for overall T cell function, IL-2 secretion was measured in Jurkat T cells exposed to DCPA and its metabolites. DCPA decreased IL-2

secretion, up to 78%, in a concentration dependent manner. Exposure of Jurkat T cells to DCA also decreased IL-2 secretion in a concentration dependent manner but only at higher concentrations (100 μ M and 200 μ M). A 78% decrease in IL-2 production observed with 100 μ M DCPA required 200 μ M DCA to achieve the same inhibition. This suggests that DCPA is a more potent inhibitor of IL-2 secretion. Metabolic transformation can increase the toxicity of a chemical and, in NOH-DCA treated Jurkat T cells, IL-2 levels are decreased significantly more than that of DCPA. Although 50 μ M NOH-DCA decreased IL-2 by 90%, 50 μ M DCPA only inhibited IL-2 by 50%, indicating that metabolism of DCPA produces a metabolite more toxic than its parent. In addition, NFAT activity is also inhibited by DCPA and its metabolites, although to varying degrees. Interestingly, although a concentration of 50 μ M NOH-DCA inhibited IL-2 secretion by 90%, NFAT activity is only reduced 2 fold. In contrast, 50 μ M DCPA inhibited IL-2 secretion by almost 50% but NFAT activity was decreased 2.5 fold. In addition, 100 μ M DCPA decreased IL-2 secretion by 78% with a 8-fold decrease in NFAT activity. These data suggest that NOH-DCA and DCPA may inhibit IL-2 production through different mechanisms. To support this hypothesis, DCPA and DCA were also shown to inhibit Ca influx in a concentration-dependent manner but again, DCPA appears to be a more potent inhibitor. However, even at the highest concentrations of 6OH-DCA (50 μ M) and NOH-DCA (50 μ M) Ca influx was not altered. The mechanism by which NOH-DCA and 6OH-DCA inhibit IL-2 production is currently unknown, however, measurements of NFAT activity and IL-2 secretion require incubation times of 5hrs and 24hrs, respectively, and calcium flux was measured immediately after adding the treatment. It is possible that NOH-DCA alters Ca

homeostasis at later time points and that may account for the decreases in NFAT and IL-2. It is also possible that NOH-DCA and 6OH-DCA alter T cell function through a different mechanism than DCPA and DCA. Further studies are required to determine the mechanism for these differential effects.

Several mechanisms could account for the Ca-dependent changes observed in Jurkat T cells exposed to DCPA and DCA. T cells express 2 plasma membrane potassium (K) channels, Kv1.3 and KCa, which are involved in regulation of Ca flux in T cells. Kv1.3 channels are voltage-gated and activated when the membrane potential is depolarized, whereas KCa channels are activated by increases in intracellular Ca. Both channels function to move K ions out of the cell to increase the driving force of Ca. Alterations that prevent the efflux of K ions reduce the Ca driving force resulting in a decrease in intracellular Ca. Using a membrane potential dye we determined that DCPA-induced decreases in Ca influx were not mediated by alterations in the membrane potential.

In T cells, CRAC channels are responsible for the Ca influx observed in activated cells. In T cells from patients with SCID syndrome, Feske et al reported that these T cells retained only 1-2% of the normal Ca influx observed in stimulated cells (Feske 2009). Using electrophysiological and molecular biological methods they determined that this defect was due solely to the absence of functional CRAC channels. 2-APB is a widely used modulator of CRAC channels. Low concentrations ($<5\text{ }\mu\text{M}$) of 2-APB enhance Ca influx through CRAC channels following store depletion without altering IP₃-mediated release of Ca from the ER (Prakriya and Lewis 2001). High concentrations of 2-APB ($>10\text{ }\mu\text{M}$) inhibit CRAC channels. We used 2-APB to further

understand the effects of DCPA in Ca signaling. We demonstrate that, at CRAC-enhancing concentrations of 2-APB (2.5 μ M), we could inhibit Ca influx with DCPA. This combined data supports our hypothesis that DCPA alters Ca influx through inhibition of CRAC channels.

The toxicity of some chlorine compounds has been well documented and toxicity is dependent on the number, structure and location of the chlorine(s) (Naumann 2000). Vinyl chloride, 1,2 dichloroethane, polychlorinated biphenyls, DDT and heptachlor are common chlorine products with known toxicities. DCPA is a small lipophilic compound with 2 chlorines at the 3 and 4 positions on the benzene ring. Notably, although DCPA is lipophilic, it has recently been reported that it becomes localized in the cytosolic fraction of T cells and hepatocytes (Hanson, Peer et al. 2010). The effect of the chlorines on its toxicity is unknown. We report that, substitution of the 2 chlorines, in DCPA and DCA, with fluorines, to produce DFPA and DFA, respectively, resulted in a complete reversal of all effects. This clearly establishes a mechanism whereby the presence of the chlorines alters T cell function. Several possibilities exist for the differential effects observed with chlorine and fluorine. First, fluorines are highly electronegative and act only as hydrogen acceptors whereas chlorine and other halogens act as both hydrogen acceptors and donors (Purser, Moore et al. 2008). This increase in electronegativity could alter the distribution of charge so that the fluorine analogs do not interact with its target in the same manner as DCPA and DCA. Second, trifluoro-substitution has been reported to increase lipophilicity but mono- or difluoro-substitution can to decrease it (Purser, Moore et al. 2008). Changes in the lipophilicity may also alter interactions with the target of DCPA. Since DCPA is

targeted to the cytosol, increases in lipophilicity may prevent access into the cytosol. Lastly, fluorines are similar in size to hydrogen and may not produce the 3-D confirmation required to elicit the effects of DCPA. Further studies are required to determine the mechanism of DCPA, in particular, how the position and location of chlorines alters T cell function.

Based on the data collected we hypothesized that the inhibition of Ca influx observed in T cells exposed to DCPA was due to alterations in CRAC channel activity. Only in recent years has it become possible to assess changes in CRAC activity by methods other than electrophysiology. The discovery of two critical proteins in CRAC activation, Stim1 and Orai1, allows for a more detailed investigation into possible inhibitors of CRAC channels. Stim1 is an ER transmembrane protein that aggregates into puncta when ER stores are depleted and interacts with Orai1 proteins in the plasma membrane to activate CRAC channels. Puncta formation can be visualized in cells that express YFP-tagged Stim1 and disruption of the puncta formation results in loss of CRAC activity and Ca influx (Liou, Fivaz et al. 2007; Hewavitharana, Deng et al. 2008). In the absence of Stim1, mouse CD4⁺ T cells produce little Ca influx when ER stores are depleted and TCR-stimulated T cells produce significantly less IL-2 (Oh-Hora, Yamashita et al. 2008). Knockdown of Stim1 in HEK 293 cells also results in decreased Ca influx and CRAC activity (Roos, DiGregorio et al. 2005). Here we report that HEK 293 cells, treated with DCPA, failed to produce significant puncta formation upon ER store depletion. Some puncta formation does occur in the DCPA-treated cells but the puncta rich regions observed in control cells are not observed. HEK 293 cells are adherent, easy to transfect cells and have a relatively large cytosolic space to

identify puncta formation so these were chosen for our studies. Characterization of the role of Stim1 and Orai1 proteins in CRAC channel activation has been thoroughly investigated in both HEK 293 and Jurkat T cells. In both cell types, mutational and overexpression studies reported that Stim1 puncta formation, its interaction with Orai1 and activation of CRAC channels all occur in a similar manner. We extend our results in HEK 293 cells and conclude that DCPA inhibits IL-2, NFAT and Ca²⁺ influx by inhibiting Stim1 puncta formation in Jurkat T cells. Localization of DCPA to the cytosolic fraction in Jurkat T cells supports the possibility that DCPA can interact directly with the cytosolic domain of Stim1 to block puncta formation. In the cytosol, the C terminus of Stim1 contains a CRAC-activating domain (CAD) that, when deleted, does not form Stim1 puncta (Park, Hoover et al. 2009). Although little is known about this domain, it is possible that DCPA could alter this interaction leading to the inhibition of Stim1 puncta. However, there are several other possibilities that may also explain these results. A reduction in the mobility of the plasma membrane hydrocarbon chains, in stimulated Jurkat T cells exposed to DCPA, has been reported and it may be possible for this to occur on the ER membrane and possibly inhibit Stim1 dimerization (Brundage, Barnett et al. 2003). Interaction with cytoskeletal proteins could also prevent Stim1 puncta formation. Some studies indicate a role for the cytoskeleton in Stim1 aggregation. Stim1 can colocalize with alpha-tubulin and disruption of microtubules with nocodazole resulted in a diffuse distribution of Stim1 (Smyth, DeHaven et al. 2007). Since DCPA inhibits Stim1 puncta formation, it seems unlikely that DCPA directly alters Orai1 as aggregation of Stim1 is independent of Orai1 clustering (Xu, Lu et al. 2006). In fact, Stim1 puncta formation has been shown to

occur before Orai1 aggregation. Further studies are required to determine the exact nature of DCPA inhibition of Stim1.

In conclusion, exposure of Jurkat T cells to DCPA and its metabolites alters T cell function through inhibition of IL-2 and NFAT. The parent compound, DCPA, and its metabolite, DCA, both inhibit T cell function by decreasing IL-2 secretion in a NFAT and Ca-dependent manner. DCPA appears to be a more potent inhibitor of IL-2 secretion as greater decreases in IL-2 are observed at lower concentrations of DCPA than that of DCA. N-OH-DCA almost eliminated IL-2 production but in a Ca-independent manner. The positioning and presence of chlorines on DCPA and DCA are critical in eliciting these toxic effects. Finally, DCPA appears to be a novel inhibitor of CRAC channels and its use as such may have widespread implications for immunotherapy.

REFERENCES

- Barnett, J. B. and J. Gandy (1989). "Effect of acute propanil exposure on the immune response of C57Bl/6 mice." Fundam Appl Toxicol **12**(4): 757-64.
- Barnett, J. B., J. Gandy, et al. (1992). "Comparison of the immunotoxicity of propanil and its metabolite, 3,4-dichloroaniline, in C57Bl/6 mice." Fundam Appl Toxicol **18**(4): 628-31.
- Bauer, E. R., H. H. Meyer, et al. (1998). "Application of an androgen receptor assay for the characterisation of the androgenic or antiandrogenic activity of various phenylurea herbicides and their derivatives." Analyst **123**(12): 2485-7.
- Blyler, G., K. S. Landreth, et al. (1994). "Selective myelotoxicity of propanil." Fundam Appl Toxicol **22**(4): 505-10.
- Brundage, K. M., J. B. Barnett, et al. (2003). "The amide class herbicide 3,4-dichloropropionanilide (DCPA) alters the mobility of hydrocarbon chains in T-lymphocyte but not macrophage membranes." J Toxicol Environ Health A **66**(23): 2253-65.
- Brundage, K. M., R. Schafer, et al. (2004). "Altered AP-1 (activating protein-1) activity and c-jun activation in T cells exposed to the amide class herbicide 3,4-dichloropropionanilide (DCPA)." Toxicol Sci **79**(1): 98-105.
- Feske, S. (2009). "ORAI1 and STIM1 deficiency in human and mice: roles of store-operated Ca²⁺ entry in the immune system and beyond." Immunol Rev **231**(1): 189-209.
- Feske, S., Y. Gwack, et al. (2006). "A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function." Nature **441**(7090): 179-85.
- Hanson, M. L., C. J. Peer, et al. (2010). "Subcellular localization of the amide class herbicide 3,4-dichloropropionanilide (DCPA) in T cells and hepatocytes." J Toxicol Environ Health A **73**(1): 1-4.
- Hewavitharana, T., X. Deng, et al. (2008). "Location and function of STIM1 in the activation of Ca²⁺ entry signals." J Biol Chem **283**(38): 26252-62.
- Hogan, P. G., L. Chen, et al. (2003). "Transcriptional regulation by calcium, calcineurin, and NFAT." Genes Dev **17**(18): 2205-32.

- Jain, J., C. Loh, et al. (1995). "Transcriptional regulation of the IL-2 gene." Curr Opin Immunol **7**(3): 333-42.
- Liou, J., M. Fivaz, et al. (2007). "Live-cell imaging reveals sequential oligomerization and local plasma membrane targeting of stromal interaction molecule 1 after Ca²⁺ store depletion." Proc Natl Acad Sci U S A **104**(22): 9301-6.
- Luik, R. M., B. Wang, et al. (2008). "Oligomerization of STIM1 couples ER calcium depletion to CRAC channel activation." Nature **454**(7203): 538-42.
- Malerba, I., A. F. Castoldi, et al. (2002). "In vitro myelotoxicity of propanil and 3,4-dichloroaniline on murine and human CFU-E/BFU-E progenitors." Toxicol Sci **69**(2): 433-8.
- McMillan, D. C., T. P. Bradshaw, et al. (1991). "Contribution of 3,4-dichlorophenylhydroxylamine in propanil-induced hemolytic anemia." Adv Exp Med Biol **283**: 343-5.
- McMillan, D. C., J. P. Freeman, et al. (1990). "Metabolism of the arylamide herbicide propanil. I. Microsomal metabolism and in vitro methemoglobinemia." Toxicol Appl Pharmacol **103**(1): 90-101.
- McMillan, D. C., J. E. Leakey, et al. (1990). "Metabolism of the arylamide herbicide propanil. II. Effects of propanil and its derivatives on hepatic microsomal drug-metabolizing enzymes in the rat." Toxicol Appl Pharmacol **103**(1): 102-12.
- McMillan, D. C., T. A. McRae, et al. (1990). "Propanil-induced methemoglobinemia and hemoglobin binding in the rat." Toxicol Appl Pharmacol **105**(3): 503-7.
- Naumann, K. (2000). "Influence of chlorine substituents on biological activity of chemicals: a review." Pest Manag Sci **56**(1): 3-21.
- Oh-Hora, M., M. Yamashita, et al. (2008). "Dual functions for the endoplasmic reticulum calcium sensors STIM1 and STIM2 in T cell activation and tolerance." Nat Immunol **9**(4): 432-43.
- Park, C. Y., P. J. Hoover, et al. (2009). "STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1." Cell **136**(5): 876-90.
- Pastorelli, R., Catenacci, G., Guanci, M., Fannelli, R., Valoti, E., Minoia, C., Airoldi, L (1998). "3,4 Dichloroaniline-haemoglobin adducts in humans: preliminary data on agricultural workers exposed to propanil." Biomarkers **3**(3): 227-233.

- Prakriya, M. and R. S. Lewis (2001). "Potentiation and inhibition of Ca^{2+} release-activated Ca^{2+} channels by 2-aminoethyldiphenyl borate (2-APB) occurs independently of IP(3) receptors." J Physiol **536**(Pt 1): 3-19.
- Purser, S., P. R. Moore, et al. (2008). "Fluorine in medicinal chemistry." Chem Soc Rev **37**(2): 320-30.
- Rao, A. (2009). "Signaling to gene expression: calcium, calcineurin and NFAT." Nat Immunol **10**(1): 3-5.
- Roberts, D. M., R. Heilmair, et al. (2009). "Clinical outcomes and kinetics of propanil following acute self-poisoning: a prospective case series." BMC Clin Pharmacol **9**: 3.
- Roos, J., P. J. DiGregorio, et al. (2005). "STIM1, an essential and conserved component of store-operated Ca^{2+} channel function." J Cell Biol **169**(3): 435-45.
- Salazar, K. D., I. V. Ustyugova, et al. (2008). "A review of the immunotoxicity of the pesticide 3,4-dichloropropionanilide." J Toxicol Environ Health B Crit Rev **11**(8): 630-45.
- Serfling, E., F. Berberich-Siebelt, et al. (2007). "NFAT in lymphocytes: a factor for all events?" Sci STKE **2007**(398): pe42.
- Smith-Garvin, J. E., G. A. Koretzky, et al. (2009). "T cell activation." Annu Rev Immunol **27**: 591-619.
- Smyth, J. T., W. I. DeHaven, et al. (2007). "Role of the microtubule cytoskeleton in the function of the store-operated Ca^{2+} channel activator STIM1." J Cell Sci **120**(Pt 21): 3762-71.
- Turci, R., A. Barisano, et al. (2006). "Determination of dichloroanilines in human urine by gas chromatography/mass spectrometry: validation protocol and establishment of Reference Values in a population group living in central Italy." Rapid Commun Mass Spectrom **20**(17): 2621-5.
- Valentovic, M., J. G. Ball, et al. (2001). "3,4-Dichlorophenylhydroxylamine cytotoxicity in renal cortical slices from Fischer 344 rats." Toxicology **162**(3): 149-56.
- Valentovic, M. A., T. Yahia, et al. (1997). "3,4-Dichloroaniline acute toxicity in male Fischer 344 rats." Toxicology **124**(2): 125-34.

- Wittke, K., H. Hajimiragha, et al. (2001). "Determination of dichloroanilines in human urine by GC-MS, GC-MS-MS, and GC-ECD as markers of low-level pesticide exposure." J Chromatogr B Biomed Sci Appl **755**(1-2): 215-28.
- Xu, P., J. Lu, et al. (2006). "Aggregation of STIM1 underneath the plasma membrane induces clustering of Orai1." Biochem Biophys Res Commun **350**(4): 969-76.
- Zhang, B. and S. Lin (2009). "Effects of 3,4-dichloroaniline on testicle enzymes as biological markers in rats." Biomed Environ Sci **22**(1): 40-3.
- Zhao, W., R. Schafer, et al. (1998). "Cytokine production by C57BL/6 mouse spleen cells is selectively reduced by exposure to propanil." J Toxicol Environ Health A **55**(2): 107-20.
- Zhao, W., R. Schafer, et al. (1999). "Propanil affects transcriptional and posttranscriptional regulation of IL-2 expression in activated EL-4 cells." Toxicol Appl Pharmacol **154**(2): 153-9.