2014

Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones, total synthesis of methyl linderone and Wacker-type oxidation of aryl-substituted alkenes

Hong Yin

Follow this and additional works at: <https://researchrepository.wvu.edu/etd>

Recommended Citation

Yin, Hong, "Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones, total synthesis of methyl linderone and Wacker-type oxidation of aryl-substituted alkenes" (2014).

Graduate Theses, Dissertations, and Problem Reports. 7008.

<https://researchrepository.wvu.edu/etd/7008>

This Dissertation is protected by copyright and/or related rights. It has been brought to you by the The Research Repository @ WVU with permission from the rights-holder(s). You are free to use this Dissertation in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you must obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This Dissertation has been accepted for inclusion in WVU Graduate Theses, Dissertations, and Problem Reports collection by an authorized administrator of The Research Repository @ WVU. For more information, please contact researchrepository@mail.wvu.edu.

**Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or
2-halo-1,4-benzoquinones, total synthesis of methyl linderone and
Wacker-type oxidation of aryl-substituted alkenes**

Hong Yin

Dissertation submitted
to the Eberly College of Arts and Sciences
at West Virginia University
in partial fulfillment of the requirement for the degree of
Doctor of Philosophy in
Chemistry

Björn C. G. Söderberg, Ph. D., Chair
Kung K. Wang, Ph. D.
Alan M. Stolzenberg, Ph. D.
John H. Penn, Ph. D.
Patrick S. Callery, Ph. D.
C. Eugene Bennett Department of Chemistry

Morgantown, West Virginia
2014

Keywords: alkynylcyclobutenones, rearrangement, 2-alkylidene-1,3-cyclopentendiones, *N*-halosuccinimide, methyl linderone, aryl-substituted alkene, Wacker oxidation, α -arylated ketone

Copyright 2014 Hong Yin

ABSTRACT

Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones, total synthesis of methyl linderone and Wacker-type oxidation of aryl-substituted alkenes

Hong Yin

We have discovered a practical general method for the preparation of diverse 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones. An in-depth study of the reactions of 4-(ethynyl)-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones with *N*-bromo- or *N*-iodosuccinimide is described. This reaction was used in a short total synthesis of methyl linderone. Also, a series of facile, palladium catalyzed, direct, and regiospecific Wacker-type oxidations has been developed to yield various α -arylated ketones from aryl-substituted internal alkenes.

Acknowledgments

This work is dedicated to my parents, Lingxiang Yin and Yuying Hong. I am glad that they are here today to witness my accomplishments.

I would like to thank my advisor, Dr. Björn C. G. Söderberg, for his constant advising, patience, and encouragement during my graduate studies. I would also like to thank all faculty members in the C. Eugene Bennett Department of Chemistry for their assistance over the years, especially my doctoral defense committee members.

Finally, thanks to all my research group members, past and present, for all of their help and friendship.

Table of Contents

Title Page	i
Abstract	ii
Acknowledgment	iii
Table of Contents	iv
Appendix	vii
List of Figures	viii
List of Schemes	xvi
List of Tables	xix

Chapter 1

Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones via ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones

1.1 Rearrangement of 4-alkynylcyclobutenones	2
1.1.1 Thermally induced ring expansions of 4-alkynylcyclobutenones	3
1.1.2 Electrophilic reagents induced ring expansions of 4-alkynylcyclobutenones	15
1.2 Ring expansion of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones in the presence of an <i>N</i> -halosuccinimide	20
1.3 Results and discussion	23
1.4 Conclusions	31

Chapter 2

Total synthesis of methyl linderone

2.1 Introduction to linderone and methyl linderone	33
2.2 Previous synthesis of linderone and methyl linderone	38
2.3 Methyl linderone via ring expansion of 4-ene-yne-substituted cyclo-2-buten-1-one and <i>N</i> -iodosuccinimide	45
2.3.1 Retrosynthetic analysis	45
2.3.2 Results and discussion	46
2.3.3 Conclusions	47

Chapter 3

Synthesis of α -arylated ketone via Wacker-type oxidation of aryl-substituted alkenes

3.1 α -Arylated ketone background	49
3.2 Wacker-type oxidation	58
3.3 Wacker-type oxidation of aryl-substituted alkenes	73
3.4 Results and discussion	77
3.5 Conclusions	84

Chapter 4

Supporting information: experimental procedures

4.1 Supporting information Chapter 1: Ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones	86
4.2 Supporting information Chapter 2: Total synthesis of methyl linderone	103
4.3 Supporting information Chapter 3: Wacker-type oxidation of aryl-substituted alkenes	107

Appendix

^1H and ^{13}C NMR Spectra

- ^1H and ^{13}C NMR for Chapter 1: Ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones 149
- ^1H and ^{13}C NMR for Chapter 2: Total synthesis of methyl linderone 201
- ^1H and ^{13}C NMR for Chapter 3: Wacker-type oxidation of aryl-substituted alkenes 208

List of Figures

Figure 1: Structures of arnebifuranone, isoarnebifuranone, nanaomycin D, and deoxyfrenolicin	6
Figure 2: Structures of (±)-terreic acid and (±)-isoperezone	12
Figure 3: Eicosanoid core and naturally occurring eicosanoids	16
Figure 4: Postulated reaction path for palladium catalyzed ring expansion reaction	17
Figure 5: Hydrolysis products 85 and 86	27
Figure 6: Structures of linderone, methyl linderone, lucidone, methyl lucidone isolated from <i>Lindera</i> species	34
Figure 7: Structures of Stigmahamones I-II	34
Figure 8: Structure of kanakugiol	36
Figure 9: Structure of calythrone	38
Figure 10: Structures of coumestan, phenylbenzofurans, and isoflavones	49
Figure 11: Chemical structures of various palladium-complex catalysts	52
Figure 12: Typical apparatus for Wacker-type oxidation	60
Figure 13: Palladium complex accounts for the deactivation	62
Figure 14: Structures of substrates 238 , 240 , 241 , and 242	74
Figure 15: Wacker-type oxidation of 1-aryl-1-propene	75
Figure 16: Products 276 and 277	82
Figure 17: ¹ H NMR of 2,3-bis(1-methylethoxy)-5-trimethylsilyl-2,5-cyclohexadiene-1,4-dione (70)	149

Figure 18: ^{13}C NMR of 2,3-bis(1-methylethoxy)-5-trimethylsilyl-2,5-cyclohexadiene-1,4-dione (70)	150
Figure 19: ^1H NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (71)	151
Figure 20: ^{13}C NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (71)	152
Figure 21: ^1H NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (72)	153
Figure 22: ^{13}C NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (72)	154
Figure 23: ^1H NMR of 2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (74)	155
Figure 24: ^{13}C NMR of 2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (74)	156
Figure 25: ^1H NMR of 2-(iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (75)	157
Figure 26: ^{13}C NMR of 2-(iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (75)	158
Figure 27: ^1H NMR of 2-(bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (76)	159
Figure 28: ^{13}C NMR of 2-(bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (76)	160
Figure 29: ^1H NMR of 5-chloro-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (77)	161

Figure 30: ^{13}C NMR of 5-chloro-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (77)	162
Figure 31: ^1H NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-iodoethynyl)-2-cyclobutene-1-one (78)	163
Figure 32: ^{13}C NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-iodoethynyl)-2-cyclobutene-1-one (78)	164
Figure 33: ^1H NMR of 5-iodo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (79)	165
Figure 34: ^{13}C NMR of 5-iodo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (79)	166
Figure 35: ^1H NMR of 2-(diiodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (80)	167
Figure 36: ^{13}C NMR of 2-(diiodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (80)	168
Figure 37: ^1H NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-bromoethynyl)-2-cyclobutene-1-one (81)	169
Figure 38: ^{13}C NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-bromoethynyl)-2-cyclobutene-1-one (81)	170
Figure 39: ^1H NMR of 5-bromo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (82)	171
Figure 40: ^{13}C NMR of 5-bromo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (82)	172

Figure 41: ^1H NMR of 2-(1-bromo-1-iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (83)	173
Figure 42: ^{13}C NMR of 2-(1-bromo-1-iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (83)	174
Figure 43: ^1H NMR of 2-(dibromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (84)	175
Figure 44: ^{13}C NMR of 2-(dibromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (84)	176
Figure 45: ^1H NMR of 4-hydroxy-2-methyl-3-(1-methylethoxy)-4-(2-trimethylsilylethynyl)-2-cyclobutene-1-one (89)	177
Figure 46: ^{13}C NMR of 4-hydroxy-2-methyl-3-(1-methylethoxy)-4-(2-trimethylsilylethynyl)-2-cyclobutene-1-one (89)	178
Figure 47: ^1H NMR of 2-trimethylsilylmethylene-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (90) and 5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (91)	179
Figure 48: ^{13}C NMR of 2-trimethylsilylmethylene-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (90) and 5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (91)	180
Figure 49: ^1H NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (92)	181
Figure 50: ^{13}C NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (92)	182

Figure 51: ^1H NMR of 6-iodo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (93)	183
Figure 52: ^{13}C NMR of 6-iodo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (93)	184
Figure 53: ^1H NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (94)	185
Figure 54: ^{13}C NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (94)	186
Figure 55: ^1H NMR of 6-bromo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (95)	187
Figure 56: ^{13}C NMR of 6-bromo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (95)	188
Figure 57: ^1H NMR of 3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (97)	189
Figure 58: ^{13}C NMR of 3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (97)	190
Figure 59: ^1H NMR of 2-(1-iodomethylene)-4-methyl-5-(1-methylethoxy)-4-cyclopentene-1,3-dione (98)	191
Figure 60: ^{13}C NMR of 2-(1-iodomethylene)-4-methyl-5-(1-methylethoxy)-4-cyclopentene-1,3-dione (98)	192
Figure 61: ^1H NMR of 2-(1-bromomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (99)	193
Figure 62: ^{13}C NMR of 2-(1-bromomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (99)	194

Figure 63: ^1H NMR of 2-(diiodomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (101)	195
Figure 64: ^{13}C NMR of 2-(diiodomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (101)	196
Figure 65: ^1H NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-iodoethanoate (104)	197
Figure 66: ^{13}C NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-iodoethanoate (104)	198
Figure 67: ^1H NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-bromoethanoate (105)	199
Figure 68: ^{13}C NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-bromoethanoate (105)	200
Figure 69: ^1H NMR of 2,3-dimethoxy-4-hydroxy-4-[(E)-4-phenyl-3-buten-1-yn-1-yl]-2-cyclobuten-1-one (137)	201
Figure 70: ^{13}C NMR of 2,3-dimethoxy-4-hydroxy-4-[(E)-4-phenyl-3-buten-1-yn-1-yl]-2-cyclobuten-1-one (137)	202
Figure 71: ^1H NMR of 4,5-dimethoxy-[(E)-2-(1-iodo-3-phenyl-2-propen-1-ylidene)]-4-cyclopentene-1,3-dione (138)	203
Figure 72: ^{13}C NMR of 4,5-dimethoxy-[(E)-2-(1-iodo-3-phenyl-2-propen-1-ylidene)]-4-cyclopentene-1,3-dione (138)	204
Figure 73: ^1H NMR of 4,5-dimethoxy-2-[(E)-3-phenyl-2-propen-1-ylidene]-4-cyclopente-1,3-dione (139)	205

Figure 74: ^{13}C NMR of 4,5-dimethoxy-2-[(E)-3-phenyl-2-propen-1-ylidene]-4-cyclopente-1,3-dione (139)	206
Figure 75: ^1H NMR of 4,5-dimethoxy-2-[(E)-1-(N,N-dimethylamino)-3-phenyl-2-propen-1-ylidene]-4-cyclopente-1,3-dione (140)	207
Figure 76: ^1H NMR of 4-fluoro-2-nitro-1-(1-propen-1-yl)benzene (233)	208
Figure 77: ^{13}C NMR of 4-fluoro-2-nitro-1-(1-propen-1-yl)benzene (233)	209
Figure 78: ^1H NMR of 2,4-dinitro-1-(1-propen-1-yl)benzene (234)	210
Figure 79: ^1H NMR of 4-carbomethoxy-2-nitro-1-(1-propen-1-yl)benzene (235)	211
Figure 80: ^1H NMR of 4-methoxy-2-nitro-1-(1-propen-1-yl)benzene (236)	212
Figure 81: ^{13}C NMR of 4-methoxy-2-nitro-1-(1-propen-1-yl)benzene (236)	213
Figure 82: ^1H NMR of 5-methoxy-2-nitro-1-(1-propen-1-yl)-benzene (237)	214
Figure 83: ^1H NMR of 3-nitrobenzaldehyde (250) and 1-(3-nitrophenyl)-1,2-propanedione (251)	215
Figure 84: ^{13}C NMR of 3-nitrobenzaldehyde (250) and 1-(3-nitrophenyl)-1,2-propanedione (251)	216
Figure 85: ^1H NMR of 1-(4-fluoro-2-nitrophenyl)-2-propanone (257)	217
Figure 86: ^{13}C NMR of 1-(4-fluoro-2-nitrophenyl)-2-propanone (257)	218
Figure 87: ^1H NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (258)	219
Figure 88: ^{13}C NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (258)	220
Figure 89: ^1H NMR of 4-fluoro-2-nitrobenzaldehyde (259) and 1-(4-fluoro-2-nitrophenyl)-1,2-propanedione (260)	221

Figure 90: ^{13}C NMR of 4-fluoro-2-nitrobenzaldehyde (259) and 1-(4-fluoro-2-nitrophenyl)-1,2-propanedione (260)	222
Figure 91: ^1H NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (263)	223
Figure 92: ^{13}C NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (263)	224
Figure 93: ^1H NMR of 4-carbomethoxy-2-nitrobenzaldehyde (266) and 1-(4-carbomethoxy-2-nitrophenyl)-1,2-propanedione (267)	225
Figure 94: ^1H NMR of 1-(5-methoxy-2-nitrophenyl)-2-propanone (269)	226
Figure 95: ^{13}C NMR of 1-(5-methoxy-2-nitrophenyl)-2-propanone (269)	227
Figure 96: ^1H NMR of 5-methoxy-2-nitro-benzaldehyde (270) and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (271)	228
Figure 97: ^{13}C NMR of 5-methoxy-2-nitro-benzaldehyde (270) and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (271)	229
Figure 98: ^1H NMR of 4-(2-nitrophenyl)-3-methoxy-2-methyl-1-butene (277)	230
Figure 99: ^{13}C NMR of 4-(2-nitrophenyl)-3-methoxy-2-methyl-1-butene (277)	231
Figure 100: ^1H NMR of 1-(2-bromophenyl)-1,2-propanedione (282)	232
Figure 101: ^{13}C NMR of 1-(2-bromophenyl)-1,2-propanedione (282)	233

List of Schemes

Scheme 1: Rearrangement of 4-alkynylcyclobutenones	2
Scheme 2: Preparation of 4-alkynyl-4-hydroxycyclobutenones from squaric acid	2
Scheme 3: Thermally ring expansion to form substituted 1,4-benzoquinones	3
Scheme 4: Thermally ring expansion to form substituted 1,4-benzoquinones and 2-alkylidene-1,3-cyclopentenediones	4
Scheme 5: Plausible mechanisms for thermal ring expansion of 4-hydroxy-2-cyclobuteneones	5
Scheme 6: Thermally ring expansion to form 2-alkylidene-1,3-cyclopentenediones	5
Scheme 7. Synthesis of isoarnebifuranone	8
Scheme 8. Synthesis of nanaomycin D	10
Scheme 9: Synthesis of deoxyfrenolicin	10
Scheme 10: Thermal ring expansion of 4-alkynyl- <i>tert</i> -butoxycyclobutenones	11
Scheme 11: Synthesis of the natural product, (±)-terreic acid	13
Scheme 12: Synthesis of the natural product, (±)-isoperezone	14
Scheme 13: Palladium(II) catalyzed ring expansion of 4-alkynyl-4-hydroxy-cyclobutenones	15
Scheme 14: Palladium catalyzed ring expansion-coupling reaction	14
Scheme 15: Iodine mediated ring expansion of 4-alkynyl-4-hydroxycyclobutenones	18
Scheme 16: Conversion of iodomethylenecyclopentene-1,3-dione to aminomethylenecyclopentene-1,3-dione	20

Scheme 17: Synthesis of ring expansion substrates 73 , 78 , and 81	21
Scheme 18: Synthesis of ring expansion substrates 89 and 96	22
Scheme 19: Attempted synthesis of 100	22
Scheme 20: Thermal ring expansion of <i>tert</i> -butoxy analog	30
Scheme 21: Ring expansion of an electron deficient substrate	30
Scheme 22: Proposed mechanism for inhibition of human chymase by compound 112	35
Scheme 23: Lee's synthesis route of methyl linderone in 1967	39
Scheme 24: Lee's synthesis route of methyl linderone in 1986	41
Scheme 25: Aoyama's synthesis route of methyl linderone	42
Scheme 26: Langer's synthesis route of linderone	43
Scheme 27: Retrosynthetic analysis of methyl linderone	46
Scheme 28: Synthesis of methyl linderone	47
Scheme 29: Transition metal-catalyzed arylation reactions of ketone enolates	50
Scheme 30: Synthesis of 2,3-diarylbenzofurans by means of a palladium catalyst system	50
Scheme 31: Plausible reaction mechanism for the synthesis of 2,3-diarylbenzofurans	51
Scheme 32: Synthesis of the benzomorphan derivatives	53
Scheme 33: Asymmetric α -arylation of 2-methyl- α -tetralones	53
Scheme 34: Nickel-catalyzed enantioselective α -arylation of cycloalkanones	54
Scheme 35: α -Arylation of <i>N</i> -tosyl-2'-aminoacetophenone with <i>o</i> -dibromobenzene	54
Scheme 36: Pd-catalyzed synthesis of 1,2,2-triarylethanones from aryl benzyl ketones	55

Scheme 37: Pd-catalyzed synthesis of 1,2,2-triarylethanones from acetophenones	55
Scheme 38: Pd-catalyzed synthesis of the acetonylpyrine	56
Scheme 39: Pd-catalyzed synthesis of benzyl methyl ketone	56
Scheme 40: Individual reactions of the Wacker process	58
Scheme 41: Wacker process of terminal alkenes	59
Scheme 42: Overall catalytic cycle of the Wacker process	60
Scheme 43. Modified Wacker oxidation using H ₂ O ₂ as the terminal oxidant	62
Scheme 44: Mimoun et al.'s mechanism for peroxide-mediated Wacker oxidation	63
Scheme 45: Formation of biogenetic intermediate 190 and 191	64
Scheme 46: Synthesis of (+)-dihydropinidine and (-)-epidihydropinidine	65
Scheme 47: Synthesis of (-)-pinidinone	65
Scheme 48: Synthesis of tetracycle (+)- 203	66
Scheme 49: Synthesis towards key hexahydronaphthalene 206	67
Scheme 50: The synthesis of 1,2,5-trisubstituted pyrrole	68
Scheme 51: C ₃ Symmetric tripyrrolyl benzene synthesis	68
Scheme 52: Synthesis of 1,3-cyclooctanedione 219 from cyclooctanone 217	69
Scheme 53: TBHP-mediated Wacker-type oxidation	70
Scheme 54: TBHP-mediated Wacker-type oxidation to form β-hydroxy ketone	71
Scheme 55: TBHP-mediated Wacker-type oxidation to form methyl ketone	71
Scheme 56: Benzoquinone-reoxidant Wacker-type oxidation to acetophenone	72

List of Tables

Table 1: Ring-expansion of 68 , 73 , 78 , and 81 in the absence or presence of NXS	23
Table 2: Ring-expansion of 89 and 96 in the absence or presence of NIS or NBS	28
Table 3: Synthesis of 1-aryl-1-propenes	73
Table 4: Ratios of products for crude reaction mixtures from oxidation of 2-ethenyl-1-nitrobenzene	76
Table 5: Wacker-type oxidation of aryl-substituted alkenes	77

Chapter 1

Formation of 2-halomethylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones via ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones

1.1 Rearrangement of 4-alkynylcyclobutenones

1.1.1 Thermally induced ring expansions of 4-alkynylcyclobutenones

1.1.2 Electrophilic reagents induced ring expansions of 4-alkynylcyclobutenones

1.2 Ring expansion of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones in the presence of an

N-halosuccinimide

1.3 Results and discussion

1.4 Conclusions

1.1 Rearrangement of 4-alkynylcyclobutenones

The rearrangement of 4-alkynylcyclobutenones **1** is an extremely useful method for the synthesis of substituted 1,4-benzoquinones **3** and 2-alkylidene-1,3-cyclopentenediones **2** (Scheme 1).^{1,2} Substituted 1,4-benzoquinones constitute members of compounds which are reasonable precursors to electrophilic alkylating agents^{3,4} while 2-alkylidene-4-cyclopentene-1,3-diones are attractive building blocks for synthesis of cyclopentanoids.⁵⁻⁷

Scheme 1: Rearrangement of 4-alkynylcyclobutenones

Moore et al. provided a simple method for the transformation of 4-alkynylcyclobutenones into benzoquinones and/or 2-alkylidene-4-cyclopentenediones, starting with cyclobutenediones **5**, a class of compounds which are easily prepared in a variety of substitution patterns from commercially available squaric acid (**4**).⁸ Alkynyl anions add in high yield to cyclobutenediones **5**⁹ to give 4-alkynyl-4-hydroxycyclobutenones **6** (Scheme 2), and the reaction occurs with good regioselectivity with a number of unsymmetrically substituted substrates.^{10,11}

Scheme 2: Preparation of 4-alkynyl-4-hydroxycyclobutenones from squaric acid

1.1.1 Thermally induced ring expansions of 4-alkynylcyclobutenones

In 1989, Moore and co-workers developed a limited thermal rearrangement route involving a (2-alkynylethenyl)ketene¹² for the synthesis of benzoquinones and/or 2-alkylidene-4-cyclopentenediones. The synthetic scope and specific examples illustrating the synthetic utility of this method are given in Schemes 3. The syntheses in Scheme 3 began with the conversion of dimethyl squarate (**7**) to the 4-alkynyl-2,3-dimethoxy-4-hydroxyl cyclobutenones **8** in 50-90% yield using the corresponding lithium acetylides (THF, -78 °C) followed by quenching of the reaction mixture with ammonium chloride. Thermolyses of the realized cyclobutenones (*p*-xylene, 138 °C) provided the 1,4-benzoquinones **9** as the only isolable products.

Scheme 3: Thermally ring expansion to form substituted 1,4-benzoquinones

The thermal ring expansion of 4-alkynyl-2,3-dimethoxy-4-hydroxyl cyclobutenones **8** is an extremely useful method for the synthesis of benzoquinones, which have a variety of substitution patterns for further study.^{13,14} It is also important to note that this reaction is apparent an ideal and highly regioselective generality for the synthesis of trimethylsilyl-substituted 1,4-benzoquinones, which are particularly regarded as potential bioreductive alkylating agents or reasonable precursors to electrophilic quinone methides subsequent to an in vivo reduction.

Recognizing that only the corresponding benzoquinones were detected as products of

the thermolyses of all of the above examples, in which the alkyne moiety is substituted with an alkyl group or a proton, Moore et al. decided to implement the same condition in cases when the substituent R of **10** (Scheme 4) is an alkoxy, phenyl, or trimethylsilyl group. Interestingly, a mixture of the benzoquinones **11** and 2-alkylidene-1,3-cyclopentenediones **12** resulted.

Scheme 4: Thermally ring expansion to form substituted 1,4-benzoquinones and 2-alkylidene-1,3-cyclopentenediones

In order to explain the observed results, Moore et al. proposed the mechanisms of thermal ring expansion of 4-hydroxy-2-cyclobutenones **13** as shown in Scheme 5, beginning with a remarkably selective electrocyclic ring opening upon mild thermolysis to form (2-alkynylethenyl)ketenes **14**. Next, a ring closure of ketene **14** occurs to give the unique diradicals (or zwitterions) **15** and **17**. Finally, an intramolecular transfer of the proton substituent from oxygen to carbon of diradicals gives products **16** and **18**.¹²

With knowledge that this rearrangement selectivity to give either benzoquinones or cyclopentenediones is significantly influenced by the R substituent of the alkyne moiety, an effort was made by Moore et al. to drive the rearrangement completely to the cyclopentenediones **20** when the cyclobutenones **19** are employed (Scheme 6). This came to fruition.

Scheme 5: Plausible mechanisms for thermal ring expansion of 4-hydroxy-2-cyclobuteneones

Scheme 6: Thermally ring expansion to form 2-alkylidene-1,3-cyclopentenediones

Stabilization of the carbon centered radical in **17** by the adjacent R group has been used as an argument for the formation of 2-alkylidene-4-cyclopentene-1,3-diones.¹² The intermediate **17** becomes favored over **15** when the substituent R becomes more and more capable of stabilizing an adjacent radical site (Scheme 5). In contrast, the presumed aromatic stabilization associated with six-membered ring formation **15** is overcome by direct stabilization of one of the radical centers by the adjacent R substituent when five-membered ring formation **17** takes place.

In general, alkynes having an ester or a carbon-carbon double bond in conjugation give only 2-alkylidene-4-cyclopentene-1,3-diones while terminal alkynes, alkyl, and trimethylsilyl substituted alkynes produce predominantly or only 1,4-benzoquinones. Mixtures of both products are usually encountered from substrates having an aryl or an alkoxy group attached to the alkyne.

Later, in the same year, Moore et al.¹⁵ utilized their methodology described in the preceding paper, the rearrangement of 4-alkynylcyclobutenones to 1,4-benzoquinones, in the preparation of selected naturally occurring products arnebifuranone (**21**), isoarnebifuranone (**22**), nanaomycin D (**23**), and deoxyfrenolicin (**24**) to further illustrate its wide utility and synthetic scope.

Figure1: Structures of arnebifuranone, isoarnebifuranone, nanaomycin D, and deoxyfrenolicin

Arnebifuranone (**21**) is a naturally occurring 2,3-dimethoxy-1,4-benzoquinone^{16,17} which possesses *Z* stereochemistry at the stereogenic alkene site.¹⁸ Arnebifuranone, along with des-*O*-methyllasiiodiplodin, shikonofuran, arnebinol, and arnebinone have been isolated

from the roots of *Arnebia euchroma* (Royle) Johnst., which has been used as a pain killer in the Chinese culture.¹⁹ These compounds are of high interest because of their various pharmacological assay systems and they are able to function as inhibitors of prostaglandin biosynthesis.^{20,21} The attempted synthesis of arnebifuranone (**21**), was outlined in Scheme 7.

The synthesis of **22** initiated from 3-(2-bromoethyl)furan (**25**), which was converted to the corresponding Grignard reagent using magnesium in ether, and was further transformed into **26** in 75% yield upon the treatment with 2-methylpropenal. When SOCl₂ was added in hexane to **26**, the allylic chloride **27** was obtained in 65% isolated yield and excellent regioselectivity. **27** was the major isomer of a 7:1 mixture of regioisomers. Exposure of **27** to the Grignard reagent obtained from (trimethylsilyl)ethyne under copper-catalyzed conditions and subsequently deprotection provided the previously unknown alkyne **28** in 76% yield as a colorless liquid. Alkyne **28** was then underwent deprotonation using butyl lithium and was reacted with dimethyl squarate (**7**) in THF at -78 °C to give the adduct **29** as a colorless oil. Finally, the developed methodology was applied as a key step in the construction of arnebifuranone (**21**).^{1,10} However, thermolysis of **29** in refluxing *p*-xylene (138 °C) for 2 h only gave isoarnebifuranone (**22**) as an orange oil in 86% isolated yield. Isoarnebifuranone is actually stereoisomeric with the natural product arnebifuranone (**21**) and is another benzoquinone with the thermodynamically more stable *E* configuration, which is consistent with the fact that the naturally occurring quinone is quite unstable.

Scheme 7: Synthesis of isoarnebifuranone

Another reported example that further illustrates the synthetic utility of the quinone-cyclobutenone rearrangement is the regiospecific syntheses of the naphthoquinones **34** and **36**, which are key synthetic precursors to the biologically important quinones nanaomycin D (**23**) and deoxyfrenolicin (**24**).

Nanaomycin D (**23**) and deoxyfrenolicin (**24**) are members of an important family of naturally occurring isochroman-1,4-naphthoquinones (approximately 45) which are of great importance due to the significant biological activity that they have elicited in antibiotics, antimycotics and antineoplastic assays.²²⁻²⁴ The definite and concrete biological action mechanism of these compounds remains elusive. The proposed mode of their biological action involves bioreductive alkylation due to their structural features, which was regarded

necessary to allow them to be employed as precursors to electrophilic quinone methides subsequent to an in vivo reduction.^{3,4}

Starting from **30**, treatment with the lithium salt of 3-(tetrahydropyranyloxy) propyne (2:1 mixture of diastereomers) at -78 °C in THF gave **31** and its regioisomer in a respective ratio of 3:1 as a yellow oil in 58% yield (Scheme 8). Moreover, each regioisomer was obtained as a 2:1 mixture of diastereomers. Further experimentation with **30** and other lithium acetylides revealed that the reaction regioselectivities could be improved to approximately 95:5 at -100 °C in a mixed solvent of THF/diethyl ether (1:1) while yields might be lowered as a result.²⁵ Reaction of **31** with an excess of allyl iodide followed by the addition of Ag₂CO₃ in dioxane gave the allyl ether **32** in >95% yield as a yellow oil.¹⁹ Allyl ether **32** was then converted to naphthoquinone **33** as a yellow oil in 71% yield upon thermolysis in refluxing *p*-xylene (138 °C). The reaction proceeded by removal of THP protecting group using 4:2:1 CH₃COOH/THF/H₂O, resulted in the synthetic precursor naphthoquinone **34** in 73% yield. Finally, nanaomycin D (**23**) was prepared from **34** according to the published methods.²⁷

The last example to demonstrate the synthetic utility of the quinone-cyclobutenone rearrangement is the synthesis of **36**, the precursor of deoxyfrenolicin, which was accomplished by performing a procedure analogous to that described above (Scheme 9). That is, once the major regioisomer product of the alkynylation of **30** with the lithium salt of 3-(tetrahydropyranyloxy)hexyne in hand, which is actually a mixture of two diastereomeric benzocyclobutenones with only 2% of the corresponding regioisomers. The major product was conveniently purified by flash chromatography and the allylation of the major product was carried out to give **35**, which was further converted to synthetic precursor

naphthoquinone **36** by the method reported above. Finally, deoxyfrenolicin (**24**) was prepared from **36** according to the published methods.²⁷

Scheme 8: Synthesis of nanaomycin D

Scheme 9: Synthesis of deoxyfrenolicin

The methodology provided successful syntheses of naphthoquinones **34** and **36**, constituting a formal synthesis of the natural isochroman-1,4-naphthoquinones nanaomycin D (**23**), and deoxyfrenolicin (**24**).^{29,30} Quinones **34** and **36** were obtained in 28% and 22% yield from **30** respectively. Therefore, the quinone-cyclobutenone rearrangement methodology is not only efficient, brief, and simple, but also is a potentially general route to naphthoquinones of specific substitution patterns.

In 1990, Moore et al. further developed their methodology where 2,3-di-*tert*-butoxycyclobutenedione (**37**) (Scheme 10) was a useful synthon for hydroxyquinones. Besides the synthetic scope of this methodology, they also described its specific utilization in successful preparation of natural products, (±)-terreic acid (**43**), and (±)-isoperezzone (**44**) based on the thermal ring expansion of 4-alkynyl-*tert*-butoxycyclobutenones.³¹

Scheme 10: Thermal ring expansion of 4-alkynyl-*tert*-butoxycyclobutenones

Moore et al. reported the synthesis of *tert*-butoxyquinones **41** from the thermal ring expansion of 4-alkynyl-butoxycyclobutenones **40** in *p*-xylene at 138 °C and subsequently transformed to hydroxyquinones **42** upon treatment with trifluoroacetic acid at low temperature (Scheme 10).

In a typical procedure, squaric acid (**4**) was first converted to 2,3-di-*tert*-butoxycyclobutenedione (**37**) without any difficulty.¹² Next, cyclobutenedione **37** was treated with an alkynyl organolithium reagent to give the cyclobutenones **38** which were then readily hydrolyzed to the cyclobutenediones **39** upon treatment with trifluoroacetic anhydride (TFAA).³² After regiospecific addition of an alkynyllithium reagent to the more electrophilic (nonvinylogous ester) carbonyl group in **39**, cyclobutenones **40** were furnished. As mentioned above, thermolysis of **40** gave the *tert*-butoxyquinones **41**, which are members of a rare class of quinones. Finally, a variety of hydroxyquinones **42** were provided via deprotection of **41** upon treatment with trifluoroacetic acid at 0 °C.³³

Two real complex examples of the specific synthetic utility of this methodology are given below. The first example is the total synthesis of the natural antibiotic (±)-terreic acid (**43**), and the second one is total synthesis of naturally occurring (±)-isoperezzone (**44**).

Figure 2: Structures of (±)-terreic acid and (±)-isoperezzone

The efficient synthesis of (±)-terreic acid (**43**), a naturally occurring antibiotic^{34,35} was outlined in Scheme 11.

Preparation of ($\pm$)-terreic acid (**43**) via thermal ring expansion of 4-alkynyl-*tert*-butoxycyclobutenones began with the alkylation of 3-*tert*-butoxy-4-methylcyclobutenedione (**45**) with lithium acetylides followed by ammonium chloride, yielding the cyclobutenones **46** in good yield. These transformations were carried out at -78 °C in THF and the reactions were efficient. Cyclobutenones **46** were then converted to benzoquinones **47** in respective yields of 79% and 69% in refluxing acetonitrile or benzene. Reactions of these quinones **47** with *tert*-butyl hydroperoxide or Triton B³⁶ gave epoxide **48** in yields 59% and 68%. De-*tert*-butylation of **48** under previous used condition (0 °C, trifluoroacetic acid) gave ($\pm$)-terreic acid (**43**) in 84% yield. The syntheses had an overall yields of 22% and 26%, starting with the cyclobutenedione **45** which was readily available from squaric acid (**4**). Compared with the previously reported work, these syntheses are superior to the other two routes to ($\pm$)-terreic acid.^{37,38}

Scheme 11: Synthesis of the natural product, ($\pm$)-terreic acid

Final illustration of the specific synthetic utility of the thermal rearrangement of 4-alkynyl-*tert*-butoxycyclobutenones methodology was described as follows, that is, total

synthesis of the naturally occurring ($\pm$)-isoperezone (**44**) (Scheme 12). The synthesis began via an alkylation of squarate **37** with 2-lithio-6-methyl-hept-5-ene, resulting in cyclobutenone **49**,³⁹ which was then dissolved in pyridine in the presence of trifluoroacetic anhydride to give the cyclobutenedione **50** in 52% yield. Cyclobutenone **51** was then obtained by the regiospecific addition of 1-lithiopropyne to the more electrophilic carbonyl group of **50** and thermolysis of **51** in refluxing benzene furnished quinone **52** in 70% yield. The last step to complete this total synthesis of ($\pm$)-isoperezone was deprotection of **52** with 4 N HCl in refluxing dioxane, which directly provided **44** in 72% yield. The overall yield is 26% while the previous synthesis, reported in 1987, proceeded in 38% overall yield from parvifolene, a natural product isolated from *Pereziae spp.*⁴⁰

Scheme 12: Synthesis of the natural product, ($\pm$)-isoperezone

1.1.2 Electrophilic reagents induced ring expansions of 4-alkynylcyclobutenones

In addition to thermally induced ring expansion, some electrophilic reagents have also been shown to induce ring expansions of 4-alkynyl-4-hydroxy-2-cyclobuten-1-ones. For example, Liebeskind et al. proposed that the 4-alkynyl-4-hydroxy-cyclobutenone ring system **6**, was transformed to an alkylidenecyclopentenedione **53** possessing a vinyl-palladium bond via ring expansion reaction with an electrophilic source of Pd(II).^{41,42} **53** could be subsequently activated toward protonation by using the HX generated in the ring expansion step to provide access to synthetically useful alkylidenecyclopentenediones **54**. Meanwhile, the electrophilic Pd(II) reagent source rendering the reaction catalytic in Pd was regenerated (Scheme 13).

Scheme 13: Palladium(II) catalyzed ring expansion of 4-alkynyl-4-hydroxy-cyclobutenones

In 1987, Liebeskind et al.⁴³ successfully demonstrated a new, highly stereoselective and practical synthesis of highly functionalized 4-oxygenated 5-alkylidenecyclopentenones and 3-oxygenated 2-alkylideneindanones based on the palladium-catalyzed reactions. As shown in Scheme 14, Palladium(II) catalyzed reactions afford exclusively 2-alkylidene-4-cyclopentene-1,3-diones. Exposing 4-(1-hexynyl)-4-hydroxy-3-methoxy-2-methyl-cyclobut-2-enone (**55a**) to the condition (10 mol % Pd(OCOCF₃)₂ in THF at 60 °C for 1 h,) afforded (*E*)-alkylidene cyclopentenedione **56a**, which was isolated together with the *Z*-stereoisomer

in a 12:1 ratio in 45% yield. Two other 4-alkynyl-4-hydroxycyclobutenones **55b**, **55c** were exposed to the similar conditions discussed above (10 mol % $\text{Pd}(\text{OCOCF}_3)_2$ in THF at 40 °C for 5 h) and resulted in the formation of **56b** in 49% yield (11 : 1 ratio) and **56c** in 40% yield (15: 1 ratio).

Scheme 14: Palladium catalyzed ring expansion-coupling reaction

The reaction is of interest because of its potential application in the synthetic field of the naturally occurring eicosanoids as well as simpler analogues of the natural products.

Figure 3: Eicosanoid core and naturally occurring eicosanoids

Marine organisms produce an unprecedented molecular diversity by the incorporation of elements like halogen that are not readily available to terrestrial species.⁴⁴ Naturally

occurring marine eicosanoids, such as the clavulone **58**⁴⁵ (claviridenones^{45,46}), chloro,^{6,48} bromo, and iodovulones (**59**),⁴⁹ and the punaglandin **60**,⁵⁰ containing a 4-oxygenated 5-alkylidenecyclopentenone core **57** (Figure 3), are often produced and of great importance due to their remarkable bioactivity that they possess in both vitro and vivo cytotoxicity studies.⁵¹ There has been numerous reported syntheses for the 4-oxygenated 5-alkylidenecyclopentenone core **57**,⁵²⁻⁵⁴ however, the remarkable stereoselectivity challenges associated with the construction of these complex alkylidene systems are reflected in the many distinct approaches that have been reported for their racemic and enantioselective total syntheses.⁵⁵⁻⁵⁸

The observed exceptional stereoselectivity in the palladium-induced ring expansions of 4-alkynyl-4-hydroxycyclobutenones was rationalized by the author.⁴³ It was postulated that non-vinyl bond “a” in **61** (Figure 4) was migrating to the adjacent sp hybridized carbon, resulting in a cationic intermediate with the ability to best stabilize the positive charge. The reaction proceeded through the most stable cationic intermediate and the ability of the migrating group to stabilize positive charge might play an important role in controlling the selectivity of the reaction. Therefore, a plausible reaction pathway might be the trans addition of bond “a” across the alkyne bond to furnish the desired alkylidene cyclopentenedione in a stereospecific fashion.

Figure 4: Postulated reaction path for palladium catalyzed ring expansion reaction

In 1995, Eguchi et al.⁵⁹ demonstrated the ring expansion in Scheme 15 was probably

cationic in nature, and the result of coordination of an electron deficient species [Pd(II) and H^+] to the alkyne followed by acyl migration.^{60,61} Those transformations in Scheme 15 employed diethyl squarate (**62**) as a starting material, which was easily converted to the 4-alkynyl-4-hydroxycyclobutenone ring system **63** upon treatment with the corresponding lithium acetylides (THF, -78 °C) followed by quenching of the reaction mixture with ammonium chloride.^{9,32} The realized cyclobutenones **63** were converted to the corresponding cyclopentene-1,3-diones **64** via the ring expansion employing reagents which were believed to form hypoiodite. Specifically, conversion of cyclobutenones **63a**, **63b**, **63c** to the corresponding cationic intermediate using $I_2/PhI(OAc)_2$ at -15 °C, followed by deprotonation afforded cyclopentene-1,3-diones **64a**, **64b**, **64c** in 68%, 63%, 57% yields respectively.

The mechanism of the reaction was thought to begin with the transfer of positive iodine atom from a hypoiodite moiety to alkyne the terminus and concomitantly, 1,2-acyl migration happened to form cyclopentene-1,3-diones **64**. One of the driving force for this reaction was the ring strain relief which would facilitate the 1,2-acyl migration, and hence ring expansion.

Scheme 15: Iodine mediated ring expansion of 4-alkynyl-4-hydroxycyclobutenones

Electrophilic reagent induced ring expansions of 4-alkynylcyclobutenones provide an alternative practical route to cyclopentene-1,3-diones from squaric acid, switching the rearrangement of 4-alkynyl-4-hydroxycyclobutenones from radical to ionic via electron deficient species. Overall, both thermally induced ring expansions and electrophilic reagent induced ring expansions of 4-alkynylcyclobutenones have significant synthetic utility in preparing substituted 1,4-benzoquinones and 2-alkylidene-1,3-cyclopentenediones.

1.2 Ring expansion of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones in the presence of an *N*-halosuccinimide

Although the ring expansion of 4-alkynylcyclobutenones has been well investigated, very little research has been examined the thermal ring expansion of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones in the presence of an *N*-halosuccinimide or thermal ring expansion of haloalkynylcyclobutenones. Herein, we paid an effort to prepare chloro-, bromo- or iodo-substituted methylenecyclopentene-1,3-diones and benzoquinones from squaric acid, which also may be useful for further conversion or employed in palladium catalyzed cross coupling reactions and ultimately in the synthesis of functionalized carbazoles. A simple case is demonstrated below (Scheme 16).⁵⁹

Scheme 16: Conversion of iodomethylenecyclopentene-1,3-dione to aminomethylene cyclopentene-1,3-dione

We examined the thermal ring expansion of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones in the presence of an *N*-halosuccinimide. If the cationic pathway is sufficiently slow, at the elevated temperatures required for a thermal rearrangement, it may be possible to halogenate the diradical **15** (Scheme 5)⁶² in the presence of an *N*-halosuccinimide. The results of reactions of 4-alkynyl-4-hydroxycyclo-2-buten-1-ones with or without an *N*-halosuccinimide will be presented in this chapter.

Four 2,3-di(1-methylethoxy)-4-alkynyl-4-hydroxy-2-cyclobuten-1-ones having

different alkynyl groups including ethynyl (**73**), 2-trimethylsilylethynyl (**68**), 2-iodoethynyl (**78**), and 2-bromoethynyl (**81**) groups were initially examined. Compound **68** and **73** were prepared according to literature procedures⁶³ and compound **78** and **81** were prepared from **68** using NIS or NBS, respectively, in the presence of a catalytic amount of AgNO₃ (Scheme 17). Compound **68** was prepared from diisopropyl squarate (**67**).

Scheme 17: Synthesis of ring expansion substrates 73, 78, and 81

In addition to the reactions discussed above, cyclobutenones **89**, readily available from diisopropyl squarate (**67**) as well, was converted to 2-methyl-3-(1-methylethoxy)-4-ethynyl-2-cyclobuten-1-one (**96**) as outlined in Scheme 18. This was accomplished by its initial treatment with an alkynyl organolithium reagent to give cyclobutenone **87**, which are readily hydrolyzed to **88** upon treatment with hydrochloric acid in dichloromethane (CH₂Cl₂). Cyclobutenedione **88** was then converted to **89** by the regiospecific addition of an alkynyllithium reagent to the more electrophilic carbonyl group. Subsequently, cyclobutenone **89** was transformed to **96** using potassium fluoride in methanol and water. Cyclobutenone **96** was then subjected to reactions with or without NIS or NBS.

Scheme 18: Synthesis of ring expansion substrates **89 and **96****

It is noteworthy that in an attempt to prepare iodoalkynyl derivative **100** by reaction of **89** with NIS, only diiodomethylene cyclopentenedione **101** was obtained in a low isolated yield (Scheme 19). It is plausible that **100** was formed in the reaction but immediately underwent NIS induced ring expansion under the reaction conditions.

Scheme 19: Attempted synthesis of **100**

1.3 Result and discussion

As described above, we first attempted to examine four 2,3-di(1-methylethoxy)-4-alkynyl-4-hydroxy-2-cyclobuten-1-ones having different alkynyl groups including ethynyl (**73**), 2-trimethylsilylethynyl (**68**), 2-iodoethynyl (**78**), and 2-bromoethynyl (**81**).

Each compound was subjected to thermolysis either in the absence or in the presence of an *N*-halosuccinimide at selected temperatures. The results of these reactions are summarized in Table 1.

Table 1: Ring-expansion of **68, **73**, **78**, and **81** in the absence or presence of NXS**

Entry ^a	Solvent (temp./time)	Cyclobutenone	NXS	Cyclopentenedione ^b	Quinone ^b

1	Benzene (140 °C/3 h)	68	No NXS	69+70 (77%, 1:18)	
2	Toluene (150 °C/3 h)	68	No NXS	69+70 (70%, 1:14)

3	Benzene (rt/3 h)	68	NIS	71 (65%)	
4	MeCN (rt/15h)	68	NIS	71 (51%)	
5	Toluene (rt/20 h)	68	NIS	71 (65%)

6	Toluene (rt/120 h)	68	NBS	No reaction	
7	Benzene (80 °C/3 h)	68	NBS	72 (25%) ^c	
8	Benzene (80 °C/6 h)	68	NBS	72 (40%)	

Table 1 (continued)

Entry ^a	Solvent (temp./time)	Cyclobutenone	NXS	Cyclopentenedione ^b	Quinone ^b
9	Benzene (80 °C/3 h)	68	NCS		No reaction
10	Toluene (150 °C/3 h)	68	NCS		69+70 (55%, 1:20)
11	Benzene (140 °C/3 h)	73	No NXS		74 (58%)
12	Benzene (rt/3 h)	73	NIS	75 (34%)	
13	Benzene (80 °C/3 h)	73	NIS	75 (36%)	
14	Benzene (80 °C/3 h)	73	NBS	76 (35%) ^d	
15	Benzene (80 °C/17 h)	73	NBS	76 (47%) ^e	
16	Benzene (80 °C/24 h)	73	NBS	76 (55%)	
17	Toluene (150 °C/3 h)	73	NCS		77 (24%)+ 74 (55%)
18	Toluene (150 °C/3 h)	73	NCS ^f		77 (40%)+ 74 (35%)
19	Toluene (150 °C/3 h)	73	NCS ^g		77 (45%)+ 74 (20%)
20	Benzene (140 °C/3 h)	78	No NXS	75 (42%)	79 (25%)

Table 1 (continued)

Entry ^a	Solvent (temp./time)	Cyclobutenone	NXS	Cyclopentenedione ^b	Quinone ^b

21	Benzene (rt/3 h)	78	NIS	80 (41%) ^h	
22	Benzene (80 °C/3 h)	78	NIS	80 (43%)

23	Benzene (140 °C/3 h)	81	No NXS	76 (31%)	82 (34%)

24	Benzene (80 °C/3 h)	81	NIS	83 (68%)

25	Benzene (80 °C/3 h)	81	NBS	84 (46%)	

^a See Experimental section for detailed reaction conditions.

^b Yield of product after chromatographic separation on silica gel.

^c Hydrolysis product **85** (8%) and starting material (**68**, 24%) were also obtained.

^d Hydrolysis product **86** (12%) and starting material (32%) were also obtained.

^e Hydrolysis product **86** (11%) and starting material (**73**, 10%) were also obtained.

^f Three equivalents of NCS were used.

^g Six equivalents of NCS were used.

^h Starting material (**78**, 29%) was obtained.

All thermolysis reactions were performed in an ACE-Glass pressure tube capped with a Teflon screw cap. Thermolysis of **68** at 140 °C in benzene gave an inseparable mixture of 1,4-benzoquinone **70** and the corresponding cyclopentene-1,3-dione **69** in an 18:1 ratio in 77% yield as evidenced by ¹H NMR (entry 1). The reaction of **68** is somewhat more selective and

higher yielding compared to the previously reported rearrangement of the analogous 2,3-dimethoxy-substituted compound wherein an 8:1 ratio was observed in a 46% yield.¹² Toluene was also used as the solvent with a slightly lower ratio of products in a comparable yield (entry 2).

The benzoquinone and cyclopentenedione products reported herein are easily differentiated by ¹H NMR analysis. For example, the methine protons of the two methylethyl groups of 4-cyclopentene-1,3-dione **69** resonate at 5.59 and 5.54 ppm, respectively. In contrast, the corresponding 1,4-benzoquinone methine protons of compound **70** resonate upfield at 4.81 and 4.80 ppm. A similar difference in chemical shifts for methoxy groups of related 4-cyclopentene-1,3-diones and 1,4-benzoquinones has previously been reported by Moore et al.^{1,12} In addition, the color of the pure products are different, the 1,4-benzoquinones are red while the 4-cyclopentene-1,3-diones are orange.

Reaction of **68** with *N*-iodosuccinimide (NIS) proceeded much faster and gave even at ambient temperature in benzene only 2-iodomethylene-4-cyclopentene-1,3-dione **71** (entry 3). The same product was observed using acetonitrile or toluene as the solvent, the former gave a somewhat lower yield of product **71** (entries 4 and 5). Reaction of **68** with *N*-bromosuccinimide (NBS) did not afford any appreciable amount of product at ambient temperature even after prolonged reaction time (entry 6). However, 2-bromomethylene-4-cyclopentene-1,3-dione **72** was isolated as the only ring expansion product after heating for 6 h at 80 °C (entry 8). A small amount of the hydrolysis product cyclobutenedione **85** was also isolated (Figure 5). Although the solvents were carefully dried before use, adventitious water present may still be present in the solution.

Figure 5: Hydrolysis products 85 and 86¹

The reaction of **68** in the presence of *N*-chlorosuccinimide (NCS) was even slower and did not proceed at all at 80 °C. Increasing the temperature of the reaction to 150 °C did not furnish any chlorinated product but gave instead 1,4-benzoquinone **70** as the only product (entries 9 and 10). The results obtained from the reactions of **68** are interpreted as a change from a rapid cationic process seen for NIS to a slower cationic process using NBS to a radical process not involving NCS. It should be noted that no reaction was observed at temperatures below 100 °C for **68** in the absence of NBS or NIS.

Thermal reactions of **73** were examined next. Heating **73** in benzene at 140 °C gave 1,4-benzoquinone **74** as the sole product (entry 11). As was observed for **68**, reactions using either NIS or NBS produced 2-iodo- and 2-bromo-methylene-4-cyclopentene-1,3-diones, respectively (entries 12-16). Substrate **73** appeared to react somewhat slower with NBS compared to **68** since a longer reaction time was required for complete consumption of the starting material. A small amount of the hydrolysis product **86** (Figure 5) was isolated in some cases (entries 14 and 15). Thermolysis of **73** in the presence of NCS gave the chlorinated 1,4-benzoquinone **77** and 1,4-benzoquinone **74** in a 1:2 ratio (entry 17). The yield of **77** was improved by addition of an excess of NCS however, a significant amount of 1,4-benzoquinone **74** was isolated even using 6 equiv of NCS (entries 18 and 19). The latter three reactions indicate that it is possible to obtain chlorinated 1,4-benzoquinones via this

methodology although, in low to moderate yields using a large excess of NCS. Due to the absence of chlorinated product using **68** and the mediocre yield isolated from **73**, NCS was not further examined.

Thermal ring expansions of haloalkynylcyclobutenones have not previously been reported. Heating **78** at 140 °C in benzene in the absence of NXS gave a mixture of the five- and six-membered products **75** and **79** (entry 20). A similar reactivity was observed using **81** affording **76** and **82** (entry 23). Reaction of **78** with NIS proved to be somewhat slower compared to **68** and **73**. A significant amount of starting material was isolated after 3 h at ambient temperature but a complete conversion to **80** was observed at 80 °C (entries 21 and 22). Finally, **81** was transformed to the iodobromomethylene and dibromomethylene derivatives **83** and **84** in moderate yields using NIS and NBS, respectively (entries **24** and **25**).

In addition to the reactions discussed above, two 2-methyl-3-(1-methylethoxy)-4-alkynyl-2-cyclobuten-1-ones (**89** and **96**) were prepared and subjected to reactions with or without NIS or NBS. The results are summarized in Table 2.

Table 2: Ring-expansion of **89 and **96** in the absence or presence of NIS or NBS**

Entry ^a	Solvent (temp./time)	Cyclobutenone	NXS	Cyclopentenedione ^b	Quinone ^b

1	Benzene (140 °C/3 h)	89	No NXS	90 ^c (22%, 1.2:1)	91 ^c (32%)

2	Benzene (80 °C /3 h)	89	NIS	92 (61%, 1.5:1)	93 (5%)

Table 2 (continued)

Entry ^a	Solvent (temp./time)	Cyclobutenone	NXS	Cyclopentenedione ^b	Quinone ^b
3	Benzene (80 °C/3 h)		NBS
4	Benzene (140 °C/3 h)		No NXS
5	Toluene (150 °C/3 h)		NIS
6	Benzene (80 °C/3 h)		NBS

^a For detailed reaction conditions, see Chapter 4 experimental procedures.

^b Yield of product after chromatography.

^c The products could not be separated by chromatography.

Thermal ring expansion of **89** gave an inseparable 2:3 mixture of five- and six-membered cyclic products (entry 1). The cyclopentenedione **90** was obtained as a 1.2:1 mixture of double bond isomers.¹² A mixture of products was obtained from reactions of **89** with either NIS or NBS at elevated temperatures (entries 2 and 3). The result is surprising since we have not observed the formation of a quinone in any previous reaction using NIS or NBS. It is also interesting to note that Moore et al. did not report the formation of a five-membered cyclic product upon thermolysis of the *tert*-butoxy analog of **46** (Scheme 20).²

Scheme 20: Thermal ring expansion of *tert*-butoxy analog

Heating a solution of **96** either in the absence or presence of NIS or NBS gave the more predictable result. That is, quinone **97** was formed upon thermolysis of **96** in the absence of an NXS, while reactions with NIS or NBS gave cyclopentenediones **98** and **99**, respectively (entries 4-6).

As the final example, the electron deficient alkyne **102** was examined. This substrate has previously been shown by Moore et al. to exclusively afford 2-alkylidene-4-cyclopentene-1,3-dione **103** upon thermolysis in xylenes (Scheme 21).¹² This selectivity for five-membered cyclic products was also observed upon reaction with either NIS or NBS where **104** and **105** were formed, respectively.

Scheme 21: Ring expansion of an electron deficient substrate

1.4 Conclusions

An in-depth study of the reactions of 4-(ethynyl)-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones with *N*-bromo- or *N*-iodosuccinimide, affording in most cases 2-halomethylene-4-cyclopentene-1,3-dione, is described. In addition, the ring expansion of haloalkynylcyclobutenones with or without the presence *N*-halosuccinimide has been developed as well. The thermal ring expansion of 4-alkynyl-4-hydroxy cyclobutenones in the presence of *N*-halosuccinimide to form 2-halogenated methylene-4-cyclopentene-1,3-diones and/or 2-halo-1,4-benzoquinones selectively is a transformation of synthetic importance since no reliable general route to them has previously been reported.

The synthetic scope of this methodology has been investigated in this chapter and its specific utilization in the synthesis of a natural product, methyl linderone, will be described in next chapter.

Chapter 2

Total synthesis of methyl linderone

2.1 Introduction to linderone and methyl linderone

2.2 Previous synthesis of linderone and methyl linderone

2.3 Methyl linderone via ring expansion of 4-ene-yne-substituted cyclo-2-buten-1-one and *N*-iodosuccinimide

2.3.1 Retrosynthetic analysis

2.3.2 Results and discussion

2.3.3 Conclusions

2.1 Introduction to linderone and methyl linderone

A handful of 2-alkylidene-4-cyclopentene-1,3-diones have been isolated from natural sources (Figure 6). For example, linderone (**106**) and/or methyl linderone (**107**) were isolated from the root of *Lindera pipericarpa*⁶⁴ and *Lindera erythrocarpa*.⁶⁵ *Lindera erythrocarpa* Makino is one of the Lauraceae woody plants, which are distributed widely in Asia, including the People's Republic of China, Korea, Japan. *Lindera* species, the large family Lauraceae that consists of mostly trees or shrub widely spread throughout the world, from the warmer regions of the earth, in both lowland and montane forests of Malaysia and Singapore⁶⁶ and in the base of mountain forests of the Far East. They are found to be important medicinal plants including the Shiny Laurel, *Lindera lucida* (B1.) Boerl., *L. strychnifolia*, *L. aggregate*, and *L. chunii*. *L. erythrocarpa* (Lauraceae). Their fruits are used as a traditional medicine because of their analgesic, digestive, diuretic, antidote, and antibacterial properties.^{65,67}

Plant secondary metabolites have provided an important source of drugs since ancient times and now around half of the practical drugs used are derived from natural sources. Many plants of Lauraceae have been applied in folk medicine for their interesting bioactivities. Previous studies of this species have afforded in addition to the cyclopentenone derivatives linderone (**106**) and methyl linderone (**107**), lucidone (**108**) and methyl lucidone (**109**) as lignans⁶⁵ which were isolated from the bark of *L. erythrocarpa*.⁶⁸ Methyl linderone (**107**), lucidone (**108**) and methyl lucidone (**109**) were also isolated from the fruits of *Lindera lucida*,^{69,70} and methyl linderone has been reported from both *L. erythrocarpa*⁶⁷ and *L. pipericarpa*.⁷¹

Figure 6: Structures of linderone, methyl linderone, lucidone, methyl lucidone isolated from *Lindera* species

106 Linderone

107 Methyl linderone

108 Lucidone

109 Methyl lucidone

In 2000, Wu et al.⁷² reported two novel, closely related cyclopentenones, stigmahamone I (**110**) and stigmahamone II (**111**), had been isolated from the seeds of *Fissistigma oldhamii*. *Fissistigma oldhamii* (Hemsl.) Merr. (Annonaceae) is a perennial shrub, which is distributed mainly in southern China. As a folklore medicine, this plant is used for liver protection, anti-inflammatory and anti-arthritic effects, and anti-tumor action.⁷³ These compounds are of high interest because of their significant activity in preliminary bioassay, such as in antiplatelet aggregation and so on.

Figure 7: Structures of Stigmahamones I-II

110 Stigmahamone I

111 Stigmahamone II

Nowadays, more and more scientific study has been reported on the bioactivities of the cyclopentenedione derivatives, linderone (**106**), methyl lucidone (**107**), lucidone (**108**) and methyl lucidone (**109**). Methyl linderone is not only a human chymase inhibitor,⁷⁴ but also exhibits cytotoxicity against a number of cell lines⁷⁵ and the linderones are chitin synthase inhibitors.⁷⁶

Human chymase is a chymotrypsin-like serine protease that is stored in the secretory granules of mast cells.⁷⁷ Although the precise physiological and pathological functions of human chymase have not been elucidated, several studies have suggested its involvement in cardiovascular diseases⁷⁸ and chronic inflammation following fibrosis.⁷⁹ In 2001, Aoyama et al.⁷⁴ first reported the biological activity of methyl linderone as a human chymase inhibitor, which are thought to be potentially useful tools for elucidating the physiological function of chymase and therapeutic agents.

The postulated inhibition mechanism of the core structure **112** existing in methyl linderone against human chymase was as presented in Scheme 22. Mechanistically, the formation of serine hydroxy *O*-alkenyl enzyme occurred through a process involving nucleophilic substitution of the active site serine residue (serine 195) in the enzyme at the β -olefinic carbon atom activated by two carbonyl groups, meanwhile, the phenyl group in **112** docks into the S1 pocket.

Scheme 22: Proposed mechanism for inhibition of human chymase by compound 112

In 2002, Kho et al.⁷⁵ suggested the important cytotoxic compound in *lindera erythrocarpa* was methyl linderone (**107**) based on evidence from the evaluating of three lignans, linderone (**106**), methyl linderone (**107**), and kanakugiol (**113**) in vitro cytotoxicity using three cancer cell line assay. Amongst the compounds, methyl linderone exhibited significant cytotoxicity against mouse melanoma (B16-F10), human acetabulum fibrosarcoma (HT 1080), and choronic myelogenous leukemia (K562) cancer cell lines, while linderone **106**, and kanakugiol **113** showed weak inhibitory activity.

Figure 8: Structure of kanakugiol

All four linderones and lucidones are inhibitors of farnesyl transferase,⁸⁰ having anti-inflammatory activity,⁸¹ and they are PPAR-g agonists.⁸² In 2005, Kwon et al.⁸⁰ described the activity-guided isolation from the fruits of *L. erythrocarpa* and their inhibitory activities on FPTase. The results proved all four cyclopentenones mentioned above were farnesyl protein transferase (FPTase) inhibitors and anti-tumor compounds by examining their inhibitory activity against H-*ras*-transformed rat-2 cells and a variety of human cancer cell lines. Especially, compounds methyl linderone and methyl lucidone showed strong inhibition for the growth of human cancer cells and colon tumor xenografted. From their bioassay results, a conclusion was drawn that methyl lucidone can be a potential anti-cancer agent against H-*ras*-transformed tumor and the cyclopentenones will also be a good lead molecule for the development of a new class of anti-tumor drug and agents.

After Kwon and his coworkers demonstrated that four cyclopentenenediones from the methanol extract of the fruits of *L. erythrocarpa* exhibited farnesyl protein inhibitory activity, in 2007, Chang et al.⁸¹ reported their study of a bioactive investigation of the extract from *L. erythrocarpa* fruits presented a significant in vitro and in vivo anti-inflammatory activity in inhibiting significantly nitric oxide (NO) production in lipopolysaccharide induced NO in the murine macrophage cell line. Among the evaluated compounds, lucidone (**108**), cis/trans-methyl ludicone (**109**), methyl linderone (**107**) and linderone (**106**), lucidone possessed the strongest NO inhibitory activity. Furthermore, lucidone also provided anti-inflammatory activity in the croton oil-induced ear edema assay. It was indicated that lucidone has a good potential to be developed as an anti-inflammation agent.

In summary, *L. erythrocarpa* fruits possess potent anti-inflammatory activity, and according to the results obtained from large investigation, the cyclopentenenedione skeleton compounds, are good candidates for development as anti-tumor and anti-inflammatory agent. This class of compounds is of great importance in further nutraceutical applications, pharmaceutical industries and epidemiological studies.

2.2 Previous synthesis of linderone and methyl linderone

Linderone (**106**) and methyl linderone (**107**) are natural acylcyclopentenones structurally similar to calythrone (**114**)⁸³ (Figure 9) and methyl linderone belongs to a unique class of natural acylcyclopentenone pigments.^{70,84} To our best knowledge, linderone and methyl linderone have been synthesized four times since first isolated.^{74,85-87}

Figure 9: Structure of calythrone

114

In 1967, Lee group described the first total synthesis of linderone and methyl linderone.⁸⁵ It has been proposed these compounds might be derived from the appropriate precursor demethyldihydrolinderone (**118**), which could be readily generated through the alkaline rearrangement of the 1,4-quinone **117**.

Preparation of linderone and methyl linderone from a six-membered ring precursor (Scheme 23) via well known reactions involving ring-contraction⁸⁸ began with the oxidation of the chalcone **115** with nitric acid in acetic acid, indeed yielding the required quinochalcone precursor **116**. Quinochalcone precursor **116** was then readily debenzylated via catalytic hydrogenation in methanol, with simultaneous saturation of the cinnamoyl side-chain. Subsequently, the hydroquinone intermediate was transformed to the 2,6-dihydroxy-quinone **117** by aerial oxidation in alkaline solution. A yellow product identified as demethyldihydrolinderone (**118**) was obtained after the quinone stood at room temperature in aqueous sodium hydroxide for 2-2.5 h either in the presence or absence of air. With the demethyldihydrolinderone (**118**) in hand, there remained only the introduction of unsaturation into the acyl side chain to complete the total

synthesis of the natural product. Methylation of compound **118**, followed by dehydrogenation gave methyl linderone (**107**) and hence linderone (**106**). This was accomplished by first converting demethyldihydrolinderone (**118**) with diazomethane into methyldihydrolinderone (**119**) and then dehydrogenating the latter with 2,3-dichloro-5,6-dicyanoquinone, thus giving methyl linderone (**107**) in 60% yield. Finally, the partial hydrolysis of methyl linderone to linderone (**106**) completed the total synthesis of the desired naturally occurring compound.⁸⁴

Scheme 23: Lee's synthesis route of methyl linderone in 1967

In 1986, Lee et al.⁸⁶ reported they have utilized another direct approach⁸⁹ to synthesize linderone (**106**). That is converting methyl pedicinin (**125**) to **126** through the oxidative rearrangement by acetic anhydride-DMSO, which would be expected to undergo an acid-catalyzed conversion into linderone (**106**). The total synthesis employing this methodology is outlined in Scheme 24.

Methyl pedicinin (**125**), which could be isolated from the leaves of *Didymocarpus pedicellata*, was first synthesized in four steps from 2-hydroxy-3,4,6-trimethoxyacetophenone (**121**).⁹⁰ The synthesis started with a Friedel-Crafts acetylation of 1,2,3,5-tetramethoxy benzene (**120**) with acetyl chloride and aluminum chloride in dichloromethane at 30 °C, resulting in 2-hydroxy-3,4,6-trimethoxyacetophenone (**121**) in 66% yield. Treatment of **121** with benzaldehyde and potassium carbonate in alcohol for 3 days provided 2-hydroxy-3,4,6-trimethoxy-chalkone (**122**). The reaction of **122** with potassium carbonate and persulphate in pyridine at 10-15 °C, followed by refluxing in benzene in the presence of sodium sulfite and chloric acid furnished pedicin (**123**), which was then subjected to oxidation using bromine in chloroform to give pedicinin (**124**). Exposing pedicinin (**124**) to the methylation condition (diazomethane prepared from nitrosomethyl urea, ice-chest, 48 h) afforded methyl pedicinin (**125**). Subsequent rearrangement of **125** by using acetic anhydride-DMSO at 75 °C for 1 h gave compound **126** as a yellow solid identified as linderone acetate in 42% yield. Mild acid hydrolysis of compound **126** upon refluxing in 10% methanolic hydrochloric acid completed the synthesis of linderone (**106**).

Scheme 24: Lee's synthesis route of methyl linderone in 1986

In 2001, Aoyama et al.⁷⁴ performed a new and practical synthesis of methyl linderone (**107**). Methyl linderone was prepared using a palladium catalyzed rearrangement of **128** to **129** as the key step (Scheme 25). A DDQ mediated oxidation and a Michael-addition elimination sequence completed the synthesis.

This synthesis of **107** initiated from alkyne **127** prepared from hydrocinnamaldehyde by the Corey–Fuchs method.⁹¹ **127** was then converted to its lithium salt and added to dimethyl squarate (**7**) to give cyclobutenone **128**. The palladium-catalyzed ring expansion of **128** with 5% Pd(OCOCF₃)₂ in the presence of *N*-bromosuccinimide (NBS) provided compound **129** in 77% yield.⁴³ Dehydrogenation of **129** with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)^{85,86} followed by nucleophilic substitution with NaOMe gave methyl linderone (**107**). Using this protocol, total synthesis of methyl linderone (**107**) was achieved in only four steps and the overall yield from dimethyl squarate was 21%.

Scheme 25: Aoyama's synthesis route of methyl linderone

In 2004, Langer et al.⁸⁷ developed an approach to linderone using synthetic precursor **135** prepared from **130** in 5 steps in 41% overall yield (Scheme 26). When 2,4-bis-silyl enol ether **130** was treated with oxalyl chloride and trimethylsilyl triflate in dichloromethane at low temperature (-78-20 °C), compound **131** was obtained in 82% yield.

Scheme 26: Langer's synthesis route of linderone

Furanone **132** was formed in excellent yield by treating **131** with dimethyl sulfate and potassium carbonate in acetone. Exposure of **132** to trimethylsilyl triflate in diethyl ether in the

presence of triethyl amine generated the corresponding silyl enol ether **133** in 93% yield. Subsequent reaction of **133** with benzaldehyde provided compound **134**, which was then converted to **135** via dehydration. Finally, **135** was transformed to linderone (**106**), employing previously reported condition.⁹²

The total syntheses of linderone (**106**) and methyl linderone (**107**) were achieved by Lee group via ring-contraction reaction and oxidative rearrangement. However, their total syntheses consist of many steps. Aoyama group successfully performed a much shorter total synthesis compared to Lee's method. With the inspiration of their research and our study in the reactions of 4-(ethynyl)-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones with *N*-halosuccinimide, herein we report another synthetic route for the naturally occurring product methyl linderone (**107**).

2.3 Methyl linderone via ring expansion of 4-ene-yne-substituted cyclo-2-buten-1-one and N-iodosuccinimide

As discussed in Chapter 1, a new convenient methodology to form the 2-halomethylene-4-cyclopentene-1,3-diones was developed. It is of interest to extend this type of ring expansion reactions and utilize it as a convenient method for the preparation of other natural cyclopentenediones, such as methyl linderone (**107**). The specific utilization of this methodology in the synthesis of methyl linderone is described in this chapter.

2.3.1 Retrosynthetic analysis

The proposed retrosynthetic route and a retrosynthetic analysis of methyl linderone (**107**), which takes advantage of the above rearrangement, is outlined in Scheme 27. It diagrams linderone and methyl linderone can be prepared from a general precursor cyclopentenedione **138**. **138** can be seen as the product of the ring expansion of 4-ene-yne substituted cyclo-2-buten-1-one **137** in the presence of NIS. Compound **137** can be formed from deprotonation of 4-phenyl-3-butene-1-yne (**136**)⁹³ followed by addition of dimethyl squarate (**7**). This came to fruition.

Scheme 27: Retrosynthetic analysis of methyl linderone

2.3.2 Results and discussion

The synthesis was performed using the 4-ene-yne substituted cyclo-2-buten-1-one **137** and NIS (Scheme 28). Deprotonation of 4-phenyl-3-buten-1-yne (**136**) with butyllithium followed by addition of dimethyl squarate (**7**) gave 2,3-dimethoxy-4-(5-phenyl-3-penten-1-ynyl)-4-hydroxy-3-cyclobutene-1-one (**137**) in 47% yield. To our knowledge, only three cases of thermal ring expansions of 4-ene-yne-substrates have previously been reported affording five-membered cyclic products.¹² An intermolecular Diels-Alder dimerization of the diene system of the product has been reported and was initially a concern. Fortunately, thermolysis of **137** in the absence of NIS gave only **139** without a trace of Diels-Alder product. Reaction of **137** with *N*-iodosuccinimide gave the expected cyclopentenedione **138**. The iodide in **138** was smoothly replaced with a methoxy group to give the natural product methyl linderone (**107**) in a comparable 15% overall yield to the synthesis seen in Scheme 28. While the substitution of an iodide for a methoxy group was facile, we were unable to introduce a hydroxy group under basic or acidic conditions. Even the small amount of dimethylamine formed or present in dimethylformamide (DMF) reacted preferentially to give the dimethylamino derivative **140**.

Treatment of **138** using a number of other solvent systems including tetrahydrofuran-H₂O, 1,4-dioxane-H₂O, and *N*-methylpyrrolidinone-H₂O under both basic and acidic conditions only resulted in no reaction or decomposition of the starting material.

Scheme 28: Synthesis of methyl linderone

2.3.3 Conclusions

We have achieved an efficient short-step total synthesis of methyl linderone in only three steps with an overall yield of 15% from dimethyl squarate. The rearrangement of 4-ene-yne-substituted cyclo-2-buten-1-ones to halo-substituted cyclopentenediones in the presence of *N*-halosuccinimide can be utilized as key steps in the construction of cyclopentenedione derivatives having a variety of substitution patterns, which constitute members of a rare class of compounds with potential to be further developed to anti-cancer reagents and anti-tumor drugs in the future.

Chapter 3

Synthesis of α -arylated ketones via Wacker-type oxidation of aryl-substituted alkenes

3.1 α -Arylated ketone background

3.2 Wacker-type oxidation

3.3 Wacker-type oxidation of aryl-substituted alkenes

3.4 Results and discussion

3.5 Conclusions

3.1 α -arylated ketone background

The α -arylated ketone structural skeleton can be seen in a variety of biologically active natural products and they are very useful synthetic intermediates of active pharmaceutical ingredients such as coumestan (**141**),⁹⁴ phenylbenzofurans **142**,⁹⁵ and isoflavones **143** (Figure 10).⁹⁶ In organic synthesis, α -arylated carbonyl compounds could be used as synthetic building blocks in constructing more complicated substituted heterocycles such as indoles, furans, imidazoles, oxazoles, and pyrazoles.⁹⁷

Figure 10: Structures of coumestan, phenylbenzofurans, and isoflavones

The selective installation of an aryl group in the α -position of a carbonyl group has been an intense research subject over the past 40 years⁹⁸ and many synthetic methods have been reported⁹⁹⁻¹⁰² to construct these scaffolds, such as reaction of an enolate with a derivative of benzyne,⁹⁹ aryl reagents,¹⁰² Grignard reaction of benzamides with benzylmagnesium halides,¹⁰³ Friedel–Crafts reaction of phenols with various phenylacetic acids in $\text{BF}_3\text{-Et}_2\text{O}$,¹⁰⁴ and so on.

However, due to their own limitation, the transition metal-catalyzed direct coupling of aryl halides **144** or pseudohalides with enolates **145** obtained in situ from the corresponding ketones under basic conditions to synthesize of α -aryl substituted ketones **146** has emerged over the last three decades as a highly versatile method for the inter- and intramolecular construction of α -arylated ketones (Scheme 29).¹⁰⁵⁻¹⁰⁷ The most commonly employed

catalysts in the field of α -arylation of ketones with aryl halides are the palladium catalysts.

Scheme 29: Transition metal-catalyzed arylation reactions of ketone enolates

The earliest examples of transitional metal-catalyzed α -arylation of ketone enolates with aryl halides and pseudohalides was by Kuwajima,¹⁰⁸ Buchwald,¹⁰⁹ and Hartwig,¹¹⁰ in 1980s and 1990s. The reactions were usually carried out with a base and a combination of $\text{Pd}(\text{OAc})_2$ or a $\text{Pd}(0)$ compound such as $\text{Pd}_2(\text{dba})_3$ or $\text{Pd}(\text{dba})_2$ and a suitable phosphine ligand. One example of the synthetic utility of this methodology is Miura et al.¹¹¹ in 1999 prepared a variety of 2,3-diarylbenzofurans **149** via reaction of aryl benzyl ketones **147** with *o*-dibromobenzenes **148** (Scheme 30).

Scheme 30: Synthesis of 2,3-diarylbenzofurans by means of a palladium catalyst system

A plausible reaction sequence and key steps for these palladium-catalyzed reactions involves (i) the oxidative addition of the aryl halide to a $\text{Pd}(0)$ species; (ii) arylation at the benzylic position of the aryl benzyl ketone in the presence of a base, (iii) C-O bond formation, as shown in Scheme 31, (iv) a reductive elimination reaction of Pd species resulting observed diarylbenzofurans **149**.

Scheme 31: Plausible reaction mechanism for the synthesis of 2,3-diarylbenzofurans

In addition to tertiary phosphines, many excellent supporting ligands have been investigated in numerous palladium-catalyzed reactions and alternatives to the tertiary phosphines have been developed.¹¹²⁻¹¹⁴ A number of Pd-complexes catalysts show excellent activity in the α -arylation of various ketone substrates with aryl halides, such as allylchloro[1,3-bis(2,6-di-*i*-propylphenyl)-4,5-dihydroimidazol-2-ylidene]palladium [(SIPr)Pd(allyl)Cl] **150**^{112,113}, complex (IPr)Pd(OAc)₂ **151**¹¹⁴ [IPr=1,3-bis(2,6-di-*i*-propylphenyl)imidazol-2-ylidene], (IPr)Pd(acac)Cl **152**¹¹⁵, aminopalladacycle **153**,⁷⁸ complex **154**,⁷⁹ **155**,⁸⁰ **156**,⁸¹ and **157**.¹¹⁶⁻¹²⁰

Figure 11: Chemical structures of various palladium-complex catalysts

Khartulyari and Maier in 2007 highlighted the synthetic utility of the palladium-catalyzed intramolecular α -arylations of ketones in an attempt to prepare benzomorphan analogues **159**.¹²¹ Specifically, the cyclization of bromoketoesters **158a** and **159b** was carried out in refluxing toluene in the presence of K_3PO_4 as the base and a $\text{Pd}(\text{dba})_2$ and $\text{P}(t\text{-Bu})_3$ as the catalyst system to give tricyclic compounds **159a** and **159b** in 35% and 36% yield, respectively (Scheme 32).

Scheme 32: Synthesis of the benzomorphan derivatives

From 1998 until now, several asymmetric syntheses using the transition metal-catalyzed α -arylation of ketone with aryl halides have been developed.⁹⁵⁻⁹⁸ For instance, in 1998, Buchwald et al. performed the intermolecular asymmetric arylation of 2-methyl- α -tetralones **160** with 5 equiv of aryl halide at 70 °C, in the presence of 5 equiv of NaOt-Bu and Pd₂(dba)₃/(*S*)- or (*R*)-BINAP catalyst system.¹²² The enantio-enriched products **161** were obtained with good levels of the yield and enantiomeric excess (ee) (Scheme 33).

Scheme 33: Asymmetric α -arylation of 2-methyl- α -tetralones

In addition to palladium catalyzed α -arylation, nickel catalyzed asymmetric synthesis of ketones from organic halides and cycloalkanones was reported as well. In 2006, Kwong, Chan et al. used the atropoisomeric dipyridylphosphine (*R*)-P-Phos¹²³ to help pioneer **163** formation utilizing nickel-catalyzed α -arylation of cycloalkanone **162** with aryl halides **144**. Product **163** was enantio-selectively synthesized in excellent yields and ee (Scheme 34).

Scheme 34: Nickel-catalyzed enantioselective α -arylation of cycloalkanones

It is important to note that the synthesis of α -aryl substituted ketones by α -arylation of ketones with aryl halides or pseudohalides has been used extensively for the construction of key intermediates in pharmaceutically important compounds.

For example, *N*-tosyl-2'-aminoacetophenone (**164**) reacted with orthodibromobenzene (**165**) in toluene and water at 120 °C in the presence of Cs_2CO_3 and a combination of $\text{Pd}(\text{OAc})_2$ and XantPhos (Scheme 35) to provide intermediate **166** was a key step in a route to oxcarbazepine (**167**) (Trileptal).¹²⁴ Trileptal is the most widely prescribed drug for the treatment of epilepsy both in children and adults.

Scheme 35: α -Arylation of *N*-tosyl-2'-aminoacetophenone with *o*-dibromobenzene

Another example is the synthesis of a wide variety of tamoxifen-related 1,2,2-triarylethanones **170**, which function as active selective estrogen receptor modulator in the treatment of breast cancer.¹²⁵ SanMartin, Dominguez et al. in 2002 demonstrated that **170** were synthesized from the arylation reaction of aryl benzyl ketones **168** at the benzylic position with aryl bromides **169** (Scheme 36) with fair-to-good yield.

Scheme 36: Pd-catalyzed synthesis of 1,2,2-triarylethanones from aryl benzyl ketones

Later in 2004, 1,2,2-triarylethanones **170** were synthesized from acetophenones **171** and aryl bromides **172** in a better yield (Scheme 37).¹²⁶ Specifically, method A focused on the use of the system $\text{Pd}(\text{OAc})_2/\text{PPh}_3/\text{Cs}_2\text{CO}_3$, while method B entailed the use of the commercially available polymerbound catalyst FiberCat1026.

Scheme 37: Pd-catalyzed synthesis of 1,2,2-triarylethanones from acetophenones

Scheme 38: Pd-catalyzed synthesis of the acetonylpurine

In 2005, Gupta and Nair¹²⁷ applied the methodology in the synthesis of acetonylpurine (**174**), which is an important precursor in the synthesis of potent antiviral compound 2-acetonylinosine (**175**) via $\text{Pd}(\text{PPh}_3)_4$ -catalyzed reaction of 6-chloro-2-iodopurine ribonucleoside (**173**), Bu_3SnOMe , and isopropenyl acetate (Scheme 38). Benzyl methyl ketone **177** was prepared using a similar protocol. Reaction of aryl bromide **176**, isopropenyl acetate, and Bu_3SnOMe in the presence of $\text{Pd}_2(\text{dba})_3$ and 2-diphenylphosphino-2'-dimethylaminobinaphthyl catalyst system resulted in the desired product in a good yield.¹²⁸

Scheme 39: Pd-catalyzed synthesis of benzyl methyl ketone

Up to date, the transition metal-catalyzed synthesis of α -aryl substituted ketones by α -arylation of ketones with aryl halides or pseudohalides appears to be the most widely used as a very useful and powerful methodology for the formation of Csp^3 - Csp^2 bonds. However, these reactions suffer from some drawbacks. For instance, some necessary ligands, catalysts, additives for the reactions are very expensive; some substrates do not tolerate the strongly basic reaction conditions or required harsh conditions such as elevated temperatures; sometimes multiple arylation might happen and specific protecting group need to be introduced to block one α -position of the ketone.

Therefore, there is a need for the further development of direct installation of $C(sp^3)$ - $C(sp^2)$ bonds. Recently, our research in Wacker-type oxidation of aryl-substituted alkenes to furnish diverse α -aryl substituted ketones seems to be a promising alternative method for this goal.

3.2 Wacker-type oxidation

The Wacker oxidation, olefin oxidation with PdCl_2 catalysts, and related processes, has been a topic of research interest for well over a century.¹²⁹ Smidt and co-workers were the first researchers to describe the overall process, and the transition metal promoted oxidation of ethylene to give acetaldehyde was realized in the late 1950s by using palladium(II) chloride and copper(II) chloride as catalyst under an oxygen atmosphere (Scheme 40).¹³⁰ This transformation was achieved by the oxidation of an olefin with an aerobic PdCl_2 -catalyzed, Cu-mediated aqueous medium containing hydrochloric acid to a carbonyl.

The Wacker Process has remained one of the most important industrial process employing transition metal catalysts. Even though it was initially reported in 1962, it is still widely utilized for the industrial production of acetaldehyde from ethylene.

Scheme 40: Individual reactions of the Wacker process

The Wacker oxidation reaction of ethylene involves three sequential stoichiometric oxidation and reduction reactions to constitute a catalytic cycle. Eq. (1) in Scheme 40 represents an oxidation reaction involving a Pd(II) catalyst which has been known for more than a century;¹³¹ Eq. (2) indicates the formed Pd(0) in eq. (1) could be regenerated by cupric chloride in situ; the final step [Eq. (3)] is the re-oxidation of CuCl to CuCl_2 . The net reaction

of these three reactions leads to a simple air oxidation of ethene to ethanal.

In addition to ethylene, several synthesis have been developed by Smidt and coworkers specifically to produce other carbonyl compounds.¹³⁰ These studies involve a wide variety of starting materials and reaction conditions, including both an acidic aqueous solution of palladium(II) chloride and organic media.^{132,133} In general, terminal olefins are transformed to methyl ketones rather than aldehydes through Wacker process as shown in Scheme 41.

Scheme 41: Wacker process of terminal alkenes

Although a lot of efforts have focused on the mechanism study of Wacker reaction,¹³⁴⁻¹³⁶ a textbook example of a homogeneous transition-metal-catalyzed reaction, no “true” pathway of the Wacker oxidation process was described so far due to its complicated kinetics. However an abbreviated scheme of the complete catalytic cycle of the Wacker process was proposed in Scheme 42,¹³⁷ involving four key steps: the coordination of terminal alkene **179** to a palladium(II) species **178**; nucleophilic attack of water to the palladium-alkene complex **180** results intermediate **181**. Hydride shift in **181** followed by a reductive elimination furnish methyl ketone **182** and regenerate palladium(0) species, which could be reoxidized to palladium(II) catalyst. Extraordinary amount of research has provided strong evidence that subtle changes in experimental conditions, such as the structure of olefins, solvents and reoxidants, can result in substantially different reaction mechanisms,^{138,139} rates and yields.

Scheme 42: Overall catalytic cycle of the Wacker process

The oxidation on a laboratory scale can be carried out with palladium(II) catalyst and a copper salt under an oxygen atmosphere at room temperature using an apparatus as depicted in Figure 12.

Figure 12: Typical apparatus for Wacker-type oxidation

In order to apply this methodology to more complicated substrates, mixed-solvent systems have been developed and a lot of molecule-targeted syntheses were performed under

various conditions.^{130,134,139} The Wacker oxidations of higher terminal olefins are slow but this problem can be partly solved by the addition of suitable organic solvents which can mix olefins with water. However, it is worth noticing that oxidation and isomerization of the double bond are competitive reactions, and the extents of the reactions are influenced by solvents. For instance, dimethylformamide is good for the oxidation, whereas use of acetic acid facilitate the isomerization.¹³⁹ The double bond migration is facilitated by high temperatures.¹⁴⁰

A typical synthetic Wacker procedure, a mixed-solvent system of *N,N*-dimethylformamide and H₂O with catalytic PdCl₂ and stoichiometric CuCl under an aerobic atmosphere is referred to as Wacker-Tsuji oxidation. These classical conditions, using *N,N*-dimethylformamide (DMF)/H₂O mixture and stoichiometric CuCl,¹⁴¹ are developed by Tsuji based on modification and extension of the simple Wacker oxidation. Compared to other reported modified Wacker oxidation systems, these conditions are most common because of their convenience and simplicity.¹³⁴

Many modifications for Wacker oxidation have been made. Other than PdCl₂, palladium(II) trifluoroacetate, palladium(II) acetate, disodium tetrachloropalladate, palladium(II) sulfate, palladium(II) chloride-cobalt complexes, bis[acetonitrile]chloro-nitropalladium complex¹³⁹ and so on were used as catalysts as well. Sometimes, palladium catalyst deactivation occurs during the reaction by the precipitation of black palladium metal, even when no precipitation is observed. The reason accounts for this issue is the formation of palladium(II) complex. For example, palladium(II) chloride bis-dimethylamine complex deactivates the catalyst when dimethylformamide is used as a solvent,¹³³ and π -allyl

palladium complexes **183** may retard the reaction when some olefin substrates were used as a reactants.¹⁴²

Figure 13: Palladium complex accounts for the deactivation

Therefore, the activation of the palladium catalyst is a common issue in the Wacker oxidation as well as other Pd(II)-catalyzed oxidation reactions. Simple Wacker process adopts copper(II) chloride as a cocatalyst to efficiently regenerate the active Pd(II) catalyst. A number of other reoxidants have been introduced,¹³⁹ such as copper(I) chloride, copper(II) nitrate, copper(II) acetate, benzoquinone, hydrogen peroxide and *t*-butyl hydroperoxide. Compared to copper(II) catalyst, copper(I) chloride, when pretreated with oxygen, is used no chlorination of ketones takes place and the rate of the reaction is high. Benzoquinone was another good reoxidant and was first used by Moiseev et al.,¹⁴³ but a stoichiometric amount is necessary. The ability of hydrogen peroxide and *t*-butyl hydroperoxide to reoxidize palladium(0) as the terminal oxidant in the absence of Cu have been proved as well.¹⁴⁴ In 1980s, Mimoun and co-workers¹⁴⁵ reported that high conversion and selectivity were obtained from terminal straight-chain olefins by using 0.067 mol % Pd(OAc)₂ with 5 equiv of H₂O₂(aq) in acetic acid (Scheme 43). The isotopic labeling studies using ¹⁸OH₂ indicated that the O atom incorporated into the ketone originated from hydrogen peroxide.

Scheme 43: Modified Wacker oxidation using H₂O₂ as the terminal oxidant

On the basis of this finding, a mechanism was proposed wherein a palladacyclic intermediate **184** is formed from peroxide insertion into the olefin, followed by an α -hydride shift facilitated by O-O bond cleavage (Scheme 44).¹⁴⁶ An exciting aspect of the proposed catalytic cycle is that Pd does not necessarily proceed through Pd(0) because catalyst regeneration can occur via a proton transfer from a peroxide to **185**, regenerating active catalyst **186**. This potentially alleviates the probability of Pd(0) aggregation and catalyst deactivation, which is a common problem in these oxidations.

Scheme 44: Mimoun et al.'s mechanism for peroxide-mediated Wacker oxidation

Methyl ketones are important functional groups and the palladium(II)-catalyzed oxidation of terminal olefins seems to be one of the best methods to methyl ketones because terminal olefins are easily available, stable under acidic, basic or nucleophilic conditions.¹³⁹ Thus, synthetic organic chemists have paid great attention to the widespread use of the reaction as a synthetic method for ketones¹⁴⁷ and carried out a considerable amount of work on the applications to organic synthesis. The reaction is an extremely useful method for the synthesis of a variety of natural product precursors and biologically active compound

intermediates. Synthetic schemes from 45 to 52 are shown below as examples to demonstrate the usefulness of Wacker-type oxidation.

In 2006, Cook et al.¹⁴⁸ successfully converted stable (+)-macroline equivalent **188** to (-)-alstonerine (**189**) in 12.6% overall yield, starting from tryptophan methyl ester **187**, via an intramolecular Tsuji-Wacker oxidation (Scheme 45). The alkaloid **189** could be further converted into biogenetic intermediate (-)-talcarpine (**190**) and (-)-anhydromacrosalhinemethine (**191**). The transformations constituted an improved total synthesis of the (-)-anhydromacrosalhinemethine (**191**) as well as (-)-talcarpine (**190**).

Scheme 45: Formation of biogenetic intermediate 190 and 191

In 2011, Szolcsanyi et al.¹⁴⁹ have reported up to date shortest routes with highest overall yields for three naturally occurring 2,6-disubstituted piperidine alkaloids, (+)-dihydropinidine (**196**), (-)-epidihydropinidine (**195**) (as HCl salts), and (-)-pinidinone (**199**), which are naturally occurring defense chemicals, possessing antifeedant properties against

budworm and usually isolated from many spruce and pinus species and various insects. The total syntheses began with (*S*)-epichlorohydrin (**192**) as common starting material, and regioselective copper-free Wacker-Tsuji oxidation of alkenylazides **193** and **197** as key steps to exclusively generate methyl ketones **194** and **198** (Scheme 46). **194** subsequently converted to (-)-epidihydropinidine (**195**) and (+)-dihydropinidine (**196**) with 34% and 54% overall yield. The similar optimized Wacker-Tsuji transformation condition allowed the dialkenylazide **197** oxidized in good yield to the desired azidodiketone **198**. The latter was finally converted to (-)-pinidinone (**199**) in high yield (Scheme 47) upon treatment with hydrogenation on Pearlman's catalyst in 35% overall yield starting from **192**.

Scheme 46: Synthesis of (+)-dihydropinidine and (-)-epidihydropinidine

Scheme 47: Synthesis of (-)-pinidinone

In 2013, Yaragorla et al.¹⁵⁰ have demonstrated a concise approach toward the tetracycle **203**, whose structural skeleton is contained in a diverse range of shiartane-type nortriterpenoids. They display a wide ranging bioactivity profile that includes anti-oxidant, anti-HIV, anti-cancer activities and possess great potential in modern drug discovery as well as in traditional medicine. The synthesis began with the conversion of synthon (+)-**200** to the epoxy-ketone **201** in six steps, followed by exposure of **201** to Wacker–Tsuji oxidation condition provided cleanly to the desired diketone **202** in 72% yield and set the stage for further four transformations to the tetracycle **203** (Scheme 48). The total synthesis was completed in 11 steps with 8% overall yield.

Scheme 48: Synthesis of tetracycle (+)-203

In 2013, Pipelier et al.¹⁵¹ utilized the Wacker–Tsuji oxidation in the stereoselective synthesis of a hexahydronaphthalene derivative **206**, which was used as a building block or a suitable intermediate for the preparation of a variety of nardosinane-type compounds. Nardosinane sesquiterpene exhibits a wide biological activity spectrum and can be extracted from soft corals. They successfully developed two synthetic strategies to realize this goal (Scheme 49). The first route was achieved by the Wacker-type oxidation of silyl ether **204** to

give expected ketone **205** in 89% yield, which could be further transformed to bicyclic diene **206** in 60% yield (dr>95:5) under acidic condition, (para-toluenesulfonic acid in acetone/water). Then they described their improvement in the access to diene **206** by a shorter synthesis. Same to the condition used in the first strategy, allylic enol **207** was oxidized to methyl ketone **208** in 65% yield, the tertiary hydroxy group in which was subsequently undergo dehydration under acidic conditions to afford diene **206** still with high diastereoselectivity (dr>95:5) in 73% yield.

Scheme 49: Synthesis towards key hexahydronaphthalene 206

In 2004, Lubell et al.¹⁵² designed a convergent synthetic strategy from a series of structurally diverse aliphatic and aromatic esters **209** to give access to a range of pyrroles **212** by the three-step Grignard/Tsuji-Wacker/Paal-Knorr sequence in good yield and purity. Pyrroles are abundant as constituents of natural products⁵⁻⁸ and 1,2,5-trisubstitution pattern

pyrroles display interesting biological properties, including antipsychotic,¹⁰⁻¹² antiinflammatory,^{13,14} radioprotective,¹⁵ and spasmolytic^{16,17} activity. The oxidation of the olefin terminus was applied to ketone **210**, generating the 1,4-dione **211**, which could be used for next steps without further purification (Scheme 50).

Scheme 50: The synthesis of 1,2,5-trisubstituted pyrrole

Scheme 51: C₃ Symmetric tripyrrolyl benzene synthesis

10 Years later, Lubell group¹⁵³ further extended the utilization of their synthetic methodology mention above in the preparation of conjugated C₃ symmetric aryl tripyrroles, which may be promising materials for both electrochromic and fluorescence sensor applications. Three-step Grignard/Tsuji-Wacker/Paal-Knorr reaction sequence formed the

final tripyrroles **216** in 9% overall yields, starting from trimethyl 1,3,5-benzene tricarboxylate **213**. Modified Wacker-Tsuji olefin oxidation of homoallylic ketone **214** gave tris(1,4-dione) **215** in 49% yields, featuring ultrasound agitation of the substrate in the presence of PdCl₂ (0.6 equiv) and CuCl (3 equiv) in THF/H₂O (4:1) under oxygen atmosphere (Schemes 51).

Actually, a variety of terminal olefins can be oxidized to the corresponding methyl ketones via Wacker-type oxidation with functional groups remain intact, including aldehyde, carboxylic acid, ester, alcohol, ether, acetal, chloride, bromide, sulfonyl ester, sulfone and so on,¹³⁹ since the Wacker-type oxidation proceeds under mild conditions.

In 2011, Anseth et al.¹⁵⁴ first reported the successful application of the Wacker-type oxidation to a cyclic substrate to synthesis large-scale 1,3-cyclooctanedione (**219**) in five steps with 29% yield. The preparation started from converting the commercially available inexpensive starting material cyclooctanone (**217**) to enone **218** through 4 steps followed by a modified Wacker-Tsuji oxidation to gave the desired dione in a single step in 37% yield (Scheme 52).^{29,30} This synthesis was proved to be a good alternative method for the standard preparation of 1,3-cyclooctanedione.¹⁷ Moreover, this modified Wacker-Tsuji oxidation condition, which employed inexpensive oxidant, *tert*-butyl hydroperoxide was efficient for the oxidation of internal double bonds conjugated to carbonyl groups.⁴¹

Scheme 52: Synthesis of 1,3-cyclooctanedione 219 from cyclooctanone 217

A definite mechanism explaining various modified Wacker-type oxidations remains elusive because of the utilization of added ions and co-oxidants. Recent advancements have been made toward the elimination of Cu additives resulting in the development of peroxide-mediated and ligand modulation Wacker oxidations. Sigman et al. are among the researchers who paid great efforts in this area.

In 2009, Sigman et al.¹⁵⁵ discovered a highly efficient Wacker-type oxidation system capable of converting classically challenging substrates to corresponding methyl ketones, for example, protected allylic alcohols **220** to acyloins **221**, and homoallylic alcohol **222** to **223** in excellent yield (Scheme 53). In a typical reaction condition, low-loadings palladium(II)-Quinox (bidentate ligand, quinoline-2-oxazoline) complex, aqueous 70 wt% TBHP (*tert*-butyl hydroperoxide) and SbF_6^- was used as catalyst, terminal oxidant and counterion, respectively. Various terminal alkenes were selectively converted to methyl ketones under these mild conditions with high selectivity.

Scheme 53: TBHP-mediated Wacker-type oxidation

Two years later, in 2009, Sigman and co-workers¹⁵⁶ further proved the methodology they discovered before could be used as a general synthetic route to oxidize olefin substrates bearing homoallylic alcohols to their corresponding methyl ketone, regardless of substitution on the substrates. Various homoallylic alcohols, such as **224** was transformed to β -hydroxy ketone **225** using a TBHP-mediated Pd(Quinox)-catalyzed Wacker-type system in good yields with significantly reduced reaction times (Scheme 54).

Scheme 54: TBHP-mediated Wacker-type oxidation to form β -hydroxy ketone

A number of styrene derivatives are another challenging substrate class for the Wacker-Tsuji oxidation.¹⁵⁷ Therefore, more modified Wacker-type oxidations have been created for the purpose of enlarging its synthetic utility.

Exposing nitro-substituted styrene **226** to TBHP-mediated Wacker-type oxidation¹⁵⁵ mentioned above resulted in methyl ketone **227** in 60% yield because of the highly electron-deficient system and a trace amount of *m*-nitro benzaldehyde was obtained (Scheme 55).

Scheme 55: TBHP-mediated Wacker-type oxidation to form methyl ketone

In 2012, Gröger, et al.¹⁵⁸ converted styrene (**228**) to acetophenone (**229**) in excellent conversion and good selectivity of 99% and 90% respectively by using PdCl₂ as a palladium catalyst, benzoquinone as oxidation and a mixture of methanol and water as solvent. However,

side products *o*-methylphenylethan-1-ol (3%), phenylacetaldehyde dimethyl acetal (2%), methyl phenylacetate (3%), and traces of racemic phenylethan-1-ol (1%) were observed.

Scheme 56: Benzoquinone-reoxidant Wacker-type oxidation to acetophenone

Compared with the facile oxidation of terminal olefins, the reaction of Wacker-type oxidation of internal alkenes is extremely slow and the selectivity for unsymmetrical alkenes is often poor unless substrates with some functional groups at suitable positions would be oxidized regioselectively by participation of the functional groups under proper conditions. For example, the selectivity can be substantially enhanced if the alkene substrate has a pendant group able to coordinate with palladium (PdCl_2 , H_2O , O_2 (3 atm), DMA, 80 °C).¹⁵⁹ Thus, more improvement is necessary for the Wacker-type oxidation of internal olefins. The regioselective oxidation of functionalized internal olefins will be surveyed in this chapter because it would be a powerful and flexible method in the synthesis of α -aryl substituted ketones, which are quite important synthons in pharmaceutical and biomedical chemistry.

3.3 Wacker-type oxidation of aryl-substituted alkenes

Aryl-substituted alkenes are a substrate class that offers a unique set of challenges to Wacker-type oxidation. The 1-aryl-1-alkenes were prepared via a Kosugi-Migita-Stille cross coupling of an aryl-iodide or -bromide with tributyl 1-propen-1-yltin (Table 3)¹⁶⁰⁻¹⁶². Since the impetus of the present study was the palladium catalyzed oxidation, no attempts were made to optimize the yields of the cross coupling reactions.

Table 3: Synthesis of 1-aryl-1-propenes

Entry	X	R		Yield (%)
1	I	2-NO ₂	---	230 (49%)
2	Br	3-NO ₂	---	231 (49%)
3	Br	4- NO ₂	---	232 (57%)
4	I	2-NO ₂	4-F	233 (73%)
5	Br	2-NO ₂	4-NO ₂	234 (51%)
6	I	2-NO ₂	4-CO ₂ Me	235 (56%)
7	I	2-NO ₂	4-OMe	236 (32%)
8	I	2-NO ₂	5-OMe	237 (37%)
9	Br	2-NO ₂	6-Me	239 (69%)
10	I	2-Br	---	243 (40%)
11	I	---	---	244 (13%)

Three other alkene substrates, 1-nitro-2-(1-penten-1-yl)-benzene (**240**), 1-(3-methyl-1-butenyl)-2-nitrobenzene (**241**), and 1-nitro-2-(2-phenylethenyl)-benzene (**242**) (Figure 14) were synthesized using literature procedures.^{163,164} Compound (**238**), α -asarone is commercially available.

Figure 14: Structures of substrates 238, 240, 241, and 242

Our initial plan was to examine the possibility of generating α -arylated ketones directly under the typical Wacker-Tsuji oxidation condition (Figure 15), 10 mol % of PdCl₂ as a palladium catalyst, 1 equiv of copper chloride as oxidation reagent, and DMF/H₂O (10:1 (v/v) under an O₂ atmosphere at 50 °C for 20 h (Condition 1). The 1-aryl-1-propene **230** was expected to be oxidized to form α -arylated propanone **245**. But this condition yielded even no trace amounts of the expected **245**; the starting materials were recovered unchanged. A thorough study of the literature on Wacker process made us realize that the oxidation of internal olefins is extremely slow under the usual conditions and the reactions are not region-selective. Rates and yields of the oxidation are heavily dependent on the structure of olefins. Also, the proper selection of solvents and reoxidants is crucial. Due to the traditional Wacker-Tsuji oxidation condition seemed to be ineffective for this transformation, we chose first to change solvent by replacing DMF with MeOH (Condition 3). To our delight, a rapid reaction took place and α -arylated ketone **245** was isolated in a decent yield of 73% with dimethyl acetal **246** (4%) and an inseparable mixture of *o*-nitrobenzaldehyde **247** (7%) and dione **248** (7%) as side products. Subsequently, we decided to conduct an oxidant reagent screen by utilizing H₂O₂ with radical inhibitor (2,6-di-*t*-butyl-4-methylphenol) or *tert*-butyl hydroperoxide (TBHP) with radical inhibitor (Condition 4 and 5). α -Arylated ketone **245** was isolated in 72% and 92% yields respectively. We were pleased to find that the oxidation of **230** was very clean under Condition 5 (only a single product spot on the TLC) and the

desired methyl ketone was formed with a good 92% yield with no trace of any other by-products. It is important to note that the yield of methyl ketone **245** dropped off from 72% to 18% when changing the reaction condition from Condition 4 to 2. In fact, the addition of solvent (methanol and water) had a significant impact on the reaction yield (Conditions 2, 4).

Figure 15: Wacker-type oxidation of 1-aryl-1-propene

Reaction conditions:

1. Substrate (0.2 mmol), PdCl_2 (10 mol %), CuCl (1 equiv), 1.1 mL of $\text{DMF:H}_2\text{O}$ (10:1) at 50 °C, 20 h.
2. Substrate (0.2 mmol), PdCl_2 (10 mol %), H_2O_2 (10 equiv) with 50% con. in aqueous solution, 50 °C, 48h.
3. Substrate (0.2 mmol), PdCl_2 (10 mol %), CuCl (1 equiv), 1.1 mL of $\text{MeOH:H}_2\text{O}$ (10:1) at 50 °C, 20 h.
4. Substrate (0.2 mmol), PdCl_2 (10 mol %), H_2O_2 (10 equiv) with 50% con. in aqueous solution, 2,6-di-*t*-butyl-4-methylphenol (10 mol %), 1.1 mL of $\text{MeOH:H}_2\text{O}$ (10:1), 50 °C, 48 h.
5. Substrate (0.2 mmol), PdCl_2 (10 mol %), *tert*-butyl hydroperoxide (10 equiv) with 90% con. in aqueous solution, 2,6-di-*t*-butyl-4-methylphenol (10 mol %), 1.1 mL of $\text{MeOH:H}_2\text{O}$ (10:1), 50 °C, 48 h.

Inspired by these results, we next conducted a thorough solvent ratio screen using the oxidation of 2-nitro styrene (**285**) as the model (Table 4). $\text{MeOH/H}_2\text{O}$ (9:1) was found to be the best solvent system (entry 2), giving methyl ketone product in 29% yield. The reaction in H_2O was significantly slower (entry 6); only 61% of 2-nitro styrene was converted after 20 h

and 22% yield of methyl ketone **286** was isolated. Reactions using MeOH/H₂O (3/7) or DMF/H₂O (10/1) did not complete either and large amount of starting material was recovered with low yield of desired ketone obtained (entries 5 and 7). Neither anhydrous methanol (entry 1) nor DMF/MeOH (1/1) (entry 8) led to any methyl ketone product. It seemed that the ratio of water to methanol had a significant impact on the product ratio under these reaction conditions (entries 1-6). Ratios of different products from oxidation of 2-nitro styrene (**285**) were summarized in Table 4.

Table 4: Ratios of products for crude reaction mixtures from oxidation of 2-ethenyl-1-nitrobenzene

285		286	287	246	247	288
Entry	MeOH/H ₂ O					
1	1/- (100%)	0	17 (73%)	0	0	1
2	9/1	1 (29%)	2.3	0.15	0.25	0.16
3	7/3	1 (26%)	1.11	0.29	0.48	0.14
4	1/1	1 (16%)	0.60	0.73	1.28	0.26
5	3/7 ^a (21% sm)	1 (11%)	0.23	0	1.75	0.26
6	-/1 ^b (39% sm)	1 (22%)	0	0	1.31	0
7	10/1 ^c (8% sm)	1 (7%)	0	0	5.3 (37%)	0
8	1/1 ^d (36% sm)	0	1.4 (26%)	0	0.07	1 (19%)

The starting material was completely consumed in cases (1-4). PdCl₂ (10 mol %), CuCl (1 equiv), 50 °C, 20 h.

^a MeOH/H₂O (3/7) was used as the solvents and starting material (21%) was obtained after the reaction.

^b H₂O was used as the solvent and starting material (39%) was obtained after the reaction.

^c DMF/H₂O (10/1) was used as the solvents and starting material (8%) was obtained after the reaction.

^d DMF/MeOH (1/1) was used as the solvents and starting material (36%) was obtained after the reaction.

3.4 Results and Discussion

To ascertain the wide applicability of this Wacker-type oxidation methodology to target syntheses, the oxidations of a number of aryl-substituted internal alkene variants were investigated in an attempt to prepare structurally diverse α -arylated ketones (Table 5).

Table 5: Wacker-type oxidation of aryl-substituted alkenes

Entry	Alkene	Ketone	Acetal	Aldehyde	Dione
1-A	230	245 (73%)	246 (4%)	247/248 (1:1, 14%)	
1-B	230	245 (18%)		247 (8%)	
1-C	230	245 (72%)	246 (2%)	247 (21%)	
1-D	230	245 (92%)			
2-A	231		249 (30%)	250/251 (1:1, 14%)	
2-D	231	252 (34%)			
3-A	232		253 (32%)	254/255 (1:1, 20%)	
3-C	232		253 (34%)	254/255 (1:1, 14%)	
3-D	232	256 (55%)	253 (2%)		
4-A	233	257 (55%)	258 (3%)	259/260 (1:1, 10%)	
4-C	233	257 (35%)	258 (17%)		
4-D	233	257 (54%)			

Table 5 (continued)

Entry	Alkene	Ketone	Acetal	Aldehyde	Dione

5-A	234		261 (13%)	262 (39%)	263 (29%)
5-C	234	264 (31%)			
5-D	234	264 (5%)

6-A	235	265 (29%)		266/267 (1.1:1, 38%)	
6-C	235	265 (37%)		266/267 (1.1:1, 9%)	
6-D	235	265 (21%)		266/267 (1.3:1, 7%)

7-A	236	268 (53%)			
7-D	236	268 (55%)

8-A	237	269 (41%)		270/271 (1.3:1, 17%)	
8-C	237	269 (36%)		270/271 (1:1, 24%)

9-A	238	272 (7%)	273 (20%)		
9-D	238	nothing

10-A	239^c	99% recovered			
10-C	239	88% recovered			

Table 5 (continued)

Entry	Alkene	Ketone	Acetal	Aldehyde	Dione
11-A ^a	240	274 (58%)	246 (9%)		
11-B	240	274 (29%)	246 (13%)	247 (4%)	
11-C	240	274 (50%)	246 (12%)		
11-D	240	274 (64%)			
12-A ^b	241	275 (17%) (45% sm)	276 (11%)	277 (13%)	
12-B ^b	241	275 (18%) (19% sm)		247 (8%)	
12-C ^b	241	275 (25%) (36% sm)	246 (9%)	247 (6%)	
12-D	241	275 (17%)			
13-A ^b	242	278 (14%) (64% sm)			
13-C ^b	242	278 (51%) (20% sm)			
13-D	242	278 (74%)			
13-E	242	278 (27%)			
14-A	243	279 (45%)	280/281 (1:1, 9%)		282 (6%)
15-A	244	283 (34%)	284 (12%)		

^a 97% Conversion after 72h.^b Starting material was not completely consumed.^c No reaction was observed, 99% sm was recovered.

Table 5 (continued)

Conditions A: 10 mol% PdCl₂, 1 equiv. CuCl, O₂ (1 atm), MeOH:H₂O (10:1), 50 °C, 20 h.

Conditions B: 10 mol% PdCl₂, H₂O₂ (10 equiv) with 50% con. in aqueous solution, MeOH:H₂O (10:1), 50 °C, 48 h.

Conditions C: 10 mol% PdCl₂, H₂O₂ (10 equiv) with 50% con. in aqueous solution, 2,6-di-*tert*-butyl-*p*-cresol (10 mol %), MeOH:H₂O (10:1), 50 °C, 48 h.

Conditions D: 10 mol% PdCl₂, *t*-BuOOH (10 equiv) with 90% con. in aqueous solution, 2,6-di-*tert*-butyl-*p*-cresol (10 mol %), MeOH:H₂O (10:1), 50 °C, 48 h.

Conditions E: *t*-BuOOH (10 equiv) with 90% con. in aqueous solution was used in place of CuCl-O₂, no radical inhibitor.

These substrates, another challenging substrate class for the Tsuji-Wacker oxidation, were evaluated.¹⁵⁷ Substituted alkenes with various functional groups were efficiently oxidized, although highly electron-deficient systems were suspected to lead to a lower yield in previous report.¹⁵⁵ An underlying problem is the difficulty in predicting which pendant groups and which of the numerous possible variants of the Wacker-type oxidation will lead to the desired product for a given substrate. The selectivity in the Wacker-type oxidation of aryl-substituted alkenes appears to be under inherent substrate control and dependent on numerous factors, including substitution on the substrate and the precise conditions used, which can lead to a mixture of aldehyde, ketone, acetal or dione products. Highlighting this disparity, substrate such as **230** yielded methyl ketone product **245** in 73% yield in the presence of CuCl₂-O₂ (entry 1-A), while the acetal products **249** and **253** were observed as the predominant ones from similar substrates **231** and **232** under the same reaction condition (entries 2-A and 3-A). Additionally, the selectivity of these reactions can be affected by the choice of oxidant. For example, the use of the H₂O₂ favors the exclusive formation of 1-aryl-

propanone **264** in 31% yield (entry 5-C), and CuCl-O₂ favors the formation of products including acetal **261** (13%), aldehyde **262** (39%), and dione **263** (29%) (entry 5-A).

As described above, in our pursuit of the improvement of selectivity, we discovered a copper-free catalyst system for Wacker-type oxidations that utilizes palladium catalyst along with *tert*-butyl hydroperoxide (TBHP) as the oxidant and 2,6-di-*tert*-butyl-p-cresol as the radical inhibitor. This system successfully oxidized various functionalized aryl-substituted substrates with high selectivity (>98%) for the α -arylated ketone products (entries 1-4 D, 6-7 D, 11-13 D). The scope of condition A to E was then thoroughly evaluated, and it was found that, for some alkenes, it was sufficient to achieve good yields of desired ketones in 20 h (entries 4-A, 7-A, 8-A, 14-A) in the presence of 10 mol% PdCl₂, 1 equiv. CuCl, O₂ (1 atm) and MeOH:H₂O (10:1) at 50 °C. Steric hindered alkenes tended to react slower, where incomplete conversion was observed after reaction of 72 h (entries 10-13 A). No reaction was observed by ¹H NMR of the crude reaction mixture from 1-methyl-3-nitro-2-(1-propen-1-yl)-benzene (**239**) using either H₂O₂ or CuCl-O₂. The starting material was recovered in high yields (entries 10-A and 10-C). It seems steric hindrance considerably affects the rate of olefin oxidation. Mixtures of inseparable aldehyde and dione were also isolated from several reaction mixtures, as determined by ¹H NMR (entries 1-4 A, 6-A, 8-A, 3-C, 6-C, 8-C, and 6-D). With no exception, the TBHP-mediated catalyst system demonstrates high selectivity for the desired ketones, regardless of substitution on the substrates evaluated. This result is particularly interesting when compared with other modified conditions, where analogous substrates also formed other side products with different ratios and yields in a condition-dependent manner.

To investigate the effect of the position of the pendent nitro group relative to the internal olefin, nitro-1-propen-1-yl-benzene (**231** and **232**) were also evaluated. When these substrates were subjected to the Condition A [10 mol% PdCl₂, 1 equiv. CuCl, O₂ (1 atm), MeOH:H₂O (10:1), 50 °C, 20 h] (entries 2-A, 3-A), the desired ketone products were not observed; instead, the products isolated were identified acetals (**249**, **253**) and mixtures of aldehydes and diones. When these substrates were subjected to the condition D, methyl ketones **245** and **252** were obtained as the only products (entries 2-D, 3-D).

Halogens, such as F and Br (entries 4, 14) as well as methoxy and methyl ester groups (entries 6-9) are apparently well-tolerated in the meta-/ortho- position of the nitroarene substrates. Reaction of 1-(2,4-dinitrophenyl)-1-propene (**234**) gave methyl ester **261** in addition to the dione **263**, methyl 2,4-dinitrophenylbenzoate (**262**) (entry 5-A). It should be noted that the reported formation of this compound from reaction of 2,4-dinitro-1-fluorobenzene with dimethoxycarbene followed by hydrolysis is in error.¹⁶⁵ Oxidation of 1-(3-methyl-1-butenyl)-2-nitrobenzene (**241**) using condition A (entry 12) afforded isomers **276** and **277** in 11% and 13% respectively in addition to the α -arylated ketone **275**.

Figure 16: Products 276 and 277

Cornell and Sigman reported a high selectivity for acetophenone formation upon oxidation of *p*-/ *m*- or *o*-substituted styrenes with *t*-BuOOH in the presence of catalytic amounts of both a palladium(II) carbene complex and AgOTf.¹⁶⁶ However, oxidation of 1-phenyl-1-propene (**244**) was non-selective and gave an approximately 1:1 mixture of 1-

phenyl-2-propanone (**283**) and 1-phenyl-1-propanone (**284**). We observed a somewhat more selective oxidation of (**244**) affording roughly a 3:1 ratio of 1-phenyl-2-propanone and 1-phenyl-1-propanone (entry 15-A). The same 3:1 ratio of **283/284** was reported using PdCl₂, CuCl, in DMF-H₂O (1:1) at 1 atm of oxygen and ambient temperature.¹⁶⁰ A 95% yield was reported without any concurrent double bond cleavage to benzaldehyde. Cleavage of the alkene to form benzaldehyde was also observed in Sigman's paper (2.3:1 ratio of oxidation: cleavage).

Overall, the results point out that these modified Wacker processes of aryl-substituted alkenes are also quite sensitive to and dependent on reaction conditions, which are believed to associate with different mechanistic scenarios. However, *t*-butyl hydroperoxide and hydrogen peroxide have proved to be efficient reoxidants for the oxidation of internal double bonds.

3.5 Conclusions

In conclusion, we demonstrate that Wacker-type oxidation of aryl-substituted alkenes is feasible under modified Wacker-type oxidation conditions to yield α -arylated ketones that are challenging or impossible to obtain using conventional reaction conditions. The selectivity in the modified Tsuji-Wacker oxidation of aryl-substituted alkenes appears to be dependent on numerous factors, including the substrate and the precise conditions used. The reaction conditions discussed above appear to work with a fairly wide range of substrates.

Chapter 4

Supporting information: experimental procedures

- 4.1 Supporting information Chapter 1: Ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones
- 4.2 Supporting information Chapter 2: Total synthesis of methyl linderone
- 4.3 Supporting information Chapter 3: Wacker-type oxidation of aryl-substituted alkenes

General procedures. All NMR spectra were recorded in CDCl_3 . The chemical shifts are expressed in δ values relative to Me_4Si (0.0 ppm, ^1H and ^{13}C) or CDCl_3 (77.0 ppm, ^{13}C) internal standards. Tetrahydrofuran (THF) and diethyl ether were distilled from sodium benzophenone ketyl prior to use. Toluene, pyridine, hexanes, and ethyl acetate were distilled from calcium hydride. Chemicals prepared according to literature procedures have been footnoted the first time used; all other reagents were obtained from commercial sources and used as received. All reactions were performed under a nitrogen atmosphere in oven-dried glassware unless otherwise stated. The thermolysis reactions were performed in ACE-Glass pressure tubes capped with a Teflon screw cap. Solvents were removed from reaction mixtures and products on a rotary evaporator at water aspirator pressure. All alkanes, arenes, and etherial solvents are purified with activated alumina then Q5 (Q5 is an O_2 scavenging Cu-zeolite catalyst made by BASF). All other solvents with heteroatoms are dried through two columns of activated alumina.

Caution: 3,4-dialkoxy-3-cyclobutene-1,2-diones have been shown to cause contact dermatitis upon contact.²⁵

4.1 Supporting information Chapter 1: Ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones

4,5-Bis(1-methylethoxy)-2-(1-trimethylsilylmethylene)-4-cyclopentene-1,3-dione (69) and 2,3-bis(1-methylethoxy)-5-trimethylsilyl-2,5-cyclohexadiene-1,4-dione (70).

A solution of **68**⁹ (296 mg, 1.00 mmol) in benzene (3 mL) was heated at 140 °C for 3 h in an ACE-Glass pressure tube capped with a Teflon screw cap. The resulting dark mixture was cooled to ambient temperature and the solvent was removed at reduced pressure. The resulting crude product was purified by chromatography (hexanes/EtOAc, 98:2) to give an inseparable mixture of **69** and **70** (1:18 ratio, 229 mg, 0.773 mmol, 77%) as a red solid. Spectral data for **70**: mp 34-37 °C; ¹H NMR (270 MHz) δ 6.71 (s, 1H), 4.81 (sept, *J*=6.3 Hz, 1H), 4.80 (sept, *J*=6.1 Hz, 1H), 1.31 (d, *J*=6.1 Hz, 6H), 1.28 (d, *J*=6.1 Hz, 6H), 0.23 (s, 9H); ¹³C NMR (150 MHz) δ 188.1, 184.7, 150.4, 146.4, 145.3, 142.0, 75.9, 75.7, 22.6, -1.6; IR (ATR) 2978, 1657, 1583, 1251, 1197, 1108, 910, 885, 841 cm⁻¹; HRMS (ESI) calcd for C₁₅H₂₅O₄Si (M+H⁺) 297.1522; found 297.1517. Partial spectral data for **69**: ¹H NMR (270 MHz) δ 6.89 (s, 1H), 5.59 (sept, *J*=6.0 Hz, 1H), 5.54 (sept, *J*=6.1 Hz, 1H), 1.37 (d, *J*=6.1 Hz, 6H), 1.35 (d, *J*=6.1 Hz, 6H).

2-(1-Iodo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione

(71). To a solution of **68** (148mg, 0.500mmol) in benzene (1.5 mL) was added NIS (124 mg, 0.550 mmol). The tube was capped with a Teflon screw cap and stirred for 3 h at ambient temperature. Removal of solvent under reduced pressure followed by chromatography (hexanes/EtOAc, 9:1) gave **71** (137 mg, 0.324 mmol, 65%) as a yellow oil. ^1H NMR (270MHz) δ 5.62 (sept, $J=6.1$ Hz, 1H), 5.40 (sept, $J=6.1$ Hz, 1H), 1.35 (d, $J=6.1$ Hz, 6H), 1.34 (d, $J=6.1$ Hz, 6H), 0.33 (s, 9H); ^{13}C NMR (150 MHz) δ 183.9, 183.9, 151.2, 148.6, 140.8, 127.3, 75.2, 75.0, 23.1, 22.9, 1.7; IR (ATR) 2987, 1678, 1613, 1304, 1246, 1065, 841 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{24}\text{IO}_4\text{Si}$ ($\text{M}+\text{H}^+$) 423.0489; found 423.0486.

2-(1-Bromo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione

(72). Treatment of **68** (75 mg, 0.25 mmol) with NBS (59 mg, 0.33 mmol) in benzene (0.75 mL) as described for **71** (80 $^{\circ}\text{C}$, 6 h), gave after chromatography (hexanes/EtOAc, 99:1) **72** (38 mg, 0.10 mmol, 40%) as a brown oil. ^1H NMR (270 MHz) δ 5.57 (sept, $J=6.1$ Hz, 1H), 5.41 (sept, $J=6.1$ Hz, 1H), 1.36 (d, $J=6.1$ Hz, 6H), 1.36 (d, $J=6.1$ Hz, 6H), 0.34 (s, 9H); ^{13}C NMR (150 MHz) δ 184.3, 183.4, 151.4, 149.5, 148.0, 136.9, 75.2, 74.9, 23.1, 22.9, 0.3; IR

(ATR) 2993, 1682, 1622, 1385, 1306, 1098, 1068, 850 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{24}\text{BrO}_4\text{Si}$ ($\text{M}+\text{H}^+$) 375.0627; found 375.0619.

2,3-Bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (74). Treatment of **73**⁹ (56 mg, 0.25 mmol) in benzene (0.75 mL) as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2) **74** (33 mg, 0.14 mmol, 58%) as an orange-red oil. ^1H NMR (270 MHz) δ 6.60 (s, 2H), 4.82 (sept, $J=6.1$ Hz, 2H), 1.30 (d, $J=6.1$ Hz, 12H); ^{13}C NMR (67.5 MHz) δ 184.8, 145.8, 134.7, 76.1, 22.6; IR (ATR) 2983, 1662, 1588, 1375, 1291, 1187, 1098, 1063 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{17}\text{O}_4$ ($\text{M}+\text{H}^+$) 225.1127; found 225.1122.

2-(Iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (75). Treatment of **73** (112 mg, 0.500 mmol) with NIS (124 mg, 0.550 mmol) in benzene (1.5 mL) as described for **71** (ambient temperature, 3 h), gave after chromatography (hexanes/EtOAc, 19:1 then 8:2) **75** (59.7 mg, 0.170 mmol, 34%) as a pale orange oil. ^1H NMR (270 MHz) δ 7.87 (s, 1H), 5.62 (sept, $J=6.1$ Hz, 1H), 5.51 (sept, $J=6.1$ Hz, 1H), 1.38 (d, $J=6.1$ Hz, 6H), 1.36 (d, $J=6.1$ Hz, 6H); ^{13}C NMR (150 MHz) δ 183.5, 182.6, 153.4, 149.0, 136.3, 88.1, 75.4, 75.2, 23.1, 23.0;

IR (ATR) 2983, 1682, 1627, 1583, 1390, 1306, 1098, 1043, 910, 781 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{IO}_4$ ($\text{M}+\text{H}^+$) 351.0093; found 351.0090.

2-(Bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (76). Treatment of **73** (57 mg, 0.25 mmol) with NBS (55.0 mg, 0.309 mmol) in benzene (1.5 mL) as described for **70** (80 $^{\circ}\text{C}$, 17 h) gave after chromatography (hexanes/EtOAc, 95:5), **76** (36 mg, 0.12 mmol, 47%) as a yellow oil. ^1H NMR (270 MHz) δ 7.12 (s, 1H), 5.42 (sept, $J=6.2$ Hz, 1H), 5.36 (sept, $J=6.1$ Hz, 1H), 1.23 (d, $J=6.1$ Hz, 6H), 1.21 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (150 MHz) δ 182.7, 182.6, 153.2, 149.7, 131.8, 116.4, 75.5, 75.2, 23.1, 23.0; IR (ATR) 2993, 1682, 1637, 1390, 1316, 1103, 910, 796 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{BrO}_4$ ($\text{M}+\text{H}^+$) 303.0232; found 303.0227.

5-Chloro-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (77) and 2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (74). Treatment of **73** (110 mg, 0.491 mmol) with NCS (72 mg, 0.539 mmol) in toluene (3 mL) as described for **71** (150 $^{\circ}\text{C}$, 3 h) gave after chromatography (hexanes then hexanes/EtOAc, 98:2), in order of elution, **77** (30 mg, 0.116

mmol, 24%) and **74** (60 mg, 0.27 mmol, 55%) both as orange oils. Analytical data for **77**: ^1H NMR (270 MHz) δ 6.82 (s, 1H), 4.87 (sept, $J=6.1$ Hz, 1H), 4.76 (sept, $J=6.1$ Hz, 1H), 1.32 (d, $J=6.1$ Hz, 6H), 1.31 (d, $J=6.3$ Hz, 6H); ^{13}C NMR (150 MHz) δ 182.3, 177.1, 146.1, 145.6, 142.0, 131.5, 76.7, 22.6; IR (ATR) 2987, 1659, 1590, 1389, 1254, 1096, 872, 779 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{ClO}_4$ ($\text{M}+\text{H}^+$) 259.0737; found 259.0732.

4-Hydroxy-2,3-bis(1-methylethoxy)-4-(2-iodoethynyl)-2-cyclobutene-1-one (78). A solution of **68** (914 mg, 3.08 mmol) under N_2 in DMF (30 mL) was cooled to 0 $^\circ\text{C}$ and AgNO_3 (38mg, 0.22 mmol) was added in one portion. After stirring for 5 min, NIS (764mg, 3.40 mmol) was added and the reaction was slowly warmed to room temperature overnight. The mixture was cooled to 0 $^\circ\text{C}$ and ice-water (20mL) was added. The mixture was extracted with Et_2O (6 x 50 mL). The organic layers were washed with water followed by brine and dried (MgSO_4). The solvent was removed under reduced pressure. Chromatography (hexanes/ EtOAc , 7:3) gave **78** (777 mg, 2.22 mmol, 72%) as a pale yellow solid. Mp 108-110 $^\circ\text{C}$; ^1H NMR (270 MHz) δ 4.96 (sept, $J=6.1$ Hz, 1H), 4.84 (sept, $J=6.1$ Hz, 1H), 3.75 (s, 1H), 1.42 (d, $J=6.3$ Hz, 6H), 1.28 (d, $J=6.1$ Hz, 3H), 1.27 (d, $J=6.1$ Hz, 3H); ^{13}C NMR (150MHz) δ 180.0, 164.1, 134.3, 88.9, 79.7, 78.2, 74.3, 22.7, 22.6, 22.5, 7.9; IR (ATR) 2988, 1771, 1617, 1390, 1325, 1098, 999, 954, 910, 737 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{IO}_4$ ($\text{M}+\text{H}^+$) 351.0093; found 351.0088.

2-(Iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (75) and 5-iodo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (79). Treatment of **78** (84 mg, 0.24 mmol) in benzene (0.75 mL) as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2), in order of elution **79** (21 mg, 0.060 mmol, 25%) and **75** (35.0 mg, 0.100 mmol, 42%) both as orange oils. Analytical data for **79**: ^1H NMR (270 MHz) δ 7.50 (s, 1H), 4.83 (sept, $J=6.1$ Hz, 1H), 4.69 (sept, $J=6.1$ Hz, 1H), 1.27 (d, $J=6.1$ Hz, 6H), 1.26 (d, $J=6.1$ Hz, 6H); ^{13}C NMR (150 MHz) δ 182.4, 178.2, 146.4, 144.3, 144.0, 117.0, 76.9, 76.5, 22.7; IR (ATR) 2983, 1667, 1578, 1390, 1251, 1182, 1103, 910 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{IO}_4$ ($\text{M}+\text{H}^+$) 351.0093; found 351.0088.

2-(Diiodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (80). Treatment of **78** (76 mg, 0.20 mmol) with NIS (66.6 mg, 0.296 mmol) in benzene (0.75 mL) as described for **70** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 95:5) **80** (45 mg, 0.094 mmol, 43%) as a pale yellow solid and **78** (23.4 mg, 0.0668, 30%). Data for **80**: mp 89-94 °C; ^1H NMR (270 MHz) δ 5.51 (sept, $J=6.1$ Hz, 2H), 1.37 (d, $J=6.2$ Hz, 12H); ^{13}C NMR (150 MHz) δ 182.7, 149.8, 133.2, 75.5, 23.0, 17.8; IR (ATR) 2983, 1672, 1617, 1385, 1311, 1103, 1078,

910, 796 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{15}\text{I}_2\text{O}_4 (\text{M}+\text{H}^+)$ 476.9060; found 476.9055.

4-Hydroxy-2,3-bis(1-methylethoxy)-4-(2-bromoethynyl)-2-cyclobutene-1-one (81).

A solution of **68** (1.00 g, 3.38 mmol) under a nitrogen atmosphere in DMF (30 mL) was cooled to 0 °C and AgNO_3 (40 mg, 0.24 mmol) was added. After stirring for 5 min, NBS (696 mg, 3.91 mmol) was added and the reaction was slowly warmed to room temperature overnight. The mixture was cooled to 0 °C and quenched with ice-water (20 mL). The mixture was extracted with ether (3 x 75 mL). The combined organic layers were dried (MgSO_4), filtered, and the solvent was removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to afford **81** (485 mg, 1.60 mmol, 47%) as a pale yellow solid. Mp 109-110 °C; ^1H NMR (600 MHz) δ 4.95 (sept, $J=6.0$ Hz, 1H), 4.88 (sept, $J=6.0$ Hz, 1H), 2.76 (s, 1H), 1.44 (d, $J=6.0$ Hz, 3H), 1.44 (d, $J=6.0$ Hz, 3H), 1.31 (d, $J=6.6$ Hz, 3H), 1.30 (d, $J=6.6$ Hz, 3H); ^{13}C NMR (150 MHz) δ 180.1, 164.2, 134.0, 79.1, 78.1, 75.0, 74.2, 49.8, 22.6, 22.6, 22.5, 22.4; IR (ATR) 3303, 1775, 1621, 1378, 1331, 1096, 1007, 953, 907, 783 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{15}\text{BrNaO}_4 (\text{M}+\text{Na}^+)$ 325.0051; found: 325.0045.

5-Bromo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (82) and 2-(bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (76). A solution of **81** (58 mg, 0.19 mmol) in benzene (1.5 mL) was treated as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2), in order of elution, **82** (20 mg, 0.066 mmol, 34%) as a red-orange oil and **76** (18.0 mg, 0.0594 mmol, 31%) as a yellow solid. Analytical data for **82**: ^1H NMR (600 MHz) δ 7.10 (s, 1H), 4.87 (sept, $J=6.0$ Hz, 1H), 4.75 (sept, $J=6.0$ Hz, 1H), 1.32 (d, $J=6.0$ Hz, 6H), 1.31 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (150 MHz) δ 182.3, 177.1, 146.2, 145.5, 135.9, 135.3, 76.8, 76.6, 22.6; IR (ATR) 2987, 1652, 1582, 1254, 1181, 1096, 907, 802, 725 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{16}\text{BrO}_4$ ($\text{M}+\text{H}^+$) 303.0232; found: 303.0224.

2-(1-Bromo-1-iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (83).

Treatment of **81** (78 mg, 0.26 mmol) with NIS (62.4 mg, 0.276 mmol) in benzene (0.75 mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 95:5) **83** (75 mg, 0.18 mmol, 68%) as a pale yellow solid. Mp 109-113 °C; ^1H NMR (600 MHz) δ 5.49 (sept, $J=6.0$ Hz, 1H), 5.49 (sept, $J=6.0$ Hz, 1H), 1.35 (d, $J=6.0$ Hz, 6H), 1.35 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (150 MHz) δ 182.1, 181.6, 150.2, 149.6, 131.4, 75.5, 75.4, 61.5, 23.0; IR (ATR) 2983,

1679, 1617, 1304, 1084, 903, 791, 745, 718 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{15}\text{BrIO}_4$ ($\text{M}+\text{H}^+$) 428.9198; found: 428.9195.

2-(Dibromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (84). Treatment of **81** (75 mg, 0.25 mmol) with NBS (50.2 mg, 0.282 mmol) in benzene (0.75 mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2) **84** (43 mg, 0.11 mmol, 46%) as an orange solid. Mp 103-106 °C; ^1H NMR (600 MHz) δ 5.49 (sept, $J=6.0$ Hz, 2H), 1.36 (d, $J=6.0$ Hz, 12H); ^{13}C NMR (150 MHz) δ 181.3, 150.4, 128.8, 101.2, 75.5, 23.0; IR (ATR) 2987, 1682, 1613, 1378, 1304, 1084, 899, 802, 748, 671 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{14}\text{Br}_2\text{NaO}_4$ ($\text{M}+\text{Na}^+$) 402.9157; found: 402.9152.

4-Hydroxy-2-methyl-3-(1-methylethoxy)-4-(2-trimethylsilylethynyl)-2-cyclobutene-1-one (89). Butyllithium (8.29 mL, 1.6 M in hexane, 13.3 mmol) was added drop wise to a -78 °C cold solution of trimethylsilylethyne (2.7 mL, 13.9 mmol) in THF (16 mL) under a nitrogen atmosphere. The solution was stirred (2 h, -78 °C) where after it was transferred via a cannula to a -78 °C cold solution of 3-(1-methylethoxy)-4-methylcyclobut-3-ene-1,2-dione **88** (1.24 g,

7.95 mmol) in THF (40 mL). After stirring for 2 h, NH₄Cl (70 mL) was added and the mixture was extracted with Et₂O (5 x 50mL). The combined organic layers were dried (MgSO₄) and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) affording **89** (1.26 g, 4.99 mmol, 63%) as a pale yellow solid. Mp 84-88 °C; ¹H NMR (270 MHz) δ 5.04 (sept, *J*=6.1 Hz, 1H), 2.88 (s, 1H), 1.67 (s, 3H), 1.48 (d, *J*=6.1 Hz, 3H), 1.45 (d, *J*=6.3 Hz, 3H), 0.18 (s, 9H); ¹³C NMR (150 MHz) δ 186.7, 179.8, 125.0, 99.1, 95.3, 83.4, 78.3, 22.9, 22.6, 6.5, 0.4; IR (ATR) 2963, 1766, 1622, 1395, 1340, 1098, 850, 766 cm⁻¹; HRMS (ESI) calcd for C₁₃H₂₁O₃Si (M+H⁺) 253.1260; found 253.1254.

2-Trimethylsilylmethylene-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (90) and 5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (91). Treatment of **89** (65 mg, 0.26 mmol) in benzene (0.75 mL) as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2) an inseparable mixture of **90** (14 mg, 0.056 mmol, 22%, 1.2:1 ratio of isomers) and **91** (21 mg, 0.083 mmol, 32%) as a pale yellow oil. Analytical data for **90** (both isomers) from the mixture: ¹H NMR (600 MHz) δ 7.04 (s, 1H), 6.95 (s, 1H), 5.70 (sept, *J*=6.1 Hz, 1H), 5.68 (sept, *J*=6.2 Hz, 1H), 1.97 (s, 3H), 1.96 (s, 3H), 1.37 (d, *J*=6.2 Hz, 6H), 1.36 (d, *J*=6.2 Hz, 6H), 0.25 (s, 9H), 0.24 (s, 9H); Analytical data for **91** from the mixture: ¹H NMR (600 MHz) δ 6.70 (s, 1H), 4.83 (sept, *J*=6.2 Hz, 1H), 1.93 (s,

3H), 1.29 (d, $J=6.2$ Hz, 6H), 0.23 (s, 9H). Analytical data for **90** and **91**: ^{13}C NMR (150 MHz) δ 191.7, 190.7, 189.4, 188.5, 186.9, 183.1, 167.3, 166.2, 154.6, 152.2, 143.2, 142.9, 142.6, 142.5, 142.0, 75.7, 74.7, 23.2, 22.9, 9.2, 7.0, 7.0, -0.8, -1.0, -1.6; IR (ATR) 2963, 1687, 1632, 1603, 1390, 1330, 1251, 1187, 1108, 845, 766 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{21}\text{O}_3\text{Si}$ ($\text{M}+\text{H}^+$) 253.1260; found 253.1255.

2-(1-Iodo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (92) and 6-iodo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (93). Treatment of **89** (64 mg, 0.25 mmol) with NIS (67 mg, 0.30 mmol) in benzene (0.75 mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 95:5) in order of elution **92** (58.8 mg, 0.155 mmol, 61%, 1.5:1 ratio of isomers) as a yellow oil and **93** (5 mg, 0.013 mmol, 5%) as a light yellow oil. Analytical data for **92**: ^1H NMR (270 MHz) δ (600 MHz) 5.72 (sept, $J=6.1$ Hz, 1H), 5.54 (sept, $J=6.1$ Hz, 1H), 1.96 (s, 3H), 1.89 (s, 3H), 1.37 (d, $J=6.2$ Hz, 6H), 1.35 (d, $J=6.1$ Hz, 6H), 0.37 (s, 9H), 0.36 (s, 9H); ^{13}C NMR (150 MHz) δ 189.1, 188.2, 185.7, 185.6, 167.6, 163.2, 142.1, 142.0, 136.6, 132.6, 131.4, 130.9, 75.1, 74.7, 23.3, 23.1, 7.7, 6.5, 1.7, 1.4; IR (ATR) 2988, 1682, 1622, 1385, 1330, 1123, 1019, 771 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{20}\text{IO}_3\text{Si}$ ($\text{M}+\text{H}^+$) 379.0226; found 379.0222. Analytical data for **93**: ^1H NMR (270 MHz) δ 4.90 (sept, $J=6.2$ Hz, 1H), 1.91 (s, 3H), 1.41 (d, $J=6.1$ Hz, 3H), 1.32 (d, $J=6.1$ Hz, 3H), 0.23 (s, 9H); ^{13}C NMR

(150 MHz) δ 173.3, 168.5, 151.2, 107.7, 97.1, 89.0, 74.2, 22.9, 22.7, 8.9, 0.5; IR (ATR) 2988, 1781, 1662, 1306, 1251, 1108, 1053, 841 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{19}\text{INaO}_3\text{Si}$ ($\text{M}+\text{Na}^+$) 401.0046; found 401.0039.

2-(1-Bromo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (94) and 6-bromo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (95). Treatment of **89** (64 mg, 0.26 mmol) with NBS (65 mg, 0.37 mmol) in benzene (0.75 mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 95:5), in order of elution, **94** (55 mg, 0.17 mmol, 65%, 2:1 ratio of isomers) as a pale yellow oil and **95** (13 mg, 0.039 mmol, 15%) as a colorless oil. Analytical data for **94**: ^1H NMR (270 MHz, major isomer) δ 5.69 (sept, $J=6.1$ Hz, 1H), 1.90 (s, 3H), 1.37 (d, $J=5.9$ Hz, 6H), 0.36 (s, 9H); ^1H NMR (270 MHz, minor isomer) δ 5.54 (sept, $J=6.1$ Hz, 1H), 1.96 (s, 3H), 1.35 (d, $J=5.9$ Hz, 6H), 0.35 (s, 9H); ^{13}C NMR (150 MHz) δ 189.1, 187.8, 186.5, 185.1, 167.3, 164.0, 151.0, 137.9, 137.7, 136.9, 133.5, 75.0, 74.6, 23.3, 23.1, 7.4, 6.7, 0.28, 0.023; IR (ATR) 2988, 1687, 1627, 1390, 1330, 1133, 1103, 870 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{20}\text{BrO}_3\text{Si}$ ($\text{M}+\text{H}^+$) 331.0365; found 331.0362. Analytical data for **95**: ^1H NMR (270 MHz) δ 4.94 (sept, $J=6.1$ Hz, 1H), 2.01 (s, 3H), 1.45 (d, $J=6.1$ Hz, 6H), 0.55 (s, 9H); ^{13}C NMR (150 MHz) δ 171.1, 166.5, 160.9, 136.5, 114.4, 104.0, 75.7, 22.8, 21.0, 9.6, 2.0; IR (ATR) 2993, 1786, 1632, 1316, 1256, 1029, 855, 742 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{19}\text{BrNaO}_3\text{Si}$ ($\text{M}+\text{Na}^+$) 353.0185;

found 353.0176.

3-Methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (97). Treatment of **96**⁶² (64 mg, 0.35 mmol) in benzene (0.75 mL) as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 98:2) **97** (38 mg, 0.21 mmol, 85%) as a yellowish-brown oil. ¹H NMR (270 MHz) δ 6.68 (d, $J=10.1$ Hz, 1H), 6.59 (d, $J=10.1$ Hz, 1H), 4.83 (sept, $J=6.1$ Hz, 1H), 1.95 (s, 3H), 1.30 (d, $J=6.1$ Hz, 6H); ¹³C NMR (150 MHz) δ 188.4, 183.5, 154.7, 136.4, 134.8, 130.9, 76.1, 22.9, 22.5, 9.0; IR (ATR) 2983, 1652, 1593, 1385, 1306, 1192, 1098, 1019, 850, 801 cm⁻¹; HRMS (ESI) calcd for C₁₀H₁₃O₃ (M+H⁺) 181.0865; found 181.0859.

2-(1-Iodomethylene)-4-methyl-5-(1-methylethoxy)-4-cyclopentene-1,3-dione (98).

Treatment of **96** (120 mg, 0.66 mmol) and NIS (164 mg, 0.729 mmol) in toluene (5 mL) as described for **71** (150 °C, 3 h) gave after chromatography (hexanes/EtOAc, 19:1) **98** (130 mg, 0.426 mmol, 64%, 13:1 ratio of double bond isomers) as an orange solid. Mp 47-52 °C; Major isomer: ¹H NMR (270 MHz) δ 8.08 (s, 1H), 5.75 (sept, $J=6.1$ Hz, 1H), 1.91 (s, 3H), 1.38 (d, $J=6.1$ Hz, 6H); ¹³C NMR (67.5 MHz) δ 187.3, 185.8, 168.2, 137.1, 133.2, 92.1, 75.2,

23.2, 6.7; Minor isomer: ^1H NMR (270 MHz) δ 8.01 (s, 1H), 5.67 (pent, $J=6.1$ Hz, 1H), 1.95 (s, 3H), 1.35 (d, $J=6.5$ Hz, 6H); partial data ^{13}C NMR (67.5 MHz) δ 91.4, 75.0, 23.1, 7.35; IR (ATR) 1734, 1680, 1618, 1326, 1106 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{12}\text{IO}_3$ ($\text{M}+\text{H}^+$) 306.9826; found 306.9822.

2-(1-Bromomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (99).

Treatment of **96** (50 mg, 0.28 mmol) with NBS (51 mg, 0.28 mmol) in benzene (3 mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes) **99** (42 mg, 0.16 mmol, 59% as a 2.8:1 ratio of double bond isomers) as a yellow oil. ^1H NMR (270 MHz, major isomer) δ 7.58 (s, 1H), 5.74 (sept, $J=6.1$ Hz, 1H), 1.93 (s, 3H), 1.39 (d, $J=6.1$ Hz, 6H); ^1H NMR (270 MHz, minor isomer) δ 7.51 (s, 1H), 5.66 (sept, $J=6.2$ Hz, 1H), 1.97 (s, 3H), 1.36 (d, $J=6.1$ Hz, 6H); ^{13}C NMR (67.5 MHz) δ 187.2, 187.2, 183.9, 184.9, 184.7, 167.9, 164.5, 139.1, 134.3, 132.4, 132.3, 119.3, 119.0, 75.3, 75.1, 23.2, 23.1, 7.3, 6.9; IR (ATR) 2986, 1689, 1636, 1393, 1327, 1269, 1015, 764 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{12}\text{BrO}_3$ ($\text{M}+\text{H}^+$) 258.9971; found 258.9964.

2-(Diiodomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (101).

Treatment of a 0 °C cold solution of **89** (549 mg, 3.05 mmol) in DMF (30 mL) under N₂ with AgNO₃ (48 mg, 0.28 mmol) and NIS (762 mg, 3.39 mmol) was added and the reaction was slowly warmed to room temperature overnight. The mixture was cooled to 0 °C and ice-water (20 mL) was added and the resulting mixture was allowed to warm to ambient temperature. The mixture was extracted with ether (6 x 50 mL). The combined organic layers were washed with water and brine, dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 9:1) to afford **101** (209 mg, 0.483 mmol, 28%) as a yellow solid. Mp 75-80 °C; ¹H NMR (270 MHz) δ 5.65 (sept, *J*=6.1 Hz, 1H), 1.90 (s, 3H), 1.37 (d, *J*=6.1 Hz, 6H); ¹³C NMR (150 MHz) δ 187.6, 184.3, 165.0, 134.6, 134.3, 75.4, 23.2, 22.7, 7.1; IR (ATR) 2988, 1687, 1622, 1558, 1390, 1321, 1118, 1098, 900, 677 cm⁻¹; HRMS (ESI) calcd for C₁₀H₁₁I₂O₃ (M+H⁺) 432.8798; found 432.8791.

Ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-iodoethanoate (104).

Treatment of **102**¹² (60 mg, 0.25 mmol) with NBS (63.2 mg, 0.281 mmol) in benzene (0.75

mL) as described for **71** (80 °C, 3 h), gave after chromatography (hexanes/EtOAc, 9:1) **104** (65 mg, 0.18 mmol, 71%) as a yellow solid. Mp 103-107 °C; ¹H NMR (270 MHz) δ 4.38 (q, *J*=7.1 Hz, 2H), 4.29 (s, 3H), 4.24 (s, 3H), 1.35 (t, *J*=7.1 Hz, 3H); ¹³C NMR (150 MHz) δ 181.5, 180.3, 166.1, 153.8, 149.9, 131.7, 93.7, 62.9, 60.2, 60.2, 13.7; IR (ATR) 1726, 1675, 1629, 1589, 1465, 1431, 1329, 1236, 1146, 1093, 1017, 953, 852, 787 cm⁻¹; HRMS calcd for C₁₁H₁₂IO₆ (M+H⁺) 366.9679; found: 366.9676.

Ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-bromoethanoate (105).

Treatment of **102** (60 mg, 0.25 mmol) with NBS (50 mg, 0.28 mmol) in benzene (0.75 mL) as described for **71** (80 °C, 3 h), gave after chromatography (hexanes/EtOAc, 9:1) **105** (30 mg, 0.095 mmol, 38%) as a yellow oil; ¹H NMR (270 MHz) δ 4.40 (q, *J*=7.1 Hz, 2H), 4.30 (s, 3H), 4.26 (s, 3H), 1.37 (t, *J*=7.1 Hz, 3H); ¹³C NMR (150 MHz) δ 181.5, 180.3, 166.1, 153.8, 149.9, 131.7, 93.7, 62.9, 60.24, 60.17, 13.7; IR (ATR) 1734, 1686, 1642, 1594, 1467, 1339, 1255, 1155, 1103, 1027 cm⁻¹; HRMS calcd for C₁₁H₁₁BrNaO₆ (M+Na⁺) 340.9637; found 340.9634.

4.2 Supporting information Chapter 2: Total synthesis of methyl linderone

2,3-Dimethoxy-4-hydroxy-4-[(E)-4-phenyl-3-buten-1-yn-1-yl]-2-cyclobuten-1-one (137).

Butyllithium (0.82 mL, 1.30 mmol, 1.60 M in hexane) was added drop wise to a -78 °C cold solution of *E*-4-phenyl-but-3-en-1-yne (175 mg, 1.37 mmol) in THF (8 mL). The reaction mixture was stirred (-78 °C, 2 h) followed by transfer via a cannula to a -78 °C cold solution of 3,4-dimethoxy-3-cyclobutene-1,2-dione (**7**) (189 mg, 1.33 mmol) in THF (20 mL). After stirring for 2 h (-78 °C), a saturated solution of NH₄Cl (50 mL, aqueous) was added and the mixture was extracted with ether (3 x 70 mL). The combined organic layers were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **137** (170 mg, 0.628 mmol, 47%) as a yellow oil. ¹H NMR (600 MHz) δ 7.36-7.27 (5H), 7.01 (d, *J*=16.2 Hz, 1H), 6.18 (d, *J*=16.2 Hz, 1H), 4.22 (s, 3H), 3.98 (s, 3H);²⁷ ¹³C NMR (150 MHz) δ 180.8, 164.8, 143.0, 135.7, 135.5, 129.0, 128.7, 126.4, 106.5, 88.3, 84.9, 79.0, 60.2, 58.6; IR (ATR) 3365, 2958, 2180, 1772, 1627, 1351, 1046, 957, 868, 746, 685 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₅O₄ (M+H⁺) 271.0970, found: 271.0966.

4,5-Dimethoxy-[(E)-2-(1-iodo-3-phenyl-2-propen-1-ylidene)]-4-cyclopentene-1,3-dione

(138). Treatment of **137** (55 mg, 0.20 mmol) with NIS (50mg, 0.22mmol) in benzene (0.75mL) as described for **71** (80 °C, 3 h) gave after chromatography (hexanes/EtOAc, 9:1) **138** (30 mg, 0.075 mmol, 37%) as a yellow oil. ^1H NMR (600 MHz) δ 8.38 (d, $J=14.4$ Hz, 1H), 7.65 (d, $J=7.8$ Hz, 2H), 7.41-7.36 (m, 4H), 4.29 (s, 3H), 4.26 (s, 3H); ^{13}C NMR (150 MHz) δ 184.1, 184.0, 151.3, 150.8, 148.4, 135.5, 130.2, 129.0, 128.7, 126.8, 126.4, 115.4, 60.1, 60.1; IR (ATR) 2929, 1674, 1632, 1468, 1332, 1093, 1041, 760, 695 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{IO}_4$ ($\text{M}+\text{H}^+$) 396.9937; found: 396.9934.

4,5-Dimethoxy-2-[(E)-3-phenyl-2-propen-1-ylidene]-4-cyclopentene-1,3-dione (139).

Treatment of **137** (55 mg, 0.20 mmol) in benzene (0.75 mL) as described for **69** (140 °C, 3 h) gave after chromatography (hexanes/EtOAc, 85:15) **139** (30 mg, 0.11 mmol, 54%) as a yellow solid. Mp 114-119 °C; ^1H NMR (600 MHz) δ 8.11 (dd, $J=15.6$ and 11.4 Hz, 1H), 7.58 (d, $J=7.8$ Hz, 2H), 7.39-7.35 (3H), 7.13 (d, $J=12.0$ Hz, 1H), 7.09 (d, $J=15.6$ Hz, 1H), 4.28 (s, 3H), 4.27 (s, 3H); ^{13}C NMR (150 MHz) δ 186.0, 185.8, 151.8, 150.3, 147.0, 136.1, 135.8, 130.0, 128.9, 128.0, 124.5, 123.2, 59.9, 59.9; IR (ATR) 2962, 1669, 1632, 1608, 1468, 1435,

1322, 1102, 1060, 999, 788, 760 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{O}_4$ ($\text{M}+\text{H}^+$): 271.0970; Found: 271.0967.

4,5-Dimethoxy-[(E)-2-(1-methoxy-3-phenyl-2-propen-1-ylidene)]-4-cyclopentene-1,3-dione (methyl linderone). Treatment of **138** (28 mg, 0.072 mmol) with K_2CO_3 (10 mg, 0.073 mmol) in MeOH (5 mL) at 60°C for 3 days gave after solvent removal and chromatography (hexanes/EtOAc, 9:1) methyl linderone (19 mg, 0.064 mmol, 89%) as a yellow oil. ^1H NMR and ^{13}C NMR data were in complete accordance with literature values.¹⁵

4,5-Dimethoxy-2-[(E)-1-(N,N-dimethylamino)-3-phenyl-2-propen-1-ylidene]-4-cyclopentene-1,3-dione (140). Treatment of **138** (10 mg, 0.0263 mmol) with *p*-toluenesulfonic acid (0.2 mg, 0.001 mmol) in a mixture of DMF (1.8 mL) and H_2O (0.2 mL) at 100°C . Water (5 mL) was added and the mixture was extracted with EtOAc (3 x 50 mL). The combined organic phases were dried (MgSO_4) and filtered. The solvent was removed under reduced pressure. Purification by chromatography (hexanes/EtOAc, 85:15) gave **140** (8 mg, 0.025 mmol, 95%) as a yellow oil. ^1H NMR (600 MHz) δ 7.52 (dd, $J=7.8$ and 1.8 Hz, 2H), 7.39-7.33 (m, 3H),

7.04 (d, $J=16.2$ Hz, 1H), 6.75 (d, $J=16.2$ Hz, 1H), 4.05 (s, 6H), 3.21 (s, 6H); ^{13}C NMR (150 MHz) δ 187.5, 160.5, 145.0, 142.6, 139.3, 135.2, 129.9, 128.8, 128.0, 120.9, 96.5, 59.8, 44.0; IR (ATR) 2929, 1646, 1543, 1482, 1407, 1313, 1004, 765 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{19}\text{NNaO}_4$ ($\text{M}+\text{Na}^+$) 336.1212; found: 336.1210.

4.3 Supporting information Chapter 3: Wacker-type oxidation of aryl-substituted alkenes

2-Nitro-1-(1-propen-1-yl)benzene (230).¹⁶⁷ To a solution of Pd(dba)₂ (870 mg, 1.51 mmol) and PPh₃ (1.60 g, 6.08 mmol) in toluene (80 mL) under N₂ atmosphere was added tributyl-1-propen-1-yltin¹⁶⁸ (12.62 g, 38.12 mmol) and 1-iodo-2-nitrobenzene (7.55 g, 30.3 mmol). The mixture was heated at reflux for 48 h. The solvent was removed under reduced pressure and Et₂O (200 mL) was added. The solution was washed with NH₄OH (10%-aqueous, 6 x 100 mL), H₂O (3 x 100 mL), brine (100 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 99:1) to give **230** (3.02 g, 18.5 mmol, 49%) as a pale yellow oil.

3-Nitro-1-(1-propen-1-yl)benzene (231).¹⁶⁹ Treatment of 1-bromo-3-nitrobenzene (716 mg, 3.55 mmol) with tributyl-1-propen-1-yltin (1.40 g, 4.22 mmol) a solution of Pd(dba)₂ (105 mg, 0.183 mmol) and PPh₃ (183mg, 0.699 mmol) in toluene (50 mL), as described for **1**, gave after work up and chromatography (hexanes/EtOAc, 98:2) **231** (339 mg, 2.08 mmol, 49%) as

a pale yellow oil.

4-Nitro-1-(1-propen-1-yl)benzene (232).¹⁷⁰ Treatment of 1-bromo-4-nitrobenzene (713 mg, 3.53 mmol) with tributyl 1-propen-1-yltin (1.40 g, 4.23 mmol) a solution of Pd(dba)₂ (107 mg, 0.186 mmol) and PPh₃ (183mg, 0.698 mmol) in toluene (50 mL), as described for **230**, gave after work up and chromatography (hexanes/EtOAc, 99:1) **232** (396 mg, 2.43 mmol, 57%) as a pale yellow oil.

4-Fluoro-2-nitro-1-(1-propen-1-yl)benzene (233). Treatment of 4-fluoro-2-nitro-iodobenzene (967 mg, 3.62 mmol) with tributyl 1-propen-1-yltin (1.45 g, 4.38 mmol) a solution of Pd(dba)₂ (106 mg, 0.185 mmol) and PPh₃ (194 mg, 0.740 mmol) in toluene (50 mL), as described for **230**, gave after work up and chromatography (hexanes/EtOAc, 98:2) **233** (579 mg, 3.20 mmol, 73%) as a pale yellow oil. Spectral data from a 1.9:1 mixture of *E/Z*-**233**. *E*-**233**: ¹H NMR (600 MHz) δ 7.61 (dd, *J*=8.4, 2.4 Hz, 1H), 7.57 (dd, *J*=8.4, 5.4 Hz, 1H), 7.26 (dt, *J*=8.4, 2.4 Hz, 1H), 6.81 (d, *J*=15.6 Hz, 1H), 6.19 (dq, *J*=15.6, 7.2 Hz, 1H), 1.93 (dd, *J*=7.2, 1.8 Hz, 3H); ¹³C NMR δ (150 MHz) 160.7 (d, *J*_{CF}=248.7 Hz), 147.4, 131.6, 130.0 (d, *J*_{CF}=8.1 Hz), 129.7 (d, *J*_{CF}=3.8 Hz), 125.2, 120.4 (d, *J*_{CF}=21.3 Hz), 111.6 (d, *J*_{CF}=26.1 Hz),

18.7. **Z-233**: ^1H NMR δ (600 MHz) 7.74 (dd, $J=8.4, 2.4$ Hz, 1H), 7.37 (dd, $J=8.4, 5.4$ Hz, 1H), 7.31 (dt, $J=8.4, 2.4$ Hz, 1H), 6.66 (d, $J=11.4$ Hz, 1H), 5.94 (dq, $J=11.4, 7.2$ Hz, 1H), 1.71 (dd, $J=7.2, 1.8$ Hz, 3H); ^{13}C NMR (150 MHz) δ 160.7 (d, $J_{\text{CF}}=249.3$ Hz), 148.7, 133.4 (d, $J_{\text{CF}}=7.5$ Hz), 129.2, 128.7 (d, $J_{\text{CF}}=4.4$ Hz), 125.1, 120.0 (d, $J_{\text{CF}}=20.7$ Hz), 112.0 (d, $J_{\text{CF}}=26.1$ Hz), 14.2. IR (ATR) 2918, 1527, 1490, 1343, 1240, 1131, 944, 874, 816, 789, 753, 676 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_9\text{H}_9\text{FNO}_2$ ($\text{M}+\text{H}^+$) 182.0617; found 182.0612.

2,4-Dinitro-1-(1-propen-1-yl)benzene (234). Treatment of 1-bromo-2,4-dinitrobenzene (865 mg, 3.50 mmol) with tributyl 1-propen-1-yltin (1.40 g, 4.24 mmol) a solution of $\text{Pd}(\text{dba})_2$ (101 mg, 0.175 mmol) and PPh_3 (186 mg, 0.711 mmol) in toluene (50 mL), as described for **230**, gave after work up and chromatography (hexanes/EtOAc, 9:1) **234** (453 mg, 2.18 mmol, 51%) as a pale orange-brown oil. Spectral data from a 1:1.8 mixture of *E/Z*-**234**. *E*-**234**: ^1H NMR (600 MHz) δ 8.74 (d, $J=2.4$ Hz, 1H), 8.36 (dd, $J=9.0, 2.4$ Hz, 1H), 7.79 (d, $J=9.0$ Hz, 1H), 6.91 (dd, $J=15.6, 1.8$ Hz, 1H), 6.51 (dq, $J=15.6, 6.6$ Hz, 1H), 2.02 (dd, $J=6.6, 1.8$ Hz, 3H). *Z*-**234**: ^1H NMR (600 MHz) δ 8.86 (d, $J=2.4$ Hz, 1H), 8.42 (dd, $J=8.4, 2.4$ Hz, 1H), 7.62 (d, $J=8.4$ Hz, 1H), 6.76 (dd, $J=12.0, 1.8$ Hz, 1H), 6.13 (dq, $J=12.0, 7.2$ Hz, 1H), 1.79 (dd, $J=7.2, 1.8$ Hz, 6H). IR (ATR) 2918, 1596, 1522, 1446, 1340, 959, 915, 834, 767, 736, 716, 698 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_9\text{H}_9\text{N}_2\text{O}_4$ ($\text{M}+\text{H}^+$) 209.0562; found 209.0557.

4-Carbomethoxy-2-nitro-1-(1-propen-1-yl)benzene (235). 1-Bromo-4-carbomethoxy-2-nitrobenzene (865 mg, 3.50 mmol) was treated with Pd(dba)₂ (84mg, 0.15 mmol), PPh₃ (154 mg, 0.587mmol), and tributyl 1-propen-1-yltin (1.17 g, 3.52mmol) in toluene (50 mL) as described for **230**. Work up and purification by chromatography (hexanes/EtOAc, 95:5) gave **235** (364 mg, 1.64 mmol, 56%) as a pale yellow oil. Spectral data from a 1:1.2 mixture of *E/Z*-**235**. *E*-**235**: ¹H NMR (270 MHz) δ 8.50 (d, *J*=1.6 Hz, 1H), 8.15 (dd, *J*=7.9, 1.6 Hz, 1H), 7.66 (d, *J*=8.3 Hz, 1H), 6.87 (dd, *J*=15.4, 1.8 Hz, 1H), 6.41 (dq, *J*=15.4, 6.7 Hz, 1H), 3.95 (s, 3H), 1.98 (dd, *J*=6.7, 1.8 Hz, 3H). *Z*-**235**: ¹H NMR (270 MHz) δ 8.63 (d, *J*=1.6 Hz, 1H), 8.21(dd, *J*=8.1, 1.6 Hz, 1H), 7.48 (d, *J*=8.1 Hz, 1H), 6.73 (dd, *J*=11.5, 1.8 Hz, 1H), 6.03 (dq, *J*=11.5, 7.1 Hz, 1H), 3.97 (s, 3H), 1.76 (dd, *J*=7.1, 1.8Hz, 3H). HRMS (ESI) calcd for C₁₁H₁₂NO₄ (M+H⁺) 222.0766; found 222.0761.

4-Methoxy-2-nitro-1-(1-propen-1-yl)benzene (236). 4-Methoxy-2-nitro-1-iodobenzene (884 mg, 3.17 mmol) was treated with Pd(dba)₂ (91 mg, 0.16 mmol), PPh₃ (166 mg, 0.634 mmol), and tributyl 1-propen-1-yltin (1.27 g, 3.80 mmol) in toluene (50 mL) as described for **230**. Work up and purification by chromatography (hexanes/EtOAc, 95:5) gave **236** (195 mg, 1.01mmol, 32%) as a pale yellow oil. Spectral data from a 1.7:1 mixture of *E/Z*-**236**. *E*-**236**:

^1H NMR (270 MHz) δ 7.48 (d, $J=8.7$ Hz, 1H), 7.38 (d, $J=2.6$ Hz, 1H), 7.08 (dd, $J=8.1$, 2.8 Hz, 1H), 6.78 (dd, $J=15.7$, 1.8 Hz, 1H), 6.13 (dq, $J=15.4$, 6.5 Hz, 1H), 3.85 (s, 3H), 1.91 (dd, $J=6.5$, 1.8 Hz, 3H); ^{13}C NMR (67.5 MHz) δ 158.5, 147.9, 129.7, 129.3, 126.0, 125.7, 120.0, 108.4, 55.8, 18.7. **Z-236**: ^1H NMR (270 MHz) δ 7.52 (d, $J=2.6$ Hz, 1H), 7.28 (d, $J=8.7$ Hz, 1H), 7.12 (overlapping *E*-isomer, 1H), 6.64 (dd, $J=11.3$, 1.8 Hz, 1H), 5.88 (dq, $J=11.5$, 7.1 Hz, 1H), 3.88 (s, 3H), 1.71 (dd, $J=7.1$, 1.8 Hz, 3H); ^{13}C NMR (67.5 MHz) δ 158.6, 148.8, 132.7, 128.1, 125.7, 125.0, 119.4, 108.9, 55.8, 14.3. IR (ATR) 2915, 2841, 1619, 1522, 1346, 1286, 1253, 1033, 961, 839, 814, 784, 679 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{12}\text{NO}_3$ ($\text{M}+\text{H}^+$) 194.0817; found 194.0812.

5-Methoxy-2-nitro-1-(1-propen-1-yl)-benzene (237). 5-Methoxy-2-nitro-1-iodobenzene (283 mg, 1.01 mmol) was treated with $\text{Pd}(\text{dba})_2$ (29.8 mg, 0.052 mmol), PPh_3 (57.6 mg, 0.22 mmol), and tributyl 1-propen-1-yltin (411 mg, 1.24 mmol) in toluene (50 mL) as described for **230**. Work up and purification by chromatography (hexanes/EtOAc, 8:2) affording **237** (89.7 mg, 0.464 mmol, 37%) as a pale yellow oil. Spectral data from a 1.8:1 mixture of *E/Z*-**237**. **E-237**: ^1H NMR (600 MHz) δ 7.99 (d, $J=9.6$ Hz, 1H), 6.99 (dd, $J=15.6$, 1.2 Hz, 1H), 6.96 (d, $J=2.4$ Hz, 1H), 6.19 (dq, $J=15.6$, 6.6 Hz, 1H), 3.89 (s, 3H), 1.94 (dd, $J=6.6$, 1.2 Hz, 3H). **Z-237**: ^1H NMR (600 MHz) δ 8.11 (d, $J=9.6$ Hz, 1H), 6.86 (dd, $J=9.0$, 2.4 Hz, 1H), 6.77 (dd, $J=12$, 1.8 Hz, 1H), 5.91 (dq, $J=12$, 7.2 Hz, 1H), 3.89 (s, 3H), 1.74 (dd, $J=7.2$, 1.8 Hz, 3H). 6.82-6.80 (m, 2H) could not be assigned to the specific proton for the two isomers. IR

(ATR) 2917, 2845, 1620, 1524, 1351, 1285, 1255, 1034, 962, 837, 813, 783, 678 cm^{-1} ;
 HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{11}\text{NNaO}_3$ ($\text{M}+\text{H}^+$) 216.0631; found 216.0630.

2-Bromo-1-(1-propen-1-yl)-benzene (243).¹⁷¹ 2-Bromo-1-iodobenzene (1.02g, 3.10 mmol) was treated with $\text{Pd}(\text{dba})_2$ (72.2 mg, 0.126 mmol), PPh_3 (132 mg, 0.504 mmol), and tributyl 1-propen-1-yltin (728 mg, 2.57 mmol) in toluene (50 mL) as described for **230**. Work up and purification by chromatography (hexanes/EtOAc, 8:2) gave **243** (203 mg, 1.03 mmol, 40%) as a pale yellow oil.

Typical oxidation procedure using Condition A: 1-(2-Nitrophenyl)-2-propanone (245),¹⁷² **1-(1,1-dimethoxymethyl)-2-nitrobenzene (246),**¹⁷³ **2-nitrobenzaldehyde (247),**¹⁷⁴ **and 1-(2-nitrophenyl)-1,2-propandione (248).**¹⁷⁵ To a suspension of PdCl_2 (15.5 mg, 0.087mmol) and CuCl (69 mg, 0.70 mmol) in a solution of MeOH (1 mL) and H_2O (200 μL) was added **230** (100 mg, 0.614mmol) dissolved in MeOH (1 mL). The reaction flask was attached to a balloon filled with oxygen and the mixture was stirred at 50 $^\circ\text{C}$ for 20 h. The resulting mixture was diluted with H_2O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO_4), filtered, and the solvents were removed under reduced

pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **246** (4.9 mg, 0.025 mmol, 4%), an inseparable mixture of **247** and **248** (1:1 ratio, 13.9 mg; 6.10 mg, 0.0404 mmol, 6.4% **247**; 7.80 mg, 0.0404 mol, 6.4% **248**) and **245** (79.9 mg, 0.45 mmol, 73%) all as a pale yellow oils.

Typical oxidation procedure using Condition B: 1-(2-Nitrophenyl)-2-propanone (245) and 2-nitrobenzaldehyde (247). To a suspension of PdCl₂ (10.2 mg, 0.058) in H₂O₂ (50%-aqueous, 300 μ L, 6.40 mmol) was added **230** (99.2 mg, 0.608 mmol). After 48 h at 50 °C, the resulting mixture worked up and purified as described using Conditions A to give after chromatography (hexanes/EtOAc, 8:2) in order of elution **247** (7.4 mg, 0.049mmol, 8%) and **245** (20.1 mg, 0.11 mmol, 18%) both as pale yellow oils.

Typical oxidation procedure using Condition C: 1-(2-Nitrophenyl)-2-propanone (245), 1-(1,1-dimethoxymethyl)-2-nitrobenzene (246), and 2-nitrobenzaldehyde (247). To a suspension of PdCl₂ (14.2 mg, 0.080 mmol) and 2,6-di-tert-butyl-p-cresol (15.0 mg, 0.068 mmol) in a solution of MeOH (1 mL), H₂O (200 μ L) and H₂O₂ (50%-aqueous, 300 μ L, 6.40 mmol) was added to **230** (106 mg, 0.648 mmol) dissolved in MeOH (1 mL). The reaction

mixture was stirred at 50 °C for 48 h. The resulting mixture worked up and purified as described using Conditions A to give in order of elution after chromatography (hexanes/EtOAc, 8:2) **246** (2.3 mg, 0.012 mmol, 2%), **247** (20.8 mg, 0.138 mmol, 21%), and **245** (83.4 mg, 0.466 mmol, 72%) all as pale yellow oils.

Typical oxidation procedure using Condition D: 1-(2-Nitrophenyl)-2-propanone (245). To a suspension of PdCl₂ (5.8 mg, 0.033 mmol) and 2,6-di-tert-butyl-p-cresol (6.3 mg, 0.029 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and *t*-butyl hydrogenperoxide (TBHP, 90%-aqueous, 267 μL, 2.49 mmol) was added **230** (31.0 mg, 0.190 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **245** (31.3 mg, 0.175 mmol, 92%) as a pale yellow oil.

1-(1,1-Dimethoxymethyl)-3-nitrobenzene (**249**),¹⁷⁶ *3-nitrobenzaldehyde* (**250**),¹⁷⁷ and *1-(3-nitrophenyl)-1,2-propanedione* (**251**).¹⁷⁸ A solution of **231** (116 mg, 0.709 mmol) in MeOH (1 mL) was added to a suspension of PdCl₂ (12.5 mg, 0.0705 mmol) and CuCl (69.0 mg, 0.70 mmol) in a solution of MeOH (1 mL) and H₂O (200 μ L) under an atmosphere of oxygen, as described for **230** in Condition A (50 $^{\circ}$ C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 8:2) gave in order of elution **249** (41.4 mg, 0.210 mmol, 30%) as a colorless oil and an inseparable mixture of **250** and **251** (1:1 ratio, 17.2 mg; 7.55 mg, 0.050 mmol, 7% **250**; 9.66 mg, 0.050 mmol, 7% **251**) as a yellow solid. Analytical data for **251**: ¹H NMR (600 MHz) δ 8.89 (t, J =2.4 Hz, 1H), 8.48 (dt, J =7.2, 2.4 Hz, 1H), 8.40 (td, J =8.4, 1.8 Hz, 1H), 7.71 (t, J =8.4 Hz, 1H), 2.58 (s, 3H); ¹³C NMR (150 MHz) 198.5, 187.7, 148.4, 135.7, 133.3, 130.0, 128.4, 125.3, 26.0; IR (ATR) 1704, 1615, 1533, 1352, 1170, 1093, 818, 711, 677 cm⁻¹.

1-(3-Nitrophenyl)-2-propanone (**252**).¹⁷⁹ To a suspension of PdCl₂ (5.6 mg, 0.032 mmol) and 2,6-di-tert-butyl-p-cresol (6.1 mg, 0.028 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and TBHP (90%-aqueous, 267 μ L, 2.49 mmol) was added **231** (40.6 mg, 0.249 mmol)

dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 $^{\circ}$ C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **252** (15.3 mg, 0.140 mmol, 34%) as a pale yellow oil.

1-(1,1-Dimethoxymethyl)-4-nitrobenzene (**253**),¹⁸⁰ *4-nitrobenzaldehyde* (**254**),¹⁷⁷ and *1-(4-nitrophenyl)-1,2-propanedione* (**255**).¹⁸¹ A solution of **232**¹⁸² (57.0 mg, 0.349 mmol) in MeOH (500 μ L) was added to a suspension of PdCl₂ (6.3 mg, 0.036 mmol) and CuCl (35.2 mg, 0.36 mmol) in a solution of MeOH (500 μ L) and H₂O (100 μ L) under an atmosphere O₂, as described for **230** in Condition A (50 $^{\circ}$ C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 95:5) gave in order of elution, **253** (21.8 mg, 0.11 mmol, 32%) as a colorless oil and an inseparable mixture of **254** and **255** (1:1 ratio, 11.7 mg; 5.1 mg, 0.034 mmol, 10% **254**; 6.6 mg, 0.034 mmol, 10% **255**) as a yellow solid.

1-(1,1-Dimethoxymethyl)-4-nitrobenzene (**253**), *4-nitrobenzaldehyde* (**254**), and *1-(4-nitrophenyl)-1,2-propanedione* (**255**). To a suspension of PdCl₂ (7.5 mg, 0.042 mmol) and 2,6-di-tert-butyl-p-cresol (7.9 mg, 0.036 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and H₂O₂ (50%-aqueous, 150 μL, 3.20 mmol) was added **232** (56.6 mg, 0.347 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **253** (23.0 mg, 0.117 mmol, 34%) as a colorless oil and an inseparable mixture of **254** and **255** (1:1 ratio, 8.7 mg; 3.8 mg, 0.025 mmol, 7% **254**; 4.9 mg, 0.025 mmol, 7% **255**) as a yellow solid.

1-(1,1-Dimethoxymethyl)-4-nitrobenzene (**253**) and *1-(4-nitrophenyl)-2-propanone* (**256**).¹⁸³ To a suspension of PdCl₂ (6.6 mg, 0.037 mmol) and 2,6-di-tert-butyl-p-cresol (6.1 mg, 0.028 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and TBHP (90%-aqueous, 267 mL, 2.49 mmol) was added **232** (40.6 mg, 0.249 mmol) dissolved in MeOH (500 μL).

The reaction mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to give in order of elution **253** (0.9 mg, 0.0046 mmol, 2%) as a colorless oil and **256** (24.7 mg, 0.138 mmol, 55%) as a pale yellow oil.

1-(4-Fluoro-2-nitrophenyl)-2-propanone (257), 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (258), 4-fluoro-2-nitrobenzaldehyde(259), and 1-(4-fluoro-2-nitrophenyl)-1,2-propanedione (260). A solution of **233** (65.2 mg, 0.36 mmol) in MeOH (500 μ L) was added to a suspension of PdCl₂ (6.5 mg, 0.037 mmol) and CuCl (34.8 mg, 0.352 mmol) in a solution of MeOH (500 μ L) and H₂O (100 μ L) under an atmosphere of oxygen, as described for **230** in Condition A (50 °C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 9:1) gave in order of elution, **258** (2.2 mg, 0.010mmol, 3%) as a colorless oil, an inseparable mixture of **259** and **260** (1:1 ratio, 6.8 mg; 3.0 mg, 0.018 mmol, 5% **259**; 3.8 mg, 0.018 mmol, 5% **260**) as a pale yellow oil, and **257** (39.2 mg, 0.199 mmol, 55%) as a pale yellow oil. Spectral data for **257**: ¹H NMR (600 MHz) δ 7.86 (dd, J =8.4, 2.4 Hz, 1H), 7.32 (dt, J =9.0, 3.0 Hz, 1H), 7.57 (dd, J =9, 5.4 Hz, 1H), 4.10 (s, 2H), 2.33 (s, 3H); ¹³C NMR (150 MHz) δ 203.2, 162.2, 160.5, 134.8 (d, J_{CF} =8.0 Hz), 126.3 (d,

$J_{CF}=4.7$ Hz), 120.8 (d, $J_{CF}=20.6$ Hz), 112.9 (d, $J_{CF}=26.3$ Hz), 47.9, 29.9. IR (ATR) 1720, 1532, 1499, 1360, 1237, 1163, 944, 878, 807 cm^{-1} . Spectral data for **258**: ^1H NMR (600 MHz) δ 7.80 (dd, $J=9.0, 5.4$ Hz, 1H), 7.56 (dd, $J=8.4, 2.4$ Hz, 1H), 7.31 (td, $J=7.8, 2.4$ Hz, 1H), 5.88 (s, 1H), 3.40 (s, 6H); ^{13}C NMR (150 MHz) δ 161.9 (d, $J_{C,F}=251.3$ Hz), 149.4, 130.1 (d, $J_{C,F}=8.1$ Hz), 129.0, 119.5 (d, $J_{C,F}=20.6$ Hz), 112.1 (d, $J_{C,F}=26.4$ Hz), 99.6, 54.6; IR (ATR) 2961, 1736, 1540, 1364, 1262, 1233, 1105, 1058, 877, 806 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_9\text{H}_{11}\text{FO}_4$ ($\text{M}+\text{H}^+$) 216.0672; found 216.0667. Spectral data for diketone **260** from the mixture with **259** (1.26:1 ratio): ^1H NMR (600 MHz) δ 7.88 (dd, $J=7.8, 1.8$ Hz, 1H), 7.63 (dd, $J=8.4, 5.4$ Hz, 1H), 7.53-7.48 (m, 1H), 2.56 (s, 3H); Spectral data for mixture of **259** and **260**: ^{13}C NMR (150 MHz) δ 195.9, 188.5, 186.5, 165.6, 164.7, 163.9, 163.0, 132.3, 132.2, 132.2, 132.2, 128.2, 128.2, 127.5, 127.5, 122.1, 122.0, 121.4, 121.2, 112.6, 112.4, 112.0, 111.8, 24.2; IR (ATR) 3099, 1701, 1610, 1535, 1347, 1219, 1231, 1167, 882, 826, 809 cm^{-1} . **259**: HRMS (ESI) calcd for $\text{C}_7\text{H}_4\text{FNNaO}_3$ ($\text{M}+\text{H}^+$) 192.0073; found 192.0069. **260**: HRMS (ESI) calcd for $\text{C}_9\text{H}_7\text{FNO}_4$ ($\text{M}+\text{H}^+$) 212.0359; found 212.0354.

1-(4-Fluoro-2-nitrophenyl)-2-propanone (257) and 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (258). Compound **233** (47.6 mg, 0.263 mmol) dissolved in MeOH (500 μL) was added to a suspension of PdCl_2 (7.8 mg, 0.044 mmol), 2,6-di-tert-butyl-p-cresol (7.8 mg, 0.035 mmol), and H_2O_2 (50%-aqueous, 150 μL , 3.20 mmol) in MeOH (500 μL) and H_2O

(100 μ L). After 48 h at 50 $^{\circ}$ C, the mixture was worked up and purified as described in Condition C to give after chromatography (hexanes/EtOAc, 9:1), in order of elution, **258** (9.4 mg, 0.044 mmol, 17%) and **257** (18.3 mg, 0.093 mmol, 35%) both as a yellow oils.

1-(4-Fluoro-2-nitrophenyl)-2-propanone (257). Compound **233** (45.8 mg, 0.253 mmol) dissolved in MeOH (500 μ L) was added to a suspension of PdCl₂ (5.8 mg, 0.033 mmol), 2,6-di-tert-butyl-p-cresol (5.7 mg, 0.026 mmol) and TBHP (90%-aqueous, 267 μ L, 2.49 mmol) in a solution of MeOH (500 μ L) and H₂O (100 μ L). After 48 h at 50 $^{\circ}$ C, the mixture was worked up and purified as described in Conditions D to give after chromatography (hexanes/EtOAc, 9:1) **257** (26.7 mg, 0.135mmol, 54%) as a pale yellow oil.

Methyl 2,4-dinitrobenzoate (261),¹⁸⁴ **2,4-dinitrobenzaldehyde (262),**¹⁸⁶ **1-(2,4-dinitrophenyl)-1,2-propanedione (263).** A solution of **234** (68.7 mg, 0.330 mmol) in MeOH (500 μ L) was added to a suspension of PdCl₂ (7.0 mg, 0.039mmol) and CuCl (35.4mg, 0.358mmol) in a solution of MeOH (500 μ L) and H₂O (100 μ L) under an atmosphere of oxygen, as described for **230** in Condition A (50 $^{\circ}$ C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 95:5) gave in order of elution **263** (23.5 mg, 0.096mmol, 29%), **261** (9.6 mg,

0.042mmol, 13%), and **262** (25.3 mg, 0.129 mmol, 39%) all as pale yellow oils. Spectral data for **263**: ^1H NMR (600 MHz) δ 9.02 (dd, $J=2.4$, 1H), 8.65 (dd, $J=8.4$, 2.4 Hz, 1H), 7.79 (dd, $J=8.4$ Hz, 1H), 2.60 (s, 1H); ^{13}C NMR (150 MHz) δ 195.3, 187.4, 149.5, 147.6, 137.9, 131.4, 129.1, 119.4, 24.0; IR (ATR) 1717, 1601, 1536, 1345, 1170, 903, 835, 736, 721 cm^{-1} .

1-(2,4-Dinitrophenyl)-2-propanone (264).¹⁸⁵ To a suspension of PdCl_2 (7.8 mg, 0.044 mmol) and 2,6-di-tert-butyl-p-cresol (7.5 mg, 0.034 mmol) in a solution of MeOH (500 μL), H_2O (100 μL) and H_2O_2 (50%-aqueous, 150 μL , 3.20 mmol) was added **234** (40.9 mg, 0.196mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 $^\circ\text{C}$ for 48 h. The mixture was diluted with H_2O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO_4), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to give **264** (13.5 mg, 0.060 mmol, 31%) as a yellow oil.

1-(4-Carbomethoxy-2-nitrophenyl)-2-propanone (265),¹⁸⁶ **4-carbomethoxy-2-nitrobenzaldehyde (266),**¹⁸⁷ and **1-(4-carbomethoxy-2-nitrophenyl)-1,2-propanedione (267).**

A solution of **235** (68.2 mg, 0.308 mmol) in MeOH (500 μL) was added to a suspension of

PdCl₂ (6.5 mg, 0.037 mmol) and CuCl (30.3mg, 0.306 mmol) in a solution of MeOH (500 μ L) and H₂O (100 μ L) under an atmosphere of oxygen, as described **230** in Condition A (50 °C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 95:5) gave in order of elution an inseparable mixture of **266** and **267** (1.1:1 ratio, 26.9 mg; 12.9 mg, 0.0615 mmol, 20 % **266**; 14.0 mg, 0.0559 mmol, 18% **267**) and **265** (21.5mg, 0.0906 mmol, 29%) both as pale yellow oils.

1-(4-Carbomethoxy-2-nitrophenyl)-2-propanone (265), 4-carbomethoxy-2-nitrobenzaldehyde (266), and 1-(4-carbomethoxy-2-nitrophenyl)-1,2-propanedione (267).

To a suspension of PdCl₂ (5.0 mg, 0.028 mmol) and 2,6-di-tert-butyl-p-cresol (5.8 mg, 0.026 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and H₂O₂ (50%-aqueous, 150 μ L, 3.20 mmol) was added **235** (55.5 mg, 0.251 mmol) dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to give in order of elution an inseparable mixture of **266** and **267** (1.1:1 ratio, 5.2 mg; 2.7 mg, 0.012 mmol, 5% **266**; 2.7 mg, 0.011 mmol, 4% **267**) and **265** (22.3mg, 0.094mmol, 37%) all as pale yellow oils.

1-(4-Carbomethoxy-2-nitrophenyl)-2-propanone (265), 4-carbomethoxy-2-nitrobenzaldehyde (266) and 1-(4-carbomethoxy-2-nitrophenyl)-1,2-propanedione (267).

To a suspension of PdCl₂ (5.8 mg, 0.033 mmol) and 2,6-di-tert-butyl-p-cresol (5.5 mg, 0.025 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and TBHP (90%-aqueous, 267 μL, 2.49 mmol) was added **235** (55.3 mg, 0.250 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 9:1) to give in order of elution an inseparable mixture of **266** and **267** (1.3:1 ratio, 3.4 mg; 1.8 mg, 0.0084 mmol, 4% **266**; 1.63 mg, 0.0065 mmol, 3% **267**) and **265** (12.5 mg, 0.0527 mmol, 21%) as pale yellow oils. Partial spectral data of **267** from the mixture: ¹H NMR (600 MHz) δ 8.81 (d, *J*=1.6 Hz, 1H), ~8.4 (overlapping with **266**), 6.96 (d, *J*=3.0 Hz, 1H), 4.01 (s, 3H), 2.58 (s, 3H).

1-(4-Methoxy-2-nitrophenyl)-2-propanone (268).¹⁸⁸ A solution of **236** (41.1 mg, 0.213 mmol) in MeOH (500 μL) was added to a suspension of PdCl₂ (4.1 mg, 0.023 mmol) and CuCl (22.6 mg, 0.2283 mmol) in a solution of MeOH (500 μL) and H₂O (100 μL) under O₂, as described

for **230** in Condition A (50 °C, 20h). Extraction and purification by chromatography (hexanes/EtOAc, 95:5) gave **268** (23.7 mg, 0.113 mmol, 53%) as a pale yellow oil.

1-(4-Methoxy-2-nitrophenyl)-2-propanone (268). To a suspension of PdCl₂ (5.6 mg, 0.032 mmol) and 2,6-di-tert-butyl-p-cresol (5.5 mg, 0.025 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and TBHP (90%-aqueous, 267 μL, 2.49 mmol) was added **236** (40.3 mg, 0.209 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 9:1) to give **268** (23.8 mg, 0.114 mmol, 55%) as a pale yellow oil.

1-(5-Methoxy-2-nitrophenyl)-2-propanone (269), 5-methoxy-2-nitro-benzaldehyde (270),¹⁸⁹ and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (271). A solution of **237** (65.0 mg, 0.336 mmol) in MeOH (500 μL) was added to a suspension of PdCl₂ (6.2 mg, 0.035 mmol) and CuCl (34.9 mg, 0.352 mmol) in a solution of MeOH (500 μL) and H₂O (100 μL) under O₂, as described for **230** in Condition A (50 °C, 20 h). Extraction and purification by

chromatography (hexanes/EtOAc, 95:5) gave in order of elution an inseparable mixture of **270** and **271** (1.3:1 ratio, 11.5 mg; 5.9 mg, 0.033 mmol, 10% **270**; 5.6 mg, 0.0251 mmol, 7% **271**) and **269** (28.7 mg, 0.137 mmol, 41%) both as pale yellow oils. Spectral data for **269**: ^1H NMR (600 MHz) δ 8.19 (d, $J=9.0$ Hz, 1H), 6.89 (dd, $J=9.6$, 3.0 Hz, 1H), 6.44 (d, $J=2.4$ Hz, 1H), 4.09 (s, 2H), 3.88 (s, 3H), 2.33 (s, 3H); ^{13}C NMR (150 MHz) δ 203.7, 163.4, 141.4, 133.5, 128.1, 118.7, 112.9, 55.9, 49.2, 30.0; IR (ATR) 1721, 1581, 1509, 1335, 1292, 1261, 1162, 1085, 843 cm^{-1} . Spectral data for **271** from the mixture of **270/271**: ^1H NMR (600 MHz) δ 8.17 (d, $J=9.0$ Hz, 1H), 7.10 (dd, $J=9.0$, 3.0 Hz, 1H), 6.96 (d, $J=3.0$ Hz, 1H), 3.94 (s, 3H), 2.57 (s, 3H); ^{13}C NMR (100 MHz) δ 196.0, 189.6, 164.7, 135.6, 126.6, 116.7, 114.6, 56.4, 24.2; IR (ATR) 2945, 1706, 1582, 1513, 1485, 1333, 1290, 1239, 1190, 1026, 933, 849, 743 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_9\text{NNaO}_5(\text{M}+\text{Na}^+)$ 246.0378; found 246.0374.

1-(5-Methoxy-2-nitrophenyl)-2-propanone (269), 5-methoxy-2-nitro-benzaldehyde (270), and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (271). To a suspension of PdCl_2 (5.7 mg, 0.032 mmol) and 2,6-di-tert-butyl-p-cresol (5.6 mg, 0.025 mmol) in a solution of MeOH (500 μL), H_2O (100 μL) and H_2O_2 (50%-aqueous, 150 μL , 3.20 mmol) was added **237** (20.1 mg, 0.104 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 $^\circ\text{C}$ for 48 h. The mixture was diluted with H_2O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO_4), filtered, and the solvents were removed under

reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to give in order of elution an inseparable mixture of **270** and **271** (1:1 ratio, 5.0 mg; 2.3 mg, 0.012 mmol, 12% **270**; 2.8 mg, 0.012 mmol, 12% **271**) and **269** (7.8 mg, 0.037 mmol, 36%) both as pale yellow oils.

*1-(2,4,5-Trimethoxyphenyl)-2-propanone (272)*¹⁹⁰ and *1-(2,4,5-trimethoxyphenyl)-1-propanone (273)*.¹⁹¹ To a suspension of PdCl₂ (24.2 mg, 0.136 mmol) and CuCl (140 mg, 1.41 mmol) in a solution of MeOH (2 mL) and H₂O (400 μ L) was added **238** (292 mg, 1.40 mmol) dissolved in MeOH (2 mL). The reaction flask was attached to a balloon filled with oxygen and the mixture was stirred at 50 °C for 20 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **273** (62 mg, 0.276 mmol, 20%) and **272** (21.7 mg, 0.097 mmol, 7%) both as pale yellow oils.

*1-(2-Nitrophenyl)-2-pentanone (274)*¹⁹² and *1-(1,1-dimethoxymethyl)-2-nitrobenzene (246)*.

To a suspension of PdCl₂ (12.0 mg, 0.068 mmol) and CuCl (61.4 mg, 0.620 mmol) in a solution of MeOH (1 mL) and H₂O (200 μ L) was added 1-(2-nitrophenyl)-1-pentene (**240**)¹⁶³ (113 mg, 0.589mmol) dissolved in MeOH (1 mL). The reaction flask was attached to a balloon filled with oxygen and the mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **246** (10.7 mg, 0.054mmol, 9%) and **274** (71.6 mg, 0.45 mmol, 58%) as pale yellow oils.

1-(2-Nitrophenyl)-2-pentanone (274), 1-(1,1-dimethoxymethyl)-2-nitrobenzene (246), and 2-nitrobenzaldehyde (247). To a suspension of PdCl₂ (13.0 mg, 0.073 mmol) and H₂O₂ (50%-aqueous, 300 μ L, 6.40 mmol) in a solution of MeOH (1 mL) and H₂O (200 μ L) was added **240** (117 mg, 0.610 mmol) dissolved in MeOH (1 mL). The reaction mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **246** (15.9 mg, 0.081mmol, 13%), **247** (3.2 mg, 0.021mmol, 4%), and **274** (36.7 mg, 0.177mmol, 29%) as pale yellow oils.

1-(2-Nitrophenyl)-2-pentanone (274) and 1-(dimethoxymethyl)-2-nitrobenzene (246). To a suspension of PdCl₂ (6.7 mg, 0.038 mmol) and 2,6-di-tert-butyl-p-cresol (5.4 mg, 0.025 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and H₂O₂ (aqueous-50%, 150 μ L, 3.20 mmol) was added **240** (48.3 mg, 0.252 mmol) dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 °C for 96 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution **246** (5.9 mg, 0.030mmol, 12%) as a pale yellow oil and **274** (26.4 mg, 0.127 mmol, 50%) as a pale yellow oil.

1-(2-Nitrophenyl)-2-pentanone (274). To a suspension of PdCl₂ (6.3 mg, 0.0355 mmol) and 2,6-di-tert-butyl-p-cresol (6.3 mg, 0.0295 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and TBHP (90%-aqueous, 267 μ L, 2.49 mmol) was added **240** (47.3 mg, 0.247mmol) dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure.

The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **274** (32.8 mg, 0.158 mmol, 64%) as a pale yellow oil.

3-Methyl-1-(2-nitrophenyl)-2-butanone (**275**),¹⁹³ **1-(2-nitrophenyl)-3-methoxy-3-methyl-1-butene** (**276**), **4-(2-nitrophenyl)-3-methoxy-2-methyl-1-butene** (**277**). To a suspension of PdCl₂ (11.4 mg, 0.064 mmol) and CuCl (62.1 mg, 0.627 mmol) in a solution of MeOH (1 mL) and H₂O (200 μ L) was added 1-(2-nitrophenyl)-3-methyl-1-butene (**241**)¹⁹⁴ (116 mg, 0.334 mmol) dissolved in MeOH (1 mL). The reaction flask was attached to a balloon filled with oxygen and the mixture was stirred at 50 $^{\circ}$ C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was dissolved in 5 mL of Sat. NaHSO₄ to remove the aldehyde and extracted with 5 mL of ether. The product was then purified by chromatography (hexanes/EtOAc, 95:5) to give in order of elution starting material (51.7 mg, 0.27 mmol, 45%), **277** (17.7 mg, 0.080 mmol, 13%), **276** (15.1 mg, 0.068 mmol, 11%), and **275** (21.5 mg, 0.10 mmol, 17%) as pale yellow oils. Spectral data for **277**: ¹H NMR (600 MHz) δ 7.88 (dd, J =8.4, 1.2 Hz, 1H), 7.50 (td, J =7.8, 1.2 Hz, 1H), 7.38 (dd, J =7.2, 1.2 Hz, 1H), 7.36 (td, J =7.8, 1.8 Hz, 1H), 4.93 (pent, J =1.2 Hz, 1H), 4.89 (pent, J =0.7 Hz, 1H), 3.76 (dd, J =8.5, 4.5 Hz, 1H), 3.15 (dd, J =13.8, 8.4 Hz, 1H), 3.12 (s, 3H), 3.08 (dd, J =13.8, 4.2 Hz, 1H), 1.74 (t, J =1.2 Hz, 3H); ¹³C NMR (150

MHz) δ 149.8, 143.7, 133.8, 133.3, 132.4, 127.3, 124.4, 113.8, 85.1, 56.3, 37.7, 16.8; IR (ATR) 2934, 1578, 1524, 1449, 1348, 1096, 905, 861, 785, 739, 668 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{15}\text{NNaO}_3$ ($\text{M}+\text{Na}^+$) 244.0944; found 244.0944.

3-Methyl-1-(2-nitrophenyl)-2-butanone (275) and 2-nitrobenzaldehyde (247). To a suspension of PdCl_2 (13.1 mg, 0.074 mmol) and H_2O_2 (50%-aqueous, 300 μL , 6.40 mmol) in a solution of MeOH (1 mL) and H_2O (200 μL) was added **241** (117 mg, 0.611 mmol) dissolved in MeOH (1 mL). The reaction mixture was stirred at 50 $^\circ\text{C}$ for 72 h. The mixture was diluted with H_2O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO_4), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution starting material (21.8 mg, 0.11 mmol, 19%), and inseparable mixture of **275** and **247** (2.2:1 ratio, 31.1 mg; 23.3 mg, 0.112 mmol, 18% **275**; 7.7 mg, 0.0512 mmol, 8% **247**) as pale yellow oils.

3-Methyl-1-(2-nitrophenyl)-2-butanone (275), 1-(1,1-dimethoxymethyl)-2-nitrobenzene

(**246**), and **2-nitrobenzaldehyde (247)**. To a suspension of PdCl₂ (5.7 mg, 0.032 mmol) and 2,6-di-tert-butyl-p-cresol (5.8 mg, 0.026 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and H₂O₂ (50%-aqueous, 150 μL, 3.20 mmol) was added **241** (47.1 mg, 0.246 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 96 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give in order of elution starting material (16.9 mg, 0.088 mmol, 36%), **246** (4.5 mg, 0.023mmol, 9%) and an inseparable mixture of **275** and **247** (4.2:1 ratio, 14.9 mg; 12.7 mg, 0.613 mmol, 25% **275**; 2.2 mg, 0.0146 mmol, 6% **247**) as yellow oils.

3-Methyl-1-(2-nitrophenyl)-2-butanone (275). To a suspension of PdCl₂ (5.4 mg, 0.030 mmol) and 2,6-di-tert-butyl-p-cresol (5.6 mg, 0.025 mmol) in a solution of MeOH (500 μL), H₂O (100 μL) and TBHP (90%-aqueous, 267 μL, 2.49 mmol) was added **241** (47.8 mg, 0.250 mmol) dissolved in MeOH (500 μL). The reaction mixture was stirred at 50 °C for 96 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **275** (9.0 mg, 0.043mmol, 17%) as a pale yellow oil.

2-(2-Nitrophenyl)-1-phenyl-ethanone (278).¹⁷² To a suspension of PdCl₂ (14.5 mg, 0.0818 mmol) and CuCl (70.7 mg, 0.714 mmol) in a solution of MeOH (1 mL) and H₂O (200 μL) was added 2-nitro-1-(2-phenylethenyl)benzene (**242**)¹⁹⁵ (152 mg, 0.675 mmol) dissolved in MeOH (1 mL). The reaction flask was attached to a balloon filled with oxygen and the mixture was stirred at 50 °C for 48 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 95:5) to give **278** (23.4 mg, 0.097 mmol, 14%) as a light brown oil.

2-(2-Nitrophenyl)-1-phenyl-ethanone (278). To a suspension of PdCl₂ (14.4 mg, 0.0812 mmol) and 2,6-di-tert-butyl-p-cresol (17.8 mg, 0.0808 mmol) in a solution of MeOH (1 mL), H₂O (200 μL) and H₂O₂ (50%-aqueous, 300 μL, 6.40 mmol) was added **242** (15.0 mg, 0.067 mmol) dissolved in MeOH (1 mL). The reaction mixture was stirred at 50 °C for 68 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced

pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **278** (8.2 mg, 0.034 mmol, 51%) as a light brown oil.

2-(2-Nitrophenyl)-1-phenyl-ethanone (278). To a suspension of PdCl₂ (5.5 mg, 0.031 mmol) and 2,6-di-tert-butyl-p-cresol (5.8 mg, 0.026 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and TBHP (90%-aqueous, 267 μ L, 2.49 mmol) was added **242** (49.1 mg, 0.218 mmol) dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **278** (38.9 mg, 0.161mmol, 74%) as a pale brown oil.

2-(2-Nitrophenyl)-1-phenyl-ethanone (278). To a suspension of PdCl₂ (6.6 mg, 0.037 mmol) in a solution of MeOH (500 μ L), H₂O (100 μ L) and TBHP (90%-aqueous, 267 μ L, 2.49 mmol) was added **242** (50.5 mg, 0.224mmol) dissolved in MeOH (500 μ L). The reaction mixture was stirred at 50 °C for 72 h. The mixture was diluted with H₂O (50 mL) and extracted with

EtOAc (4 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The crude product was purified by chromatography (hexanes/EtOAc, 8:2) to give **278** (14.6 mg, 0.060 mmol, 27%) as a light brown oil.

1-(2-Bromophenyl)-2-propanone (**279**),¹⁹⁶ *1-bromo-2-(1,1-dimethoxymethyl)benzene* (**280**),¹⁹⁷ and *2-bromobenzaldehyde* (**281**), *1-(2-bromophenyl)-1,2-propanedione* (**282**). A solution of **243** (133 mg, 0.672 mmol) in MeOH (1 mL) was added to a suspension of PdCl₂ (12.9 mg, 0.073 mmol) and CuCl (69.8 mg, 0.71 mmol) in a solution of MeOH (1 mL) and H₂O (200 μ L) under an atmosphere of oxygen, as described for **230** in Condition A (50 $^{\circ}$ C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 98:2) gave in order of elution, an inseparable mixture of **280** and **281** (7:1 ratio, 13.1 mg; 11.7 mg, 0.051 mmol, 8% **280**; 1.34 mg, 0.0073 mmol, 1% **281**), **282** (9 mg, 0.040 mmol, $\leq$ 6%), and **279** (64.5 mg, 0.30 mmol, 45%) as pale yellow oils. Spectral data for impure 1-(2-bromophenyl)-1,2-propanedione **282**: ¹H NMR (600 MHz) δ 7.61 (dd, J =7.2, 1.8 Hz, 1H), 7.57 (dd, J =7.2, 2.4 Hz, 1H), 7.45 (td, J =7.2, 1.8 Hz, 1H), 7.41 (td, J =7.2, 1.8 Hz, 1H), 2.59 (s, 3H); ¹³C NMR (100 MHz) 197.9, 193.4, 136.3, 133.6, 133.1, 131.4, 127.7, 121.0, 25.1; IR (ATR) 2970, 1715, 1688, 1587, 1434, 1356, 1172, 1053, 1027, 903, 802, 738 cm⁻¹; HRMS (ESI) calcd for C₉H₈BrO₂ (M+H⁺) 226.9708; found 226.9702.

1-Phenyl-2-propanone (283)¹⁹⁸ and **1-phenyl-1-propanone (284)**. A solution of 1-propen-1-yl-benzene (**244**)¹⁹⁹ (95.3 mg, 0.806 mmol) in MeOH (1 mL) was added to a suspension of PdCl₂ (14.3 mg, 0.081 mmol) and CuCl (79.8 mg, 0.81 mmol) in a solution of MeOH (1 mL) and H₂O (200 μL) under an atmosphere of oxygen, as described for **230** in Condition A (50 °C, 20 h). Extraction and purification by chromatography (hexanes/EtOAc, 9:1) gave in order of elution, **284** (16.0 mg, 0.099 mmol, 12%) and **283** (44.2 mg, 0.27 mmol, 34%) as colorless oils.

References and Notes

- ¹ Karlsson, J. O.; Nguyen, N. V.; Foland, L. D.; Moore, H. W. *J. Am. Chem. Soc.* **1985**, *107*, 3392.
- ² Moore, H. W.; Decker, O. H. W. *Chem. Rev.* **1986**, *86*, 821.
- ³ Moore, H. W. *Science* **1977**, *197*, 521.
- ⁴ Moore, H. W.; Czernick, R. *Med. Res. Rev.* **1981**, *1*, 249.
- ⁵ Kikuchi, H.; Tsukitani, Y.; Iguchi, K.; Yamada, Y. *Tetrahedron Lett.* **1982**, *23*, 25.
- ⁶ Iguchi, K.; Kaneta, S.; Mori, K.; Yamada, Y.; Honda, A.; Mori, Y. *Tetrahedron Lett.* **1985**, *26*, 5787.
- ⁷ Baker, B. J.; Okuda, R. K.; Yu, P. T. K.; Scheuer, P. J. *J. Am. Chem. Soc.* **1985**, *107*, 2976.
- ⁸ Reed, M.; Perri, S. T.; Pollart, D.; Foland, L. D.; Moore, H. W. *J. Org. Chem.* **1988**, *53*, 2411.
- ⁹ Liebeskind, L. S.; Fengl, R. W.; Wirtz, K. R.; Shawe, T. T. *J. Org. Chem.* **1988**, *53*, 2482.
- ¹⁰ Perri, S. T.; Foland, L. D.; Decker, O. H. W.; Moore, H. W. *J. Org. Chem.* **1986**, *51*, 3067.
- ¹¹ Decker, O. H. W.; Moore, H. W. *J. Org. Chem.* **1987**, *52*, 1174.
- ¹² Foland, L. D.; Karlsson, J. O.; Perri, S. T.; Schwabe, R.; Xu, S. L.; Patil, S.; Moore, H. W. *J. Am. Chem. Soc.* **1989**, *111*, 975.
- ¹³ Hashimoto, T. *Yakugaku Azsshi* **1967**, *87*, 535.
- ¹⁴ Back, H.; Alt, H. *Angew. Chem., Int. Ed. Engl.* **1967**, *6*, 944.
- ¹⁵ Foland, L. D.; Decker, O. H. W.; Moore, H. W. *J. Am. Chem. Soc.* **1989**, *111*, 989.
- ¹⁶ Thomson, R. H. *Naturally Occurring Quinones III*; Chapman and Hall: London, **1987**.
- ¹⁷ Thomson, R. H. *Naturally Occurring Quinones*; Academic: New York, **1976**.

- ¹⁸ Xin-Sheng, Y.; Ebizuka, Y.; Noguchi, H.; Kiuchi, F.; Seto, H.; Sankawa, U. *Tetrahedron Lett.* **1984**, 5541.
- ¹⁹ Zhon Yao Zhi, Chinese Academy of Medicinal Sciences, Ed.; People's Hygenic Publisher: **1979**, 1, 569.
- ²⁰ Kruchi, F.; Shibuya, M.; Sankawa, U. *Chem. Pharm. Bull.* **1982**, 30, 754.
- ²¹ Kruchi, F.; Shibuya, M.; Sankawa, U. *Ibid.* **1982**, 30, 2279.
- ²² Hayashi, M.; Unemoto, T.; Minami-Kakinuma, S.; Tanaka, H.; Omura, S. *J. Antibiot.* **1982**, 35, 1078.
- ²³ Watanabe, S.; Shimizu, H. *J. Antibiot.* **1985**, 38, 1447.
- ²⁴ Chang, C. J.; Floss, H. G.; Soong, P.; Gang, C. T. *J. Antibiot.* **1975**, 28, 156.
- ²⁵ Liebeskind, L. S.; Jewell, C. F.; Iyer, S. *Ibid.* **1986**, 51, 3066.
- ²⁶ Bernady, K. F.; Floyd, M. B.; Poletto, J. F.; Weiss, M. J. *J. Org. Chem.* **1979**, 44, 1438.
- ²⁷ Naruta, Y.; Uno, H.; Maruyama, K. *Chem. Lett.* **1982**, 609.
- ²⁸ Ellestad, G. Z.; Kuntsmann, M. P.; Whaley, H. A.; Patterson, E. L. *J. Am. Chem. Soc.* **1968**, 90, 1325.
- ²⁹ Ichihara, A.; Ubukata, M.; Oikawa, H.; Murakami, K.; Sakamura, S. *Tetrahedron Lett.* **1980**, 4469.
- ³⁰ Kraus, G. A.; Roth, B. *J. Org. Chem.* **1978**, 43, 4923.
- ³¹ Enhsen, A.; Karabelas, K.; Heerding, J. M.; Moore, H. W. *J. Org. Chem.* **1990**, 55, 1177.
- ³² Reed, M. W.; Perri, S. T.; Pollart, D. J.; Foland, L. D.; Moore, H. W. *J. Org. Chem.* **1988**, 53, 2477.
- ³³ Beyerman, H. C.; Bontekoe, J. S. *Proc. Chem. Soc.* **1961**, 249.

- ³⁴ Read, G.; Westlake, D. W. S.; Vining, L. C. *Can. J. Biochem.* **1969**, *47*, 1071.
- ³⁵ Yamamoto, H.; Moriyama, K.; Jinnouchi, H.; Yagishita, K. *J. Antibiot.* **1980**, *33*, 320.
- ³⁶ Moore, H. W. *J. Org. Chem.* **1967**, *32*, 1996.
- ³⁷ Rashid, A.; Read, G. *J. Chem. Soc.* **1967**, 1323.
- ³⁸ Sheehan, J. C.; Lo, Y. S. *J. Med. Chem.* **1974**, *17*, 371.
- ³⁹ Oppolzer, W.; Zutterman, F.; Battig, K. *Helv. Chim. Acta* **1983**, *66*, 522.
- ⁴⁰ E. Garcia, G.; Mendoza, V.; A. Guzman B. *J. Nat. Prod.* **1987**, *50*, 1055.
- ⁴¹ Clark, G. R.; Thiensathit, S. *Tetrahedron Lett.* **1985**, 2503.
- ⁴² Liebeskind, L. S.; Chidambaram, R. *J. Am. Chem. Soc.* **1987**, *109*, 5025.
- ⁴³ Liebeskind, L. S.; Mitchell, D.; Foster, B. S. *J. Am. Chem. Soc.* **1987**, *109*, 7908.
- ⁴⁴ Gul, W.; Hamann, M. T. *Life Sciences.* **2005**, *78*, 442.
- ⁴⁵ Kikuchi, H.; Tsukitani, Y.; Iguchi, K.; Yamada, Y. *Tetrahedron Lett.* **1982**, 5171.
- ⁴⁶ Kobayashi, M.; Yasuzawa, T.; Yoshihara, M.; Akutsu, H.; Kyogoku, Y.; Kitagawa, I. *Tetrahedron Lett.* **1982**, 5331.
- ⁴⁷ Kobayashi, M.; Yasuzawa, T.; Yoshihara, M.; Son, B. W.; Kyogoku, Y.; Kitagawa, I. *Chem. Pharm. Bull.* **1983**, *11*, 1440.
- ⁴⁸ Nagaoka, H.; Iguchi, K.; Miyakoshi, T.; Yamada, N.; Yamada, Y. *Ibid.* **1986**, 223.
- ⁴⁹ Iguchi, K.; Kaneta, S.; Mori, K.; Yamada, Y.; Honda, A.; Mori, Y. *J. Chem. Soc., Chem. Commun.* **1986**, 981.
- ⁵⁰ Nagaoka, H.; Miyaoka, H.; Miyakoshi, Y.; Yamada, Y. *Ibid.* **1986**, *108*, 5019.
- ⁵¹ Fukushima, M.; Kato, T.; Yamada, Y.; Kitagawa, I.; kurozumi, S.; Scheuer, P. J. *Proc. Am. Assoc. Cancer Res.* **1985**, *26*, 249.

- ⁵² Katsin, N.; R. Ikan, R. *Synth. Commun.* **1977**, 7, 185.
- ⁵³ Pohmakotr, M.; Chancharunee, S. *Tetrahedron Lett.* **1984**, 4141.
- ⁵⁴ Block, E.; Wall, A.; Zubieta, J. *J. Am. Chem. Soc.* **1985**, 107, 1783.
- ⁵⁵ Verlaak, J. M. J.; Klunder, A. J. H.; Zwanenburg, B. *Tetrahedron Lett.* **1982**, 5436.
- ⁵⁶ Boeckmann, R. J.; Naegely, P. C.; Arthur, S. D. *J. Org. Chem.* **1980**, 45, 752.
- ⁵⁷ Corey, E. J.; Mehrotra, M. M. *J. Am. Chem. Soc.* **1984**, 106, 3384.
- ⁵⁸ Sasai, H.; Shibasaki, M. *Tetrahedron Lett.* **1987**, 333.
- ⁵⁹ Yamanoto, Y.; Ohno, M.; Eguchi, S. *Tetrahedron Lett.* **1995**, 36, 5539.
- ⁶⁰ Yamamoto, Y.; Noda, M.; Ohno, M.; Eguchi, S. *J. Org. Chem.* **1997**, 62, 1292.
- ⁶¹ Mitchell, D.; Liebeskind, L. S. *Ibid.* **1990**, 112, 291.
- ⁶² Liebeskind, L. S.; Foster, B. S. *J. Am. Chem. Soc.* **1990**, 112, 8612.
- ⁶³ Nishikawa, T.; Shibuya, S.; Hosokawa, S.; Isobe, M. *Synlett.* **1994**, 485.
- ⁶⁴ Kiang, A. K.; Lee, H. H.; Sim, K. Y. *Proc. Chem. Soc. London* **1961**, 455.
- ⁶⁵ Liu, S. Y.; Hisada, S.; Inagaki, I. *Phytochemistry* **1973**, 12, 233.
- ⁶⁶ Corner, E. J. H. *Wayside Trees of Malaya*, 3rd ed. The Malayan Nature Society, Kuala Lumpur, Malaysia, **1988**, 1, 371.
- ⁶⁷ Ichino, K.; Tanaka, H.; Ito, K.; Tanaka, T.; Mizuno, M. *J. Nat. Prod.* **1988**, 51, 915.
- ⁶⁸ Komae, H.; Hayashi, N.; *Phytochemistry* **1972**, 11, 853.
- ⁶⁹ Lee, H. H. *Tetrahedron Lett.* **1968**, 40, 4243.
- ⁷⁰ Leong, Y. -W.; Harrison, L. J.; Bennett, G. J.; Kadir, A. A.; Connolly, J. D. *Phytochemistry* **1998**, 47, 891.
- ⁷¹ Kiang, A. K.; Lee, H. H.; Sim, K. Y. *J. Chem. Soc.* **1962**, 438.

- ⁷² Chia, Y. C.; Wu, J.-B.; Wu, Y.-C. *Tetrahedron Lett.* **2000**, *41*, 2199.
- ⁷³ Wu, J. B.; Cheng, Y. D.; Kuo, S. C.; Wu, T. S.; Iitaka, Y.; Ebizuka, Y.; Sankawa, U. *Chem. Pharm. Bull.* **1994**, *42*, 2202.
- ⁷⁴ Aoyama, Y.; Konoike, T.; Kanda, A.; Naya, N.; Nakajima, M. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1695.
- ⁷⁵ Lee, S. M.; Baek, S.W.; Lee, C.H.; Lee, H. B.; Kho, Y. H. *Nat. Prod. Sci.* **2002**, *8*, 100.
- ⁷⁶ Hwang, E. I.; Lee, Y. M.; Lee, S. M.; Yeo, W. H.; Moon, J. S.; Te, H.; Park, K. D. D.; Kim, S. U. *Planta Med.* **2007**, *73*, 679.
- ⁷⁷ Wang, Z.; Walter, M.; Selwood, T.; Rubin, H.; Schechter, N. M. *Biol. Chem.* **1998**, *379*, 167.
- ⁷⁸ Ihara, M.; Urata, H.; Kinoshita, A.; Suzumiya, J.; Sasaguri, M.; Kikuchi, M.; Ideishi, M.; Arakawa, K. *Hypertension* **1999**, *33*, 1399.
- ⁷⁹ Takai, S.; Yuda, A.; Jin, D.; Nishimoto, M.; Sakaguchi, M.; Sasaki, S.; Miyazaki, M. *FEBS Lett.* **2000**, *467*, 141.
- ⁸⁰ Oh, H. M.; Choi, S. K.; Lee, J. M.; Lee, S.-K.; Kim, H.-Y.; Han, D. C.; Kim, H. M.; Son, K. H.; Kwon, B. M. *Bioorg. Med. Chem.* **2005**, *13*, 6182.
- ⁸¹ Wang, S.-Y.; Lan, X.-Y.; Xiao, J.-H.; Yang, J.-C.; Kao, Y.-T.; Chang, S.-T. *Phytother. Res.* **2008**, *22*, 213.
- ⁸² Kim, G. -O.; Jin, Y. J.; Ko, R. -K.; Choi, H. -M.; Kang, M. -C.; Han, J. -H.; KO, K. -H. PCT Int. Appl. WO 20,110,556,037 (CAN 154: 557389), **2011**.
- ⁸³ Birch, A. J. *J. Chem. Soc.* **1951**, 3026.
- ⁸⁴ Kiang, A. K.; Lee, H. H.; Sim, K.Y. *J. Chem. Soc.* **1962**, 4338.

- ⁸⁵ Lee, H. H.; Tan, C. H. *J. Chem. Soc.* **1967**, 1583.
- ⁸⁶ Bennett, G. L.; Lee, H. H. *J. Chem. Soc. Perkin Trans. I* **1986**, 633.
- ⁸⁷ Bose, G.; Langer, P. *Synlett.* **2005**, 1021.
- ⁸⁸ Birch, A. J.; Elliot, P. *Austral. J. Chem.* **1956**, 9, 95.
- ⁸⁹ Lee, H. H.; Que, Y. T.; Ng, S. *J. Chem. Soc. Perkin Trans. I* **1985**, 453.
- ⁹⁰ Rao, K. V.; Seshadri, T. R. *Proc. Indian Acad. Sci., Sect. A* **1948**, 27, 375.
- ⁹¹ Corey, E. J.; Fuchs, P. L. *Tetrahedron Lett.* **1972**, 13, 3769.
- ⁹² Clemo, N. G.; Gedge, D. R. *J. Chem. Soc., Perkin Trans. I* **1981**, 1448.
- ⁹³ Michel, P.; Genner, D.; Rassat, A. *Tetrahedron Lett.* **1999**, 40, 8575.
- ⁹⁴ Tang, L. N.; Pang, Y. L.; Yan, Q.; Shi, L. Q.; Huang, J. H.; Du, Y. F.; Zhao, K. *J. Org. Chem.* **2011**, 76, 2744.
- ⁹⁵ Chittimalla, S. K.; Chang, T.-C.; Liu, T.-C.; Hsieh, H. P.; Liao, C.-C. *Tetrahedron* **2008**, 64, 2586.
- ⁹⁶ Kim, Y.-W.; Hackett, J. C.; Brueggemeier, R. W. *J. Med. Chem.* **2004**, 47, 4032.
- ⁹⁷ Xu, Q. L.; Gao, H. Y.; Yousufuddin, M.; Ess, D. H.; Kürti, L. *J. Am. Chem. Soc.* **2013**, 135, 14048.
- ⁹⁸ Schlummer, B.; Scholz, U. In *Modern Arylation Methods*. Ackermann, L. Ed. Wiley-VCH: Weinheim, **2009**, 69.
- ⁹⁹ Kessar, S. V. In *Comprehensive Organic Synthesis*. Trost, B. M.; Fleming, I.; Semmelhack, M. F. Eds. Pergamon Press: New York, **1991**, 4, Chapter 2.3.
- ¹⁰⁰ Semmelhack, M. F.; Bargar, T. *J. Org. Chem.* **1977**, 42, 1481.
- ¹⁰¹ Beugelmans, R.; Ginsburg, H. *Heterocycles* **1985**, 23, 1197.

- ¹⁰² Finet, J. P. *Chem. Rev.* **1989**, 89, 1487.
- ¹⁰³ Jenkins, S. S. *J. Am. Chem. Soc.* **1933**, 55, 2896.
- ¹⁰⁴ Yadav, D. K.; Gautam, A. K.; Kureel, J.; Srivastava, K.; Sahai, M.; Singh, D.; Chattopadhyay, N.; Maurya, R. *Bioorg. Med. Chem. Lett.* **2011**, 21, 677.
- ¹⁰⁵ Landers, B.; Berini, C.; Wang, C.; Navarro, O. *J. Org. Chem.* **2011**, 76, 1390.
- ¹⁰⁶ Fox, J. M.; Huang, X.; Chieffi, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2000**, 122, 1360.
- ¹⁰⁷ Miao, T.; Wang, G.-W. *Chem. Commun.* **2011**, 47, 9501.
- ¹⁰⁸ Kuwajima, I.; Urabe, H. *J. Am. Chem. Soc.* **1982**, 104, 6831.
- ¹⁰⁹ Palucki, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1997**, 119, 11108.
- ¹¹⁰ Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1997**, 119, 12382.
- ¹¹¹ Terao, Y.; Satoh, T.; Miura, M.; Nomura, M. *Bull. Chem. Soc. Jpn.* **1999**, 72, 2345.
- ¹¹² Viciu, M. S.; Germaneau, R. F.; Nolan, S. P. *Org. Lett.* **2002**, 4, 4053.
- ¹¹³ Viciu, M. S.; Germaneau, R. F.; Navarro-Fernandez, O.; Stevens, E. D.; Nolan, S. P. *Organometallics* **2002**, 21, 5470.
- ¹¹⁴ Singh, R.; Nolan, S. P. *J. Organomet. Chem.* **2005**, 690, 5832.
- ¹¹⁵ Navarro, O.; Marion, N.; Scott, N. M.; González, J.; Amoroso, D.; Bell, A.; Nolan, S. P. *Tetrahedron* **2005**, 61, 9716.
- ¹¹⁶ Schnyder, A.; Indolese, A. F.; Studer, M.; Blaser, H.-U. *Angew. Chem., Int. Ed.* **2002**, 41, 3668.
- ¹¹⁷ Lavallo, V.; Canac, Y.; Prasang, C.; Donnadiou, B.; Bertrand, G. *Angew. Chem., Int. Ed.* **2005**, 44, 5705.
- ¹¹⁸ Adjabeng, G.; Brenstrum, T.; Frampton, C. S.; Robertson, A. J.; Hillhouse, J.; McNulty,

J.; Capretta, A. *J. Org. Chem.* **2004**, *69*, 5082.

¹¹⁹ Churruca, F.; SanMartin, R.; Tellitu, I.; Dominguez, E. *Tetrahedron Lett.* **2006**, *47*, 3233.

¹²⁰ Schuster, E. M.; Botoshansky, M.; Gandelman, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 4555.

¹²¹ Khartulyari, A. S.; Maier, M. E. *Eur. J. Org. Chem.* **2007**, 317.

¹²² Ahman, J.; Wolfe, J. P.; Troutman, M. V.; Palucki, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 1918.

¹²³ Chen, G.; Kwong, F. Y.; Chan, H. O.; Yu, W.-Y.; Chan, A. S. C. *Chem. Commun.* **2006**, 1413.

¹²⁴ Carril, M.; SanMartin, R.; Churruca, F.; Tellitu, I.; Dominguez, E. *Org. Lett.* **2005**, *7*, 4787.

¹²⁵ Churruca, F.; SanMartin, R.; Tellitu, I.; Dominguez, E. *Org. Lett.* **2002**, *4*, 1591.

¹²⁶ Churruca, F.; SanMartin, R.; Carril, M.; Tellitu, I.; Dominguez, E. *Tetrahedron* **2004**, *60*, 2393.

¹²⁷ Gupta, M.; Nair, V. *Tetrahedron Lett.* **2005**, *46*, 1165.

¹²⁸ Nicolaou, K. C.; Xu, H. *Chem. Commun.* **2006**, 600.

¹²⁹ Jira, R. In *Applied Homogeneous Catalysis with Organometallic Compounds*, 2nd ed. (Eds.: B. Cornils, W. A. Herrmann), Wiley-VCH, Weinheim, **2002**, 38.

¹³⁰ Smidt, J.; Hafner, W.; Jira, R.; Sieber, R.; Sedlmeier, S.; Sabel, A. *Angew. Chem.* **1962**, *74*, 93.

¹³¹ Phillips F. C. *Am. Chem. J.* **1894**, *16*, 255.

¹³² Fahey, D. R.; Zeuch, E. A. *J. Org. Chem.* **1974**, *39*, 3276.

- ¹³³ McQuillin, F. J.; Parker, D. G. *J. Chem. Soc. Perkin Trans. 1* **1974**, 809.
- ¹³⁴ Takacs, J. M.; Jiang, X. -T. *Curr. Org. Chem.* **2003**, 7, 369.
- ¹³⁵ Henry, P. M. *Handb. Organopalladium Chem. Org. Synth.* **2002**, 2, 2119.
- ¹³⁶ Backvall, J. E.; Heumann, A. *J. Am. Chem. Soc.* **1986**, 108, 7107.
- ¹³⁷ Li, J. J. *Name Reactions: A Collection of Detailed Mechanisms and Synthetic Applications*. Springer-Verlag Berlin Heidelberg, **2009**, 564.
- ¹³⁸ Keith, J. A.; Henry, P. M. *Angew. Chem. Int. Ed.* **2009**, 48, 9038.
- ¹³⁹ Tsuji, J. *Synthesis* **1984**, 5, 369.
- ¹⁴⁰ Urbance, J.; Hrusovsky, M.; Laton, V. *Zb. Pr. Chemickotechnol. Fak. SVST* **1971**, 161.
- ¹⁴¹ Tsuji, J.; Nagashima, H.; Nemoto, H. *Org. Synth.* **1984**, 62, 9.
- ¹⁴² Huttel, R.; McNiff, M. *Chem. Ber.* **1973**, 106, 1789.
- ¹⁴³ Moiseev, I. I.; Vargaftik, M. N.; Syrkin, Y. K. *Dokl. Akad. Nauk SSS* **1960**, 133, 377.
- ¹⁴⁴ Tsuji, J.; Nagashima, H.; Hori, K. *Chem. Lett.* **1980**, 257.
- ¹⁴⁵ Mimoun, H.; Charpentier, R.; Mitschler, A.; Fischer, J.; Weiss, R. *J. Am. Chem. Soc.* **1980**, 102, 1047.
- ¹⁴⁶ Roussel, M.; Mimoun, H. *J. Org. Chem.* **1980**, 45, 5387.
- ¹⁴⁷ Tsuji, J.; Shimizu, I.; Yamanoto, K. *Tetrahedron Lett.* **1976**, 2975.
- ¹⁴⁸ Liao, X.; Zhou, H.; Yu, J.; Cook, J. M. *J. Org. Chem.* **2006**, 71, 8884.
- ¹⁴⁹ Kavala, M.; Mathia, F.; Kozísek, J.; Szolcsanyi, P. *J. Nat. Prod.* **2011**, 74, 803.
- ¹⁵⁰ Mehta, G.; Yaragorla, S. *Tetrahedron Lett.* **2013**, 54, 549.
- ¹⁵¹ Ferdjani, O. S.; Kar, A.; Chavan, S. P.; Horeau, M.; Viault, G.; Pouessel, J.; Guillory, X.; Blot, V.; Tessier, A.; Planchat, A.; Jacquemin, D.; Dubreuil, D.; Pipelier, M. *Eur. J. Org. Chem.*

2013, 7083.

- ¹⁵² Hansford, K. A.; Zanzarova, V.; Dörr, A.; Lubell, W. D. *J. Comb. Chem.* **2004**, 6, 893.
- ¹⁵³ Saroukou, M.; Skalski, T.; Skene W.G.; Lubell, W. D. *Tetrahedron*, **2014**, 70, 450.
- ¹⁵⁴ Sims E. A.; DeForest, C. A.; Anseth, K. S. *Tetrahedron Lett.* **2011**, 52, 1871.
- ¹⁵⁵ Michel, B. W.; Camelio, A. M.; Cornell, C. N.; Sigman, M. S.; *J. Am. Chem. Soc.* **2009**, 131, 6076.
- ¹⁵⁶ McCombs, J. R.; Michel, B. W.; Sigman, M. S. *J. Org. Chem.* **2011**, 76, 3609.
- ¹⁵⁷ Muzart, J. *Tetrahedron* **2007**, 63, 7505.
- ¹⁵⁸ Schnapperelle, I.; Hummel, W.; Gröger, H. *Chem. Eur. J.* **2012**, 18, 1073.
- ¹⁵⁹ Mitsudome, T.; Mitzumo, K.; Mizugaki, T.; Jitsukawa, K.; Kaneda, K. *Angew. Chem. Int. Ed.* **2010**, 49, 1238.
- ¹⁶⁰ Ross, J. P.; Couture, P.; Warkentin, J. *Can. J. Chem.* **1997**, 75, 1331.
- ¹⁶¹ Cornell, C. N. ; Sigman, M. S. *J. Am. Chem. Soc.* **2005**, 127, 2796.
- ¹⁶² Keinan, E.; Seth, K. K.; Lamed, R. *J. Am. Chem. Soc.* **1986**, 108, 3474.
- ¹⁶³ Okuro, K.; Gurnham, J.; Alper, H. *J. Org. Chem.* **2011**, 76, 4715.
- ¹⁶⁴ Gruber, M.; Chouzier, S.; Koehler, K.; Djakovitch, L. *Applied Catalysis A*: **2004**, 265, 161.
- ¹⁶⁵ Kulkarni, M. G.; Shaikh, Y. B.; Borhade, A. S.; Chavan, S. W.; Dhondge, A. P.; Gaikwad, D. D.; Desai, M. P.; Bihade, D. R.; Dhattrak, N. R. *Tetrahedron Lett.* **2013**, 54, 2293.
- ¹⁶⁶ Zhang, G.; Xie, X.; Wang, Y.; Wen, X.; Zhao, Y.; Ding, C. *Org. Biomol. Chem.* **2013**, 11, 2947.
- ¹⁶⁷ Thimmaiah, M.; Zhang, X.; Fang, S. *Tetrahedron Lett.* **2008**, 49, 5605.

- ¹⁶⁸ Labadie, J. W.; Stille, J. K. *J. Am. Chem. Soc.* **1983**, *105*, 6129.
- ¹⁶⁹ Wan, P.; Davies, M. J.; Teo, M.-A. *J. Org. Chem.* **1989**, *54*, 1354.
- ¹⁷⁰ Mariampillai, B.; Herse, C.; Lautens, M. *Org. Lett.* **2005**, *7*, 4745.
- ¹⁷¹ Padwa, A.; Rieker, W. F.; Rosenthal, R. J. *J. Am. Chem. Soc.* **1985**, *107*, 1710.
- ¹⁷² Coffman, K. C.; Palazzo, T. A.; Hartley, T. P.; Fettingner, J. C.; Tantillo, D. J.; Kurth, M. J. *Org. Lett.* **2013**, *15*, 2062.
- ¹⁷³ Gopinath, R.; Haque, S. J.; Patel, B. K. *J. Org. Chem.* **2002**, *67*, 5842.
- ¹⁷⁴ Aridoss, G.; Laali, K. K. *J. Org. Chem.* **2011**, *76*, 8088.
- ¹⁷⁵ Bovicelli, P.; Sanetti, A.; Lupattelli, P. *Tetrahedron* **1996**, *52*, 10969.
- ¹⁷⁶ Rodriguez, J. G.; Lafuente, A.; Martin-Villamil, R.; Martinez-Alcazar, M. P. *J. Phys. Org. Chem.* **2001**, *14*, 859.
- ¹⁷⁷ Aridoss, G.; Laali, K. K. *J. Org. Chem.* **2011**, *76*, 8088.
- ¹⁷⁸ Meurer, E. C.; Gozzo, F. C.; Augusti, R.; Eberlin, M. N. *Eur. J. Mass. Spectrom.* **2003**, *9*, 295.
- ¹⁷⁹ Hering, T.; Hari, D. P.; König, B. *J. Org. Chem.* **2012**, *77*, 10347.
- ¹⁸⁰ Ito, S.; Hayashi, A.; Komai, H.; Yamaguchi, H.; Kubota, Y.; Asami, M. *Tetrahedron* **2011**, *67*, 2081.
- ¹⁸¹ Bovicelli, P.; Sanetti, A. *Tetrahedron* **1996**, *52*, 10969.
- ¹⁸² Broos, R. *Bulletin des Societes Chimiques Belges* **1986**, *95*, 135.
- ¹⁸³ Hering, T.; Hari, D. P.; König, B. *J. Org. Chem.* **2012**, *77*, 10347.
- ¹⁸⁴ Ross, J. P.; Couture, P.; Warkentin, J. *Can. J. Chem.* **1997**, *75*, 131.
- ¹⁸⁵ Gallardo, I.; Guirado, G.; Marquet, J. *J. Org. Chem.* **2003**, *68*, 7334.

- ¹⁸⁶ Pamukcu, R.; Paizza, G. A. **2002**, US 6410584 [CAN137:47109].
- ¹⁸⁷ Conner, S. E.; Mantio, N. B.; Zhu, G.; Herr, R. J. *PCT Int. Appl.* **2005**, 2005066136. [CAN 143:153366].
- ¹⁸⁸ Molinaro, C.; Mowat, J.; Gosselin, F.; O'Shea, P.; Marcoux, J. -F.; Angelaud, R.; Davies, I. W. *J. Org. Chem.* **2007**, 72, 1856.
- ¹⁸⁹ Fürstner, A.; Jumbam, D. N.; Seidel, G. *Chem. Ber.* **1994**, 127, 1125.
- ¹⁹⁰ Neudorffer, A.; Mueller, M.; Martinez, C. -M.; Mechan, A.; McCann, U.; Ricaurte, G. A.; LARGERON, M. *Chem. Res. Toxicol.* **2011**, 24, 968.
- ¹⁹¹ Joshi, B. P.; Sharma, A.; Sinha, A. K. *Can. J. Chem.* **2005**, 83, 1826.
- ¹⁹² Raucher, S.; Koolpe, G. A. *J. Org. Chem.* **1983**, 48, 2066.
- ¹⁹³ Eynard, T.; Franc, C. *PCT Int. Appl.* **2006**, WO 2006120031 [CAN145:505444].
- ¹⁹⁴ Pirrung, M. C.; Liu, Y.; Deng, L.; Halstead, D. K.; Li, Z.; May, J. F.; Wedel, M.; Austin, D. A.; Webster, N. J. G. *J. Am. Chem. Soc.* **2005**, 127, 4609.
- ¹⁹⁵ Sundberg, R. J. *J. Org. Chem.* **1965**, 30, 3604.
- ¹⁹⁶ Viswanathan, R.; Prabhakaran, E. N.; Plotkin, M. A.; Johnston, J. N. *J. Am. Chem. Soc.* **2003**, 125, 163.
- ¹⁹⁷ Martin, C.; Mailliet, P.; Maddaluno, J. *J. Org. Chem.* **2001**, 66, 3797.
- ¹⁹⁸ Mitsudome, T.; Umetami, T.; Nosaka, N.; Mori, K.; Mizugaki, T.; Ebitani, K.; Kaneda, K. *Angew. Chem. Int. Ed.* **2006**, 45, 481.
- ¹⁹⁹ Thimmaiah, M.; Zhang, X.; Fang, S. *Tetrahedron Lett.* **2008**, 49, 5605.

Appendix

^1H and ^{13}C NMR Spectra

- ^1H and ^{13}C NMR for Chapter 1: Ring-expansion of 4-ethynyl-4-hydroxy-2,3-substituted-2-cyclobuten-1-ones 149
- ^1H and ^{13}C NMR for Chapter 2: Total synthesis of methyl linderone 201
- ^1H and ^{13}C NMR for Chapter 3: Wacker-type oxidation of aryl-substituted alkenes 208

Figure 17: ^1H NMR of 2,3-bis(1-methylethoxy)-5-trimethylsilyl-2,5-cyclohexadiene-1,4-dione (**70**)

Figure 18: ¹³C NMR of 2,3-bis(1-methylethoxy)-5-trimethylsilyl-2,5-cyclohexadiene-1,4-dione (**70**)

Figure 19: ¹H NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**71**)

Figure 20: ¹³C NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**71**)

cyclopentene-1,3-dione (**72**)

Figure 22: ^{13}C NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**72**)

Figure 23: ¹H NMR of 2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**74**)

Figure 24: ^{13}C NMR of 2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (74)

Figure 25: ¹H NMR of 2-(iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**75**)

Figure 26: ¹³C NMR of 2-(iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-

1,3-dione (**75**)

Figure 27: ¹H NMR of 2-(bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**76**)

Figure 28: ¹³C NMR of 2-(bromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**76**)

Figure 29: ¹H NMR of 5-chloro-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (77)

Figure 30: ^{13}C NMR of 5-chloro-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**77**)

Figure 31: ¹H NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-iodoethynyl)-2-cyclobutene-1-one (**78**)

Figure 32: ¹³C NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-iodoethynyl)-2-cyclobutene-1-one (**78**)

Figure 33: ¹H NMR of 5-iodo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**79**)

Figure 34: ^{13}C NMR of 5-iodo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**79**)

1,3-dione (**80**)

Figure 36: ¹³C NMR of 2-(diiodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-

1,3-dione (**80**)

Figure 37: ^1H NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-bromoethynyl)-2-cyclobutene-1-one (**81**)

Figure 38: ^{13}C NMR of 4-hydroxy-2,3-bis(1-methylethoxy)-4-(2-bromoethynyl)-2-cyclobutene-1-one (**81**)

Figure 39: ^1H NMR of 5-bromo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**82**)

Figure 40: ^{13}C NMR of 5-bromo-2,3-bis(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**82**)

Figure 41: ^1H NMR of 2-(1-bromo-1-iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**83**)

Figure 42: ^{13}C NMR of 2-(1-bromo-1-iodomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**83**)

Figure 43: ^1H NMR of 2-(dibromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-1,3-dione (**84**)

Figure 44: ¹³C NMR of 2-(dibromomethylene)-4,5-bis(1-methylethoxy)-4-cyclopentene-

1,3-dione (**84**)

Figure 45: ¹H NMR of 4-hydroxy-2-methyl-3-(1-methylethoxy)-4-(2-trimethylsilylethynyl)-2-cyclobutene-1-one (**89**)

Figure 46: ¹³C NMR of 4-hydroxy-2-methyl-3-(1-methylethoxy)-4-(2-trimethylsilylethynyl)-2-cyclobutene-1-one (**89**)

Figure 47: ¹H NMR of 2-trimethylsilylmethylene-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**90**) and 5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**91**)

Figure 48: ¹³C NMR of 2-trimethylsilylmethylene-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**90**) and 5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**91**)

Figure 50: ¹³C NMR of 2-(1-iodo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**92**)

cyclohexadiene-1,4-dione (**93**)

cyclohexadiene-1,4-dione (**93**)

185

Figure 54: ¹³C NMR of 2-(1-bromo-1-trimethylsilylmethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**94**)

Figure 55: ¹H NMR of 6-bromo-5-trimethylsilyl-3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (**95**)

cyclohexadiene-1,4-dione (**95**)

Figure 57: ¹H NMR of 3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (97)

Figure 58: ¹³C NMR of 3-methyl-2-(1-methylethoxy)-2,5-cyclohexadiene-1,4-dione (97)

Figure 59: ^1H NMR of 2-(1-iodomethylene)-4-methyl-5-(1-methylethoxy)-4-cyclopentene-1,3-dione (**98**)

Figure 60: ¹³C NMR of 2-(1-iodomethylene)-4-methyl-5-(1-methylethoxy)-4-cyclopentene-1,3-dione (**98**)

Figure 61: ¹H NMR of 2-(1-bromomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**99**)

Figure 62: ¹³C NMR of 2-(1-bromomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**99**)

Figure 63: ^1H NMR of 2-(diiodomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**101**)

Figure 64: ^{13}C NMR of 2-(diiodomethylene)-4-(1-methylethoxy)-5-methyl-4-cyclopentene-1,3-dione (**101**)

Figure 65: ¹H NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-iodoethanoate (**104**)

Figure 66: ^{13}C NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-iodoethanoate (**104**)

Figure 67: ¹H NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-bromoethanoate (**105**)

Figure 68: ^{13}C NMR of ethyl 2-(3,4-dimethoxy-2,5-dioxo-3-cyclopenten-1-ylene)-2-bromoethanoate (**105**)

Figure 69: ^1H NMR of 2,3-dimethoxy-4-hydroxy-4-[(E)-4-phenyl-3-buten-1-yn-1-yl]-2-cyclobuten-1-one (**137**)

Figure 70: ¹³C NMR of 2,3-dimethoxy-4-hydroxy-4-[(E)-4-phenyl-3-buten-1-yn-1-yl]-2-cyclobuten-1-one (**137**)

Figure 71: ¹H NMR of 4,5-dimethoxy-[(E)-2-(1-iodo-3-phenyl-2-propen-1-ylidene)]-4-cyclopentene-1,3-dione (**138**)

cyclopentene-1,3-dione (**138**)

Figure 73: ¹H NMR of 4,5-dimethoxy-2-[(E)-3-phenyl-2-propen-1-ylidene]-4-cyclopente-1,3-dione (**139**)

Figure 74: ¹³C NMR of 4,5-dimethoxy-2-[(E)-3-phenyl-2-propen-1-ylidene]-4-cyclopent-1,3-dione (**139**)

Figure 75: ¹H NMR of 4,5-dimethoxy-2-[(E)-1-(N,N-dimethylamino)-3-phenyl-2-propen-1-ylidene]-4-cyclopente-1,3-dione (**140**)

Figure 76: ^1H NMR of 4-fluoro-2-nitro-1-(1-propen-1-yl)benzene (233)

Figure 77: ^{13}C NMR of 4-fluoro-2-nitro-1-(1-propen-1-yl)benzene (233)

Figure 78: ¹H NMR of 2,4-dinitro-1-(1-propen-1-yl)benzene (234)

Figure 79: ¹H NMR of 4-carbomethoxy-2-nitro-1-(1-propen-1-yl)benzene (**235**)

Figure 80: ¹H NMR of 4-methoxy-2-nitro-1-(1-propen-1-yl)benzene (236)

Figure 81: ¹³C NMR of 4-methoxy-2-nitro-1-(1-propen-1-yl)benzene (**236**)

Figure 82: ¹H NMR of 5-methoxy-2-nitro-1-(1-propen-1-yl)-benzene (**8**)

Figure 83: ^1H NMR of 3-nitrobenzaldehyde (**250**) and 1-(3-nitrophenyl)-1,2-propandione (**251**)

Figure 84: ^{13}C NMR of 3-nitrobenzaldehyde (**250**) and
1-(3-nitrophenyl)-1,2-propandione (**251**)

Figure 85: ^1H NMR of 1-(4-fluoro-2-nitrophenyl)-2-propanone (**257**)

Figure 86: ^{13}C NMR of 1-(4-fluoro-2-nitrophenyl)-2-propanone (257)

Figure 87: ^1H NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-Nitrobenzene (**258**)

Figure 88: ^{13}C NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-Nitrobenzene (258)

Figure 89: ^1H NMR of 4-fluoro-2-nitrobenzaldehyde (**259**) and 1-(4-fluoro-2-nitrophenyl)-1,2-propanedione (**260**)

Figure 90: ^{13}C NMR of 4-fluoro-2-nitrobenzaldehyde (**259**) and 1-(4-fluoro-2-nitrophenyl)-1,2-propanedione (**260**)

Figure 91: ^1H NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-Nitrobenzene (**263**)

Figure 92: ^{13}C NMR of 4-fluoro-1-(1,1-dimethoxymethyl)-2-nitrobenzene (**263**)

Figure 93: ^1H NMR of 4-carbomethoxy-2-nitrobenzaldehyde (**266**) and 1-(4-carbomethoxy-2-nitrophenyl)-1,2-propanedione (**267**)

Figure 94: ¹H NMR of 1-(5-methoxy-2-nitrophenyl)-2-propanone (269)

Figure 95: ^{13}C NMR of 1-(5-methoxy-2-nitrophenyl)-2-propanone (**269**)

Figure 96: ¹H NMR of 5-methoxy-2-nitro-benzaldehyde (**270**) and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (**271**)

Figure 97: ^{13}C NMR of 5-methoxy-2-nitro-benzaldehyde (**270**) and 1-(5-methoxy-2-nitrophenyl)-1,2-propanedione (**271**)

Figure 98: ^1H NMR of 4-(2-nitrophenyl)-3-methoxy-2-methyl-1-butene (**277**)

Figure 99: ¹³C NMR of 4-(2-nitrophenyl)-3-methoxy-2-methyl-1-butene (277)

Figure 100: ¹H NMR of 1-(2-bromophenyl)-1,2-propanedione (**282**)

Figure 101: ^{13}C NMR of 1-(2-bromophenyl)-1,2-propanedione (**282**)